

A new species of ingolfiellid amphipod (Crustacea: Amphipoda) from Western Australia

Susana Gallego Martínez and Gary C. B. Poore

Marine Biology Section, Museum Victoria, GPO Box 666E, Melbourne, Victoria 3001, Australia
emails: sgallego@museum.vic.gov.au; gpoore@museum.vic.gov.au

Abstract – A third species, *Ingolfiella quokka* sp. nov., is added to the Australian fauna, the first from Western Australia. The new species is from an intertidal sandy beach environment; previously described Australian species inhabit the marine shelf of south-eastern Australia.

INTRODUCTION

The ingolfiellidean amphipods are a small group of about 30 species included in two families, the monotypic Metaingolfiellidae, and Ingolfiellidae. Their distribution ranges from the deep sea through shallow marine sediments and intertidal sands to freshwater springs and caves. The presence of an “eye-lobe” distinguishes the suborder Ingolfiellidea from other suborders of the order Amphipoda. While it is probable that the group is monophyletic their phylogenetic isolation from Gammaridea in particular is questionable (Lowry and Poore, 1989; Dahl, 1977). Stock (1979; 1977; 1976), Dojiri and Sieg (1987), and Ruffo and Taglianti (1989) have studied the systematics and zoogeography of the group. Stock (1976) divided the family Ingolfiellidae into three genera, and the largest genus, *Ingolfiella*, into subgenera. The question of subgenera was addressed again by Ruffo and Taglianti (1989). Lowry and Poore (1989) had difficulty placing their two south-eastern Australian species unambiguously in any subgenus and questioned their usefulness. We do not add further to this debate, especially as the Ingolfiellidea as a whole are subject to a phylogenetically based reclassification (R. Vonk and F. Schram, personal communication, 2002).

A new species of ingolfiellids was discovered in samples taken on a sandy beach on Rottnest Island, Western Australia, and is here described. The material comprises 16 individuals, 2 males and 14 females. The description is a composite derived after dissection of several individuals. Types are lodged in the Western Australian Museum, Perth (WAM), and Museum Victoria, Melbourne (NMV). Abbreviations are as follows: A1, A2, antennae 1, 2; G1, G2, gnathopods 1, 2; MX1, MX2, maxillae 1, 2; MD, mandible; MP, maxilliped; P3–P7, pereopods 3–7; PL1–PL3, pleopods 1–3; r, right; l, left; UR1–UR3, uropods 1–3.

Ingolfiellidae Hansen, 1903

Ingolfiella quokka sp. nov.

Figures 1–3

Holotype

Female, 1.08 mm (WAM C33548 on 3 slides), City of York Bay (32°00'S, 115°36'E), Rottnest Island, Western Australia, sandy beach, S. Griffin, July 1991.

Paratypes

Female, 1.54 mm (WAM C33549); male, 1.25 mm (WAM C33550); female, 1.48 mm (WAM C33551, with 2 slides); female, 1.27 mm (WAM C33552); female, 1.08 mm (WAM C33553, with 1 slide); male, 1.06 mm (WAM C33554, with 2 slides); female, 0.87 mm (WAM C33555); female, 1.34 mm (WAM C33556); female, 1.38 mm (WAM C33557); female, 1.31 mm (WAM C33558); female, 0.91 mm (WAM C33559); female, 1.06 mm (WAM C33560); female, 1.26 mm (NMV J52442); female, 1.22 mm (NMV J52443); female, 0.78 mm (WAM C33561). All collected with holotype.

Description

Female (based on holotype, 1.08 mm, and paratype female, 1.48 mm). Body segments laterally compressed. Head, anterodorsal margin rounded, without rostrum; “eye-lobe” semicircular. Pereonite 1 about half as long as head; posteroventral margin oblique; deeper anteriorly than posteriorly such that pereonites 1 and 2 only weakly separated. Pereonites 2–7 increasing in depth posteriorly. Pleonites 1–3 with posteriorly rounded epimera. Urosomites 1 and 2 not markedly differentiated from pleonites, of similar length; urosomite 3 with lateral plates enclosing base of telson and uropod 3.

Antenna 1, peduncular article 1 as long as head; ratio of articles 1.0:0.4:0.3; flagellum of four articles,




Figure 1 *Ingolfiella quokka* sp. nov., holotype female, WAM C33548; Rottnest Island, Western Australia.

slightly more than half length of peduncle, article 4 with one apical aesthetasc; accessory flagellum of three articles, last minute, reaching end of article 2 of flagellum. Antenna 2, peduncle as long as peduncle of antenna 1; ratio of articles 0.4:0.2:1.0:0.8:0.8; flagellum of five articles, slightly less than half length of peduncle, article 5 with one apical aesthetasc.

Mandible molar processes acute; spine row of two club-like spines; lacinia mobilis about two-thirds width of incisor process on left, half on right, each obscurely toothed; incisor process with five blunt teeth on left and right. Maxilla 1, inner plate blunt, with one seta; outer plate with four cuspidate setae, with three, two, three, five cusps respectively; palp of two articles, with two unequal apical setae. Maxilla 2, inner and outer plates each with four terminal setae. Maxilliped basal endite with two apical setae; palp articles 1–4 with three, one, one and one mesial setae respectively, article 5 with long falcate unguis, seta at midlength and at base of unguis.

Gnathopod 1 coxa at anterior of pereonite; carpus 2.4 times as long as wide, palm with three proximal spiniform setae, three pairs of pinnate setae at mid-

palm, palm without teeth; dactylus with four teeth. Gnathopod 2 palm at 45° to longitudinal axis; carpus 1.7 times as long as wide; palm defined proximally by an obtuse angle bearing two spiniform setae (proximal one longer and more curved than other), three triangular teeth along palm, three flagellate spiniform setae and five simple setae; dactylus as long as palm, with four teeth on inner margin.

Pereopods 3 and 4, propodus with three distal setae; dactylus with two distal setae and cylindrical trifid unguis. Pereopods 5 and 6, merus with spiniform seta and simple seta; carpus with two distal long stout setae, two short stout and two slender setae; propodus with five distal setae on pereopod 6, and three setae on pereopod 5; dactylus with cylindrical bifid unguis. Pereopod 7, merus with one distal stout seta and two slender setae; carpus with one distal short stout seta, three long stout setae and three slender setae; propodus with three distal setae; dactylus stout, curved with one subtriangular spike at midlength, unguis not defined. Coxal gills ovate, small, on pereonites 3, 4, and 5. No oostegites.

Pleopods 1–3 subtriangular, without distal setae.




Figure 2 *Ingolfiella quokka* sp. nov. Paratype female, WAM C33551: **A**, left gnathopod 1; **B**, right gnathopod 2, lateral; **C**, left gnathopod 2 palm and dactylus, mesial; **D**, pereopod 3; **E**, pereopod 4; **F**, pereopod 5; **G**, pereopod 6; **H**, pereopod 7. Paratype male, WAM C33554: **I**, right gnathopod 2, palm and dactylus, mesial.

Uropod 1 peduncle with two distal setae, without a long reversed seta on lower margin; inner ramus 0.9 times length of peduncle, with lateral row of eight long setae, edges of apex finely denticulate; outer ramus half as long as inner ramus, with one subdistal seta. Uropod 2 peduncle with three obliquely transverse rows of (proximal to distal) nine, seven, nine complex setae; rami uniarticulate, outer 0.7 times length of peduncle, inner ramus 0.9 times length of outer ramus; outer ramus with three setae; inner ramus with two setae. Uropod 3 with one ramus; peduncle with

three lateral setae, ramus short, broad, with long distal seta. Telson subsemicircular, with pair of long dorsal setae.

Male (based on paratype, 1.06 mm). As female except in the following. Gnathopod 2 carpus palm defined proximally by an obtuse angle bearing one long curved spiniform seta and more distally on inner surface one strong flaring tooth-like seta; palm with three subtriangular teeth and nine simple setae. Pleopods slightly narrower than in female; pleopod 1 only with two short distal setae. Uropod 2 peduncle with a short curved seta hooked distally




Figure 3 *Ingolfiella quokka* sp. nov. Paratype female, WAM C33551: A, uropod 1; B, right uropod 2, lateral; C, right uropod 2, mesial; D, detail complex setae on mesial face of uropod 2; E, antenna 1; F, antenna 2; G, maxilla 1; H, maxilla 2; I, maxilliped; J, mandible; K, ventral view of mandibles, anterior at top, lacinia mobilis on left and right stippled. Paratype male, WAM C33554: L, right uropod 2; M, pleopods 1-3; N, urosomite 3, dorsal view with telson and uropods 3.

to meet a broad triangular marginal projection of the lower margin.

Etymology

For the Quokka (*Setonix brachyurus*), a species of wallaby living on Rottnest Island in Western Australia and after which the island was named by Willem de Vlamingh in 1696.

DISCUSSION

Ingolffiella quokka is most similar to species that have been assigned to the subgenus *Antilleella* sharing characters such as a reduced or absent ocular lobe, subtriangular pleopod 1 modified in males with two apical setae, and similar pereopods 3–7. Of the species placed in this subgenus by Ruffo and Taglianti (1989) it is similar to *I. putealis*, *I. fontinalis*, *I. tabularis*, *I. margaritae*, *I. similis*, *I. unguiculata* and *I. beatricis* as follows: (1) possession of small ocular lobe (reduced or vestigial in most species of the subgenus, except for *I. similis* where it is absent and *I. beatricis* where it is developed); (2) antenna 1 flagellum of four articles and with three-articled accessory flagellum; (3) serrated palmar margin of gnathopod 2 (variously eight teeth in *I. similis*, smooth margin in *I. beatricis*); (4) gnathopod 2 sexually dimorphic; (5) gnathopods 1 and 2 dissimilar; (6) basofacial hook on peduncle uropod 2 in males; (7) subtriangular pleopods present (subtrapezoidal in *I. beatricis* and *I. unguiculata*), modified in males with two distal setae; and (8) uropod 2 with three obliquely transverse rows of spines (except for *I. putealis* with four rows and *I. fontinalis* with two rows).

Ingolffiella quokka differs from most species of the subgenus *Antilleella* in: (1) absence of a reversed modified seta in male gnathopod 2; (2) females without oostegites (except for *I. unguiculata* and *I. beatricis*) – unless supposed females of our and these species are not mature; (3) pereopods 3 and 4 unlike pereopods 5–7, that is, dactyli of pereopods 3 and 4 with long apically bifid unguis, dactyli of pereopods 5 and 6 with stronger bifid unguis, and pereopod 7 lacks unguis; (4) absence of a sharp inner spur distally on the basal part of dactyli; and (5) dactylus of gnathopods 1 and 2 with four serrations instead of three (except for *I. beatricis*).

The new species differs from *I. bassiana*, the only Australian species which could be placed in subgenus *Antilleella*. The complexity of the palmar spines on the male gnathopod 2, the plumose seta on the peduncle of uropod 1, and the two setae of the pleopod 2 in males sets *I. bassiana* apart from *I. quokka*.

Ingolffiella quokka has also some characters similar to *I. fuscina* Dojiri and Sieg, 1987, a species more akin to the subgenus *Tethydiella* Ruffo and Taglianti, 1989. These are: (1) gnathopod 1 carpus palm

bearing Y-shaped setae; (2) gnathopod 1 dactylus with four teeth; (3) pereopods 3–4 unlike pereopods 5–7 (diagnostic of *Tethydiella*); (4) pereopods 3 and 4 dactyli with trifold unguis (unlike *Antilleella*, where the unguis is bifid); (5) absence of a sharp inner spur distally on the basal part of in pereopods 3–7 dactyli; and (6) the complex nature of the mesial spines on uropod 2. However, the new species can be distinguished from *I. fuscina* by: (1) absence of unguis in pereopod 7; (2) female probably devoid of oostegites (always present in *Tethydiella*); (3) sexual dimorphism in gnathopod 2; (4) subtriangular pleopods present in both sexes (lamelliform with one or two apical setae in *I. fuscina*); (5) male pleopod 1 subtriangular with two apical setae (only one in *I. fuscina*); (6) absence of terminal, dorsal, pectinate spine on male uropod 1 (uropod 1 sexually dimorphic in *I. fuscina*); and (7) presence of a basofacial hook on the peduncle of uropod 2 in males.

Ingolffiella quokka differs from *I. australiana* Lowry and Poore, 1989, another Australian ingolffiellid. This species shares characters with species belonging to the subgenus *Trianguliella* Stock, 1976. Both species have (1) a small semicircular eyelobe; (2) antenna 2 flagellum of five articles; (3) accessory flagellum of antenna 1 of three articles, the last one minute; (4) dactylus of gnathopod 1 with four serrations; and (5) pleopods 2 and 3 subtriangular in both sexes. However, *I. australiana* is quite different from *I. quokka*: (1) antenna 1 flagellum of five articles; (2) maxilla 2, inner and outer plates with five plumose setae; (3) gnathopod 2 palm with distal triangular tooth and quadrate tooth separated by a notch; (4) pereopods 5 and 6, dactylus with unguis not defined; (5) pleopod 1 digitiform in male with two apical setae, and subtriangular in female with one seta; (6) uropod 1 has a ventral peduncular row of stout setae (not reported in any other ingolffiellidean); and (7) uropod 2 without basofacial hook on peduncle, and with five obliquely transverse rows of setae (not complex setae).

ACKNOWLEDGEMENTS

We thank Ronald Vonk, Zoological Museum of Amsterdam, The Netherlands for comments on a first draft of this paper.

REFERENCES

- Dahl, E. (1977). The amphipod functional model and its bearing upon systematics and phylogeny. *Zoologica Scripta* 6: 221–228.
- Dojiri, M. and Sieg, J. (1987). *Ingolffiella fuscina*, new species (Crustacea: Amphipoda) from the Gulf of Mexico and the Atlantic coast of North America, and partial redescription of *I. atlantisi* Mills, 1967.

- Proceedings of the Biological Society of Washington* **100**: 494–505.
- Lowry, J.K. and Poore, G.C.B. (1989). First ingolfiellids from the southwest Pacific (Crustacea: Amphipoda) with a discussion of their systematics. *Proceedings of the Biological Society of Washington* **102**: 933–946.
- Ruffo, S. and Taglianti, A.V. (1989). Description of a new cavernicolous *Ingolfiella* species from Sardinia, with remarks on the systematics of the genus (Crustacea, Amphipoda, Ingolfiellidae). *Annali del Museo Civico di Storia Naturale Giacomo Doria, Genova* **87**: 237–261.
- Stock, J. (1976). A new member of the crustacean suborder Ingolfiellidea from Bonaire, with a review of the entire suborder. *Studies on the Fauna of Curaçao and other Caribbean Islands* **50**: 56–75.
- Stock, J. (1977). The zoogeography of the crustacean suborder Ingolfiellidea with descriptions of new West Indian taxa. *Studies on the Fauna of Curaçao and other Caribbean Islands* **55**: 131–146.
- Stock, J. (1979). New data on taxonomy and zoogeography of ingolfiellid Crustacea. *Bijdragen tot de Dierkunde* **45**: 181–190.

Manuscript received January 9 2003; accepted 18 June 2003

Guide to Authors

Subject Matter:

Reviews, observations and results of research into all branches of natural science and human studies will be considered for publication. However, emphasis is placed on studies pertaining to Western Australia. Longer papers will be considered for publication as a Supplement to the *Records of the Western Australian Museum*. Short communications should not normally exceed three typed pages and this category of paper is intended to accommodate observations, results or new records of *significance*, that otherwise might not get into the literature, or for which there is a particular urgency for publication. All material must be original and not have been published elsewhere.

Presentation:

Authors are advised to follow the layout and style in the most recent issue of the *Records of the Western Australian Museum* including headings, tables, illustrations and references.

The title should be concise, informative and contain key words necessary for retrieval by modern searching techniques. An abridged title (not exceeding 50 letter spaces) should be included for use as a running head.

An abstract must be given in full length papers but not short communications, summarizing the scope of the work and principal findings. It should normally not exceed 2% of the paper and should be suitable for reprinting in reference periodicals.

The International System of units should be used.

Numbers should be spelled out from one to nine in descriptive text; figures used for 10 or more. For associated groups, figures should be used consistently, e.g. 5 to 10, not five to 10.

Spelling should follow the *Concise Oxford Dictionary*.

Systematic papers must conform with the International Codes of Botanical and Zoological Nomenclature and, as far as possible, with their recommendations.

Synonymies should be given in the short form (taxon, author, date, page) and the full reference cited at the end of the paper. All citations, including those associated with scientific names, must be included in the references.

Manuscripts:

The original and two copies of manuscripts and figures should be submitted to the Editors, c/- Publications Department, Western Australian Museum, Francis Street, Perth, Western Australia 6000. They must be in double-spaced typescript on A4 sheets. All margins should be at least 30 mm wide. Tables plus heading and legends to illustrations should be typed on separate pages. The desired position for insertion of tables and illustrations in the text should be indicated in pencil. Tables should be numbered consecutively, have headings which make them understandable without reference to the text, and be referred to in the text.

High quality illustrations are required to size (16.8 cm x 25.2 cm) or no larger than 32 cm x 40 cm with sans serif lettering suitable for reduction to size. Photographs must be good quality black and white prints, not exceeding 16.8 cm x 25.2 cm. Scale must be indicated on illustrations. All maps, line drawings, photographs and graphs, should be numbered in sequence and referred to as Figure/s in the text and captions. Each must have a brief, fully explanatory caption. On acceptance a computer disk containing all corrections should be sent with amended manuscript. The disk should be marked with program (e.g. Word, WordPerfect, etc).

In papers dealing with historical subjects references may be cited as footnotes. In all other papers references must be cited in the text by author and date and all must be listed alphabetically at the end of the paper. The names of journals are to be given in full.

Processing:

Papers and short communications are reviewed by at least two referees and acceptance or rejection is then decided by the editors.

The senior author is sent one set of page proofs which must be returned promptly.

The senior author will receive fifty free offprints of the paper. Additional offprints can be ordered at page proof stage.

Records of the Western Australian Museum

Volume 22 Part 1 2003

CONTENTS

N. Warburton, C. Wood, C. Lloyd, S. Song and P. Withers The 3-dimensional anatomy of the North-Western Marsupial Mole (<i>Notoryctes caurinus</i> Thomas 1920) using computed tomography, X-ray and magnetic resonance imaging	1
F.E. Wells and H. Morrison Description of <i>Volutoconus hargreavesi calcarelliformis</i> subsp. nov. (Mollusca: Volutidae) from northwestern Australia	9
R.L. Hoffman A new genus and species of trioniuline milliped from Western Australia (Spirobolida: Pachybolidae: Trioniulinae)	17
I. Bartsch Psammophilous halacarids (Halacaridae, Acari) from Dampier, Western Australia. Description of species and faunal comparison of the mesopsammal halacarid fauna of western and eastern Australia	23
I. Maryanto Taxonomic status of the ricefield rat <i>Rattus argentiventer</i> (Robinson and Kloss, 1916) (Rodentia) from Thailand, Malaysia and Indonesia based on morphological variation	47
G.C.B. Poore and W.F. Humphreys Second species of <i>Mangkurtu</i> (Spelaeogriphacea) from north-western Australia	67
S. Gallego Martínez and G.C.B. Poore A new species of ingolfiellid amphipod (Crustacea: Amphipoda) from Western Australia	75

Dates of Publication

Records of the Western Australian Museum

Volume 21, Part 1	1 May 2002
Volume 21, Part 2	18 June 2002
Volume 21, Part 3	20 November 2002
Volume 21, Part 4	27 June 2003