

## *Lohmannella* and *Simognathus* (Halacaridae: Acari) from Western Australia: description of two new species and reflections on the distribution of these genera

Ilse Bartsch

Deutsches Zentrum für Marine Biodiversitätsforschung, Forschungsinstitut Senckenberg, Notkestr. 85, 22607 Hamburg, Germany, e-mail: bartsch@meeresforschung.de

**Abstract** – Two new littoral species, *Lohmannella septemsetosa* sp. nov. and *Simognathus cruciferus* sp. nov. are described and the description of *L. pinggi* Otto, 1994 is supplemented. The genera *Lohmannella* and *Simognathus* are distributed world-wide. More species are recorded from the southern than northern hemisphere. *Lohmannella* is abundant in warm-temperate to polar regions but rare in the tropics. In contrast, most records of *Simognathus* are from tropical and warm-temperate areas. According to external morphological characters and distribution, different lineages of *Lohmannella* species developed in the north and south. *Simognathus* demonstrates no such north-south discrimination.

### INTRODUCTION

The two genera *Lohmannella* and *Simognathus* include 35 and 43 species and subspecies, respectively (to the end of 2004). Representatives of both are distributed world-wide, but there is a distinct difference between the faunas of the northern and southern hemisphere. *Simognathus* is more diverse in the south than in the north (Bartsch 1994, 2004). More marine *Lohmannella* species are recorded from the southern than northern hemisphere, though locally species diversity is high in the north, too. Both genera are present in Western Australia. One *Lohmannella* species has been described from Rottnest Island, off Perth (Bartsch 1993b). A record of another species and the description of a new species, from Esperance, will be presented in this paper. Ten *Simognathus* species are already known to belong to the Western Australian fauna (Bartsch 1993b, 1994, 2003b), and a new species from Esperance is described below.

### MATERIAL AND METHODS

The material studied was collected during Marine Biological Workshops held in Western Australia, on Rottnest Island and in Esperance. The fauna and flora around Rottnest Island is dominated by warm-temperate species, although strongly influenced by tropical elements (Wells and Walker 1993). Esperance lies within the warm-temperate zone (Knox 1963; Wilson and Allen 1987; Ponder and Wells 1998; O'Hara and Poore 2000).

The material was collected by the author. The mites were cleared in lactic acid and mounted in

glycerine jelly. Slides with holotypes and voucher specimens are deposited in the Western Australian Museum, Perth (WAM), apart from one *Lohmannella* specimen in the Zoological Institute and Museum in Hamburg (ZMH).

Abbreviations used in the descriptions are as follows: AD, anterior dorsal plate; AE, anterior epimeral plate; ds-1 to ds-5, first to fifth pair of dorsal setae numbered from anterior backward; GA, genitoanal plate; glp-1 to glp-5, pair of gland pores numbered 1 to 5 from anterior backward; GO, genital opening; GP, genital plate; OC, ocular plate(s); P-2 to P-4, second to fourth palpal segment; pas, parambulacral seta(e); PD, posterior dorsal plate; PE, posterior epimeral plate(s); pgs, perigenital setae; sgs, subgenital setae. The legs, their segments and claws are numbered I to IV. The leg segments 1 to 6 are trochanter, basifemur, telofemur, genu, tibia, and tarsus. The setation formula of the legs is presenting the number of setae from trochanter to tarsus.

The position of a seta is given in a decimal system, with reference to the length from anterior to posterior or basal to distal.


### SYSTEMATICS

Family Halacaridae Murray, 1877

Subfamily Lohmannellinae Viets, 1927

Genus *Lohmannella* Trouessart, 1901

*Lohmannella septemsetosa* sp. nov.  
Figures 1A–K, 2A–F



**Figure 1** A–K. *Lohmannella septemsetosa* sp. nov., holotype male. A, Idiosoma, dorsal; B, idiosoma, ventral; C, part of OC with gland pore; D, genital opening; E, P-3 and P-4, lateral; F, gnathosoma, lateral (stippled area enclosing maze-like structured integument); G, gnathosoma, ventral; H, leg I, medial; I, ambulacrum of tarsus I, ventral; J, ambulacrum of tarsus II, medial; K, telofemur to tarsus II, medial. (ma, area with maze-like epicuticula). A, B, D–H, K: scale = 50  $\mu\text{m}$ ; C, I, J: scale = 10  $\mu\text{m}$ .

#### Material Examined

##### Holotype

Male, Lucky Bay, Cape le Grand, 33°59'S, 122°13'E, near Esperance, Western Australia, Australia; algal tufts with sediment, 30 cm below water edge, 15 February 2003 (WAM T62841).

##### Paratype

One deutonymph, same data as for holotype (WAM T62842).

#### Diagnosis

Idiosomal length 410  $\mu\text{m}$ , gnathosomal length 284  $\mu\text{m}$ , ratio idiosoma:gnathosoma 0.69. Surface of plates with reticulate and maze-like ornamented epicuticula. Gland pores large. Pair of ds-1 posterior to level of gland pores. OC with gland pore in middle of plate. P-2 without spiniform ventral process. Genu and tibia I with four and seven bipectinate ventral setae. Tarsi I to IV with 2, 2, 3, 2 bipectinate ventral setae. Claws with accessory process.

## Description

## Male

*Idiosoma*. Length 410  $\mu\text{m}$ , width 302  $\mu\text{m}$ . Dorsal plates covered by delicately reticulate or maze-like epicuticula. Length of AD 142  $\mu\text{m}$ , width 159  $\mu\text{m}$ ; pair of gland pores with wide, oblong alveolus, almost 10  $\mu\text{m}$  in width (Figure 1C). Length of OC 102  $\mu\text{m}$ , width 57  $\mu\text{m}$ . Gland pore in middle of plate, pore canaliculus at the same level but in lateral margin (Figure 1A). A very faint, cornea-like structure near anterior margin. Length of PD 202  $\mu\text{m}$ , width 150  $\mu\text{m}$ , anterior margin truncate, with broadly rounded corners. With two pairs of large gland pores as illustrated. Pair of ds-1 (setae broken) posterior to the level of pair of gland pores. Following pairs of setae small; ds-2, ds-3 and ds-4 within striated integument, ds-5 immediately anterior to glp-4. Adanal setae on anal cone.

Ventral plates with maze-like or irregularly

reticulate or foveate epicuticula. Length of AE 137  $\mu\text{m}$ , width 269  $\mu\text{m}$ . Pair of platelets between AE and PE, each platelet with large gland pore (Figure 1B). Length of PE 179  $\mu\text{m}$ ; each plate with two dorsal and three ventral setae. Length of GA 207  $\mu\text{m}$ , width 144  $\mu\text{m}$ ; anterior margin truncate. Plate with about nine outlying setae and 52 setae close around GO (Figure 1D). Length of GO 63  $\mu\text{m}$ , width 43  $\mu\text{m}$ ; distance between anterior margin of GO and that of GA equalling 1.5 times length of GO. Genital sclerites with four pairs of spurlike sgs. Three pairs of internal genital acetabula shining through genital sclerites.

*Gnathosoma*. Slender, length 284  $\mu\text{m}$ , or 0.69 times of idiosomal length. Width of gnathosoma 107  $\mu\text{m}$ , length:width ratio 1:0.27. Lateral flank of gnathosomal base with an area with maze-like ornamented epicuticula (Figure 1F). Pharyngeal plate far from reaching posterior margin of gnathosoma (Figure 1G). Rostrum slender, much




Figure 2 A-F. *Lohmannella septemsetosa* sp. nov. A, Leg III, medial, male; B, leg IV, medial, male; C, tarsus I, lateral, male (medial setae and claw in broken line); D, tarsus II, medial, male (lateral setae and claw in broken line); E, idiosoma, dorsal, deutonymph; F, idiosoma, ventral, deutonymph. Scale = 50  $\mu\text{m}$ .

longer than gnathosomal base. Both pairs of maxillary setae short and situated in posterior quarter of rostrum. Palps slender. P-2 with two setae, basal setae near segment's base, at 0.2. P-3 with spur-like process (no spine). Basal portion of P-4 with two slender setae and one spur, apical portion with solenidion, small seta and spur-like tip (Figure 1E).

*Legs.* Slender, integument with maze-like sculpturing. Leg I slightly longer, leg IV shorter than idiosoma. Telofemur I about 3.4 times longer than high (Figure 1H). Leg chaetotaxy (pas excluded, solenidia included): leg I, 1, 3, 9, 9, 14, 11 (Figure 1H); leg II, 1, 3, 9, 9, 13, 9 (Figure 1K); leg III, 1, 3, 6, 5, 10, 8 (Figure 2A); leg IV, 1, 3, 4, 5, 10, 6 (Figure 2B). Several of short dorsal setae delicately serrate. Genua I to IV with 4, 4, 2, 2 ventral setae; setae of genua I, II and IV bipectinate. Tibiae I to IV with 7, 6, 5, 5 ventral setae; 7, 4, 2, 3 of these setae bipectinate. Tarsi I to IV with 4, 4, 4, 3 dorsal setae, solenidia included. Solenidion of tarsus I 11  $\mu\text{m}$  in length and in dorsolateral position (Figure 2C); on tarsus II solenidion 14  $\mu\text{m}$  in length and in dorsal position (Figure 2D). Tarsi I to IV with 2, 2, 3, 2 bipectinate ventral setae and 5, 3, 1, 1 slender setae. Tarsus I with pair of doubled pas; tarsus II with medial pas doubled, lateral pas single; tarsus III with spiniform lateral pas and a seti- plus a spiniform medial pas; pair of pas of tarsus IV spiniform.

Claws with accessory process. Central sclerite small, without claw-like process. All tarsi with carpite between end of tarsi and claws (Figure 1I and J). Carpite divaricate, 9  $\mu\text{m}$  in length.

#### *Deutonymph*

Idiosomal length 290  $\mu\text{m}$ . Shape of AD similar to that of male. OC shorter; pair of gland pores close to posterior corner of plate (Figure 2E). PD shorter and more narrow than that plate of male. AE with four pairs of setae (Figure 2F); PE with one dorsal and three ventral setae. GP and anal plate separated. GP with three pairs of pgs and two pairs of minute, seta-like sgs. Length of gnathosoma 116  $\mu\text{m}$ , i.e. 0.80 of idiosomal length. Leg chaetotaxy: leg I, 1, 3, 5-6, 5-6, 10, 9; leg II, 1, 3, 5-6, 6, 9, 7; leg III, 1, 3, 4, 4, 7, 6; leg IV, 1, 1, 2, 3, 6, 4, 4. Genu I with pair of bipectinate setae. On genu II ventromedial seta bipectinate, ventrolateral seta slender and very faintly pectinate. Ventral seta of genu III slender, that seta of genu IV bipectinate. Tibia I with two ventromedial and three ventrolateral setae, these five setae bipectinate. Tibia II with two pairs of bipectinate setae. Tibia III with one large bipectinate and one short, slightly pectinate seta, and tibia IV with two to three strong, bipectinate setae. Tarsi I-IV with 2, 2, 2, 1 bipectinate ventral setae and 3, 1, 0, 0 eupathid ventral setae.

#### **Etymology**

The specific name is derived from *septem* (Latin), seven, and *setosus* (Latin), with setae, as tibia I of this species bears seven ventral setae.

#### **Remarks**

The most conspicuous characters of *Lohmannella septemsetosa* are the gland pores with large alveoli and the seven bipectinate ventral setae on tibia I. A similar combination of characters is present in *L. dictyota* Bartsch, 1992, *L. gaussi* Lohmann, 1907, *L. kerguelensis* Lohmann, 1907, and *L. pinggi* Otto, 1994. Records of *L. dictyota* are from the southwestern Pacific, from the Society Islands, the Coral Sea and Great Barrier Reef (Bartsch 1992; Otto 2000). *L. pinggi* is known from Victoria and New South Wales, Australia (Otto 1994), *L. gaussi* from Antarctica and sub-Antarctica, from off Wilhelm II Land, the islands Crozet, Marion and South Sandwich (Lohmann 1907; Bartsch 1979a, 1993a; Newell 1984), and *L. kerguelensis* from the Kerguelen Islands and Palmer Archipelago (Lohmann 1907; Bartsch 1993a). *Lohmannella dictyota* has an unusual elongate PD with a prominent ornamentation, the ds-3 are situated on the PD, the pair of gland pores 2 are in the medial corners of the OC, and the P-2 has a spiniform ventral process. *Lohmannella gaussi* is, compared with congeners, large-sized, its legs and gnathosoma are long and slender; the glp-2 are near the lateral margin of the OC, and the PD bears a V-shaped porose area, characters not present in *L. septemsetosa*. In contrast to *L. septemsetosa* the OC of *L. kerguelensis* has a cornea and the ds-3 are in the anterior rounded margin of PD. *Lohmannella pinggi* has much shorter legs than *L. septemsetosa*, elongate OC and the ds-1 are situated distinctly anterior to the level of gland pore 1.

#### *Lohmannella pinggi* Otto, 1994

Figure 3A-G


*Lohmannella pinggi* Otto, 1994: 32-35, figures 1-8.

#### **Material Examined**

One female, Rottneest Island, Cape Vlamingh, ca 32°02'S, 115°27'E, Western Australia, Australia; from corallines on rocky platform, 10 January 1991 (WAM T62843). One female, Rottneest Island, Cape Vlamingh; from corallines on rocky platform, 9 January 1991 (WAM T62844). One female (damaged), Rottneest Island, Cape Vlamingh, Western Australia, Australia; from corallines on rocky platform, 9 January 1991 (ZMH).

#### **Diagnosis**

Idiosomal length 279-303  $\mu\text{m}$ , gnathosomal length 192-197  $\mu\text{m}$ . Dorsal plates delicately reticulated. OC oblong, its length more than twice the width. With five pairs of large gland pores; glp-



**Figure 3** A–G. *Lohmannella pinggi* Otto, 1984, female. A, Idiosoma, dorsal; B, idiosoma, ventral; C, leg I, medial; D, gnathosoma, dorsal; E, tip of tarsus II, medial; F, idiosoma, dorsal, G, gnathosoma, lateral. Scale = 50  $\mu$ m.

3 near medial margin of OC. Pair of ds-1 anterior to level of gland pores. Female GA with nine pairs of pgs. Gnathosoma 2.4 times longer than wide, length ratio idiosoma to gnathosoma 1:0.65. No ventral spiniform process on P-2. Basal seta of P-2 strong, situated in basal half. Genu I with two pairs of bipectinate setae; tibia I with seven such setae. Tarsi I to IV with 2, 2, 4, 2 bipectinate ventral setae. Accessory process of claws with minute tines.

#### Supplementary Description

Idiosomal length 279 and 303  $\mu$ m, gnathosomal length 192 and 197  $\mu$ m. Dorsal plates delicately reticulated and with scattered deep canaliculi (Figure 3A). OC oblong, length more than twice its width; anterior cornea distinct, posterior one faint. Gland pores distinct, their alveoli 7  $\mu$ m in width; glp-2 in ventral position, glp-3 on OC near

its medial margin, at about 0.57. Pair of ds-1 anterior to the level of gland pores. Two females (WAM T62843 and ZMH) with ds-2, ds-3 and ds-4 within striated integument (Figure 3A); ds-5 adjacent to glp-4. One female (WAM T62844) with pair of ds-4 situated on PD (Figure 3F). PE long, extending anteriorly beyond middle of OC. Female with 18 pgs. Interval between anterior margin of GA and GO equalling 1.1 times the length of the latter. Gnathosomal length 192 and 197  $\mu$ m, i.e., 0.65–0.69 of idiosomal length. P-2 without ventral process (Figure 3G); its basal seta situated at 0.22 (Figure 3D). Legs short. Genua and tibiae with articular membranes; tarsi with small fossa membranes. Length:height ratio of telofemora I and II 2.0, that of telofemora III and IV 1.6. Leg chaetotaxy (solenidia included, pas excluded): leg I, 1, 3, 8, 8, 13, 10; leg II, 1, 3, 8, 8, 12, 9; leg III, 1, 3,

6, 4–5, 9, 8; leg IV, 1, 3, 4, 4, 8, 5–6. Number of bipectinate ventral setae of genua I to IV: 4, 2, 0, 1; of tibia I to IV: 7, 3, 1, 4 (one of the latter slender); of tarsi I to IV: 2, 2, 4, 2. Tarsus I with four dorsal setae (solenidion included), two bipectinate ventral setae, two pairs of ventral eupathia and a pair of pas (Figure 3E); lateral pas doubled. Tarsus II with four dorsal setae, three ventral eupathidia and pair of pas; lateral pas doubled. Tarsus III with four dorsal setae and pair of pas, lateral pas setiform, medial pas including both a seti- and spiniform seta. Tarsus IV with three dorsal setae, none or one smooth ventral seta and pair of spiniform pas. Accessory process of claws with two tines (Figure 3E).

### Remarks

Three females were taken amongst shallow water corallines, of which two agree almost perfectly with the description of *L. pinggi* by Otto (1994). Differences are: the somewhat smaller size (303  $\mu\text{m}$  vs 378–437  $\mu\text{m}$ ), shape of ds-1 (long and slender vs spiniform), number of corneae (two vs one), presence of the ds-5 (vs absence), number of setae on AE (four vs three pairs) and number of dorsal setae of the legs. The five last mentioned differences are thought to be due to either a range of great variability in the type series or inadequate description; the difference in size may partly be due to the mounting.

One of the present females has the ds-4 inserted on the PD instead, as usual, within the striated integument posterior to the OC. Apart from this character, that female agrees with the other specimens. In most of the *Lohmannella* species the ds-2, ds-3 and ds-4 are situated on tiny sclerites within the striated integument; intraspecific variations, if present at all, are expected to be within a very limited range. The ds-2 in general are between the AD and OC, in some few species the ds-2 are in or on the lateral margin of the AD. The ds-3 are anterior to the PD, in ten species the setae are in the margin or on the PD; the ds-4 are inserted immediately posterior to the OC; one exception is *L. multispina* Newell, 1984 where the setae are found in the corner of the OC, other exceptions are the freshwater species, i.e., *L. andrei* (Angelier, 1951), *L. heptapegoni* Petrova, 1966, *L. curvimandibulata* (Petrova, 1969), *L. cvetkovi* (Petrova, 1965), and *L. stammeri* Viets, 1939. In these species both the ds-4 and ds-5 are inserted on the PD. These five species live in nearshore and continental waters in countries around the Mediterranean (Bartsch 1996; Pesic 2004).

### Subfamily Simognathinae Viets, 1927

#### Genus *Simognathus* Trouessart, 1889

### *Simognathus cruciferus* sp. nov.

Figure 4A–L

### Material Examined

#### *Holotype*

Female, Duke of Orleans Bay, ca 33°55'S, 122°35'E, near Esperance, Western Australia, Australia; demosponge overgrown with green algae and corallines, just below water line, 17 February 2003 (WAM T62845).

### Diagnosis


Idiosomal length 375  $\mu\text{m}$ . Dorsal plates colourless, almost uniformly foveate and with numerous delicate canaliculi. AD and OC with cornea-like structures. OC almost triangular. Pair of ds-4 on PD. Adanal setae in ventral position. AE with marginal areolae with foveae and canaliculi, integument in a cross-shaped median area almost smooth. GA with such smooth integument in a T-shaped area. Anterior margin of female GA arched. P-2 with ventral protuberance and bristle. Tibia I with wide but short, bluntly ending spine.

### Description

#### *Female*

*Idiosoma*. Length 375  $\mu\text{m}$ , width 200  $\mu\text{m}$ . Dorsal plates almost uniformly foveate (Figure 4A), each fovea surrounded by delicate canaliculi. All plates colourless. Length of AD 157  $\mu\text{m}$ , width 97  $\mu\text{m}$ ; its posterior margin truncate. First pair of gland pores small, slightly posterior to the level of insertion of leg I. An ovate smooth area near anterior margin. Length of OC 40  $\mu\text{m}$ , width 24  $\mu\text{m}$ , with ovate cornea. Length of PD 165  $\mu\text{m}$ , width 95  $\mu\text{m}$ , 1.7 times longer than wide. Dorsal setae small. Pair of ds-1 on AD close to pair of gland pores. Pair of ds-2 within striated integument immediately anterior to OC. Pair of ds-3 in distolateral corners of AD; ds-4 and ds-5 on PD; adanal setae on anal cone, in ventral position.

AE marginally foveate but smooth immediately posterior to insertion of legs I and II and in a ventral cross-shaped area (Figure 4B). Foveae surrounded by delicate canaliculi. Two pairs of internal scars (muscle attachment) in middle of cruciform area. Length of AE 142  $\mu\text{m}$ , width 199  $\mu\text{m}$ ; epimeral processes I large, slightly raised, epimeral vesicles large, with three pairs of ventral setae, posteriormost pair only slightly posterior to second pair of setae. Length of PE 177  $\mu\text{m}$ . Length of GA 155  $\mu\text{m}$ , width 108  $\mu\text{m}$ , integument in a T-shaped area almost smooth, remainder foveate. Length of GO 43  $\mu\text{m}$ , width 25  $\mu\text{m}$ ; distance between anterior margin of GO and that of GA 1.7 times length of GO. With four pairs of pgs on either side of GO.



**Figure 4** A–L. *Simognathus cruciferus* sp. nov., holotype female. A, Idiosoma, dorsal; B, idiosoma, ventral; C, gnathosoma, dorsal; D, gnathosoma, lateral; E, leg I, medial; F, leg II, medial; G, leg III, lateral; H, leg IV, lateral; I, tarsus and tibia I, lateral (medial parambulacral seta and claw omitted, other medial setae in broken line); J, tip of tarsus I, lateral (parambulacral seta, medial claw and fossary seta omitted); K, tarsus and tibia II, medial (lateral setae omitted); L, leg IV, medial. Scale = 50  $\mu\text{m}$ .

*Gnathosoma*. Length 97  $\mu\text{m}$ , width 85  $\mu\text{m}$ . Integument of gnathosomal base foveate. Rostrum short. Basal pair of maxillary setae near base of rostrum, apical pair of setae shorter, in posterior third of rostrum (Figure 4D). Tip of rostrum with

two pairs of rostral setae. Palps three-segmented; inserted adjacent. In dorsal aspect tectum with median keel, else scale-like and partly obscuring palpal base (Figure 4C). P-2 with ventral protuberance and one bristle-like seta (Figure 4D).





**Table 1** The genus *Lohmannella*. List of marine species, morphological characters and collecting data.

species	I	II	III	IV	V	VI	VI	VIII	IX
<b>Northern Hemisphere</b>									
<i>beringi</i> Newell, 1951	?	?	?	?	?	71N, 160W 52–57N, 173E–170W	Arctic Alaska	0–39	algae, sand
<i>falcata</i> (Hodge, 1863)	a	0	4	6	2	52–55N, 158–166E 47–50N, 152–56E 45–65N, 21W–21E 41–45N, 67–71W 42–44N, 3–16E 42–46N, 28–38E	Kamchatka Kuril Is N Europe NE America Mediterranean Black Sea	0–300	colonial organisms, algae, sediment,
<i>kervillei</i> (Trouessart, 1894)	a	0	2	6	1	38–53N, 28W–1E	N Europe	tidal	corallines, sand
<i>multisetosa</i> Bartsch, 1986	a	0	2	6	1	43N, 5E 49–58N, 4W–11E,	Mediterranean N Europe	10–45	sediment
<i>norvegica</i> Viets, 1927	a	0	1	6	1	50–60N, 5W–4E,	N Europe	0–sublittoral	
<i>nudipes</i> Bartsch, 1977	a	0	1	5	0	49N, 4W	N Europe	tidal	sediment
<i>rustica</i> Bartsch, 1977	a	0	2	5	1	49N, 4W	N Europe	tidal	crustose corallines
<i>steueri steueri</i> Viets, 1940	a	0	3	5	1–2	45N, 14E	Mediterranean	25–28	shell rubble, algae
<i>steueri reducta</i> Bartsch, 1986	a	0	3	4	0–2	43N, 6E	Mediterranean	20–40	sediment
<i>subfalcata</i> Bartsch, 2003	a	0	3	6	2	30N, 28W	NE Atlantic	318–321	
<b>Southern Hemisphere</b>									
<i>africana</i> Bartsch, 1992	+	1	4	6	1	33S, 28E	South Africa	0	shell rubble
<i>antarctica</i> Newell, 1984	+	0	4	6	2	64S, 64W 66S, 136E 77S, 166E	Palmer Peninsula Terre Adélie Ross Island	6–460	algae, bryozoans
<i>arenaria</i> Bartsch, 1993	+	2	4	6	2	32S, 115E	SW Australia	tidal	coarse sand
<i>bihamata</i> Viets, 1950	+	0	1	6	2	52–54S, 37–58W	Falkland, S. Georgia	shallows	sand, rubble, algae
<i>bispina</i> Newell, 1984	+	2	4	6	1	53S, 72W	Chile	tidal	coarse sand
<i>consimilis</i> Bartsch, 1993	+	0	4	6	2	57S, 27W	S. Sandwich	93–121	
<i>cygna</i> Bartsch, 1988	+	0	5	8	2	9S, 12E	Angola Basin	1430	
<i>dictyota</i> Bartsch, 1992	+	1	2	7–8	2	16S, 152W 15–19S, 145–151E	Society Islands NE Australia	tidal–17	coral rubble, coarse sand
<i>fukushimai</i> Imamura, 1968	(+)s	0	4	7	2	69S, 31E 72S, 172E 71–74S, 12–29W	Prince Harald Coast Ross Sea Weddell Sea	190–820	
<i>gaussi</i> Lohmann, 1907	+	0	4	7	2	65S 92E 56S, 27W 47S, 38E	Wilhelm II Land S. Sandwich Marion Island	95–385	

<i>grandipora</i> Newell, 1984	+	1-2	5	6	1	28-54S, 71-72W 54S, 37W	Chile S. Georgia	tidal	coarse shell, sand, holdfasts
<i>hureaui</i> Newell, 1984	+	0	3	6	?	49S, 70W	Kerguelen	shallows	holdfasts
<i>kerguelensis</i> Lohmann, 1907	+	0	4	7	2	50S, 70E 64S, 61W	Kerguelen Palmer Peninsula	shallow-58	
<i>lamellipes</i> Newell, 1984	+	4-5	?	6	2	53S, 72W	Chile	shallows	holdfasts with sand
<i>masatierrae</i> Newell, 1984	+	0	?	6	?	34S, 79W	Chile	shallows	red algae
<i>multispina</i> Newell, 1984	+	3-4	3	5-6	1-2	18S, 70W	Chile	tidal	shell rubble
<i>pinggi</i> Otto, 1994	+	0	4	7	2	35-38S, 145-150E 32S, 115E	S and SE Australia SW Australia	tidal tidal	corallines
<i>rectangulops</i> Newell, 1984	+	0	2	5	1	33S, 72W	Chile	tidal	corallines, bivalves
<i>reticulata</i> Viets, 1950	+	0	3-4	6	1	52S, 58W	Falklands	12-16	sand, rubble, algae
<i>septemsetosa</i> sp. nov.	+	0	4	7	2	34S, 122E	SW Australia	low water	sand with algae
<i>setosa</i> Newell, 1984	+	0	3	6	2	26S, 80W	Chile	tidal	corallines

- I. Number of gland pores: a, one or more pairs of pores lacking; +, five pairs present; (+)s, one or more of the five pairs replaced by setae.  
 II. P-2, number of spiniform processes:  
 III. I-4, number of bipectinate setae:  
 IV. I-5, number of bipectinate setae:  
 V. I-6, number of bipectinate setae:  
 VI. Collecting area, coordinates:  
 VII. Collecting area, state/province/island or sea area:  
 VIII. Depth (in m):  
 IX. Habitat:

**Table 2** Freshwater *Lohmannella* species and collecting data.

species	country	salinity
<i>andrei</i> (Angelier, 1951)	France	fresh
<i>curvimandibulata</i> (Petrova, 1966)	Bulgaria	fresh
<i>cvetkovi</i> (Petrova, 1965)	Bulgaria	fresh
<i>heptapegoni</i> Petrova, 1966	Israel	fresh
<i>stammeri</i> Viets, 1939	Italy, Monte Negro	fresh, slightly brackish

**Table 3** The genus *Simognathus*. List of species, morphological characters and collecting data.

species	I	II	III	IV	V	VI	VI	VIII	IX
<b>Northern Hemisphere (5–90N)</b>									
<i>adriaticus</i> Viets, 1940	la	p+s	sl	?	?	45N, 14E	Mediterranean	0.5–3	algae
<i>coreensis</i> Chatterjee and Chang, 2004	la	p/s	sl	1+1?	1+1?	36N, 135E	Korea		intertidal corallines
<i>foveolatus</i> Bartsch, 1991	la	p/s	sl	1+1	1+1	22N, 115E	S China	low water	coarse sand
<i>fuscus</i> Viets, 1936	la	p/s	sl	1+1	1+1	12–32N, 65–68W	Bermuda, Caribbean	0–11	rubble, algae
<i>leiomerus</i> Trouessart, 1894	re	p+s	sl	1+2	2+2	49–54N, 2–6W	N Europe	0–13	sediment, algae
						43N, 3–5E	Mediterranean		
<i>minor</i> Bartsch, 1979	re	p+s	sl	1+2	2+2	41N, 71W	E USA	tidal	sand
<i>minutus</i> (Hodge, 1863)	la	p/s	sl	1+1	1+1	47–60N, 10W–12E*	N Europe	tidal–65	sand, algae
<i>serratus</i> Bartsch, 2004	la	p/s	sl	1+1	1+1	29–30N, 28W	Great Meteor Bank	476–511	
<b>Equator (5N–5S)</b>									
<i>disparilis</i> Bartsch, 1977	re	p/s	sl	1+2	1+2	0, 90W	Galapagos	tidal	sand
<i>similis</i> Bartsch, 1977	la	p+s	sl	1+1	1+1	0, 90W	Galapagos	tidal	sand
<i>tropicalis</i> Chatterjee and de Troch, 2000	re	p/s	?	1+?	1+?	4S, 20E	Kenya	low water	seagrass
<b>Southern Hemisphere (5–80S)</b>									
<i>abnormalus</i> Otto, 2000	re	s	sl	1+1	1+1	16–19S, 145–150E	NE Australia	0–15	coarse sand, rubble
<i>actius</i> Otto, 2000	re	p/s	pe	1+1	1+1	16–19S, 145–147E	NE Australia	tidal	sand
<i>areolatus</i> Newell, 1984	la	p/s	sl	?	?	18–53S, 70–72W	Chile	tidal	sand, colonial organisms, algae, holdfasts
<i>aspidiotus</i> Otto, 2000	re	p/s-p	sl	1+1	1+2	15–19S, 145–152E	NE Australia	0–15	coarse sand, coral rubble
<i>clypeatus</i> Otto, 2000	re	p/s	?	1+2	1+2(d)	17S, 149W	NE Australia	5–15	coarse sand
<i>corneatus</i> Otto, 2000	la	p/s	sl	1+1	1+1	19S, 147–149E	NE Australia	3–15	coarse sand, rubble
<i>coutieri</i> (Trouessart, 1899)	re	p/s	?	?	?	12S, 43E	Djibouti	low water	shell and coral rubble
<i>cruciferus</i> sp. nov.	la	p/s	sl	1+1	1+1	34S, 122E	SW Australia	low water	demosponge with small algae
<i>delicatulus</i> Bartsch, 1994	re	p/s	sl	1+1	1+1	32S, 115E	SW Australia	tidal	sand
<i>exoticus</i> Otto, 2000	la	p/s	sl	1+1	1+1	17–18S, 148–152E	NE Australia	5–7	sand
<i>euphractus</i> Pepato and Tiago, 2004	la	p+s	sl?	1+1	1+1	24S, 45W	Brazil	tidal	rocky shore
<i>fuscus</i> Viets, 1936	la	p/s	sl	1+1	1+1	24S, 45W	Brazil	tidal	rocky shore
<i>gibberosus</i> Bartsch, 1994	la	s	sl	1+1	1+1	32S, 115E	SW Australia	low water	sand, corallines
<i>glaber</i> Bartsch, 1986	la	p/s	sl	1+2	?	36S, 175E	New Zealand	tidal	corallines
<i>glareus</i> Bartsch, 1986	la	p/s	sl	1+1	1+1	36S, 175E	New Zealand	tidal	gravel
						35S, 19E	South Africa		

<i>gracilis</i> Bartsch, 1994	re	s	sl	1+1	1+1	32S, 115E	SW Australia	30	medium to coarse sand
<i>hulingsi</i> Newell, 1984	la	p+s?	sl	?	?	18-53S, 70-72W	Chile	tidal-shallows	worm tubes, holdfasts
<i>latitarsus</i> Proches, 2002	la	p+s?	w	1+2	1+2?	30S, 30E	South Africa	tidal	algae
<i>maculatus</i> Bartsch, 1994	la	p/s	sl	1+1	1+1	32S, 115E	SW Australia	0.5	seagrasses
<i>magellanicus</i> Newell, 1984	la	p/s	sl	?	?	53-55S, 71W	Chile	tidal-278	algae, corallines, holdfasts, barnacles
						52S, 58W	Falklands		
						54S, 159W		88-278	
						55S, 159E	Macquarie Ridge	92-113	
<i>obtusus</i> Newell, 1971	la	p/s	sl	?	?	26-33S, 72-80W	Chile	tidal-190	corallines, mussels, coarse sand
<i>pectinatus</i> Newell, 1984	re	p+s?	sl	?	?	17-27S, 70-71W	Chile	tidal	shell fragments, worm tubes
<i>platyaspis</i> Otto, 2000	la	p/s	sl	1+1	1+1	15-28S, 145-147E	NE Australia	0-6	sand, rubble, coral fragments
						21S, 117E	NW Australia	low water	coral block
						10S, 124E	Philippines	10-15	sand and algae
<i>pygmaeus</i> Otto, 2000	re	s	sl	1+2	1+2	16S, 145E	NE Australia	tidal	sand
<i>salebrosus</i> Bartsch, 2003	re	s	sl	1+1	1+1	21S, 117E	NW Australia	tidal	sand
<i>scutatus</i> Bartsch, 1993	re	s	sl	1+1	1+1	32S, 115E	SW Australia	tidal	coarse sand
<i>specialis</i> Otto, 2000	la	p/s	sl	1+1	1+1	15-18S, 147-150E	NE Australia	0.5-15	algae, coarse sand, coral rubble
<i>subobtusus</i> Newell, 1984	la	p/s	sl	?	?	34S, 79W	Chile	tidal	algae
<i>tener</i> Bartsch, 2003	re	p/s	sl	1+2	1+2	21S, 117E	NW Australia	tidal	coarse sand
<i>trachys</i> Otto, 2000	la	s	sp	1+1	1+1	19S, 149E	NE Australia	10	coarse sand, coral rubble
<i>uniscutatus</i> Bartsch, 1994	re	p/s+p	sl	1+1	1+1	32S, 115E	SW Australia	2-13	sand
						21S, 116E	NW Australia		
<i>variolosus</i> Bartsch, 1994	re	s	sl	1+2	1+2	32S, 115E	SW Australia	tidal	sand
<i>versicolor</i> Otto, 2000	la	p/s	sl	1+1	1+1	19S, 149E	NE Australia	10	coral rubble
<i>xandarus</i> Otto, 2000	la	p/s	sl	1+1	1+1	19-22S, 147-153E	NE Australia	3-15	medium to coarse sand

I. Shape of OC: re, reduced; la, large, triangular or rounded.

II. P-2: p/s, with protuberance and distal seta; p/s-p, with protuberance and distal seta and a small distal protuberance; p+s, protuberance and seta at same level; s, seta only, protuberance lacking.

III. I-5, shape of ventromedial seta: pe, bipectinate; sl, slender; sp, spiniform; w, wide though setiform.

IV. III-6, number of ventral and parambulacral setae: 1+1, either one ventral and one parambulacral seta or two ventral setae and parambulacral setae lacking; 1+ 2, one ventral seta and pair of parambulacral setae; 1+2(d), one ventral seta and pair of parambulacral setae, one of pas doubled; 2+2, two ventral and pair of parambulacral setae.

V. IV-6, number of ventral and parambulacral setae: character states as above.

VI. Collecting area, coordinates:

VII. Collecting area, state/province/island or sea area:

VIII. Depth (in m):

IX. Habitat:

\*, doubtful records excluded; ?, information lacking or in need of confirmation.

bathyal (0–500 m). Representatives of the genus inhabit coarse sand, rubble, crustose and other algae, seagrasses, colonies of polychaetes and barnacles. There is a strikingly high diversity in the Australian fauna (Figure 6), 12 species being recorded from Queensland and 10 species from Western Australia. Though knowledge of the halacarid fauna of New Zealand is meagre, two of the 23 marine species recorded are representatives of *Simognathus*. From the southern South American continent six *Simognathus* species are recorded (Newell, 1984). Almost 70 marine halacarid species are known in total (Newell 1984; Bartsch 1988b, 1989). In contrast, only two out of 90 species in the Mediterranean, and two out of the known 120 shallow water halacarid species in the northeastern Atlantic, belong to the genus *Simognathus*. From eastern North America there is just a single record (Bartsch 1979b).

Table 3 presents a list of *Simognathus* species, their geographical areas and habitats, and some of their morphological characters. *Simognathus* species can roughly be divided into those with large OC, round or triangular in shape, and those with the OC reduced to narrow sclerites which often are obscured by the striated integument. Other characters are the shape of P-2, with or without any protuberance, and the number and arrangement of tarsal setae. Both in the north and south there are species with reduced OC, and the reduction seems to be correlated with a mesopsammal life style. Palps with the seta of P-2 and the protuberance situated at the same level are present in northern as well as in southern species. According to the setation of tarsi III and IV, *S. leiomerus* and *S. minor* are closely related. Other siblings are *S. abnormalus* and *S. salebrosus*; they share the characters: tibia I widest near its base, tarsus I very short and rotated versus the leg's axis, and presence of epimeral fossae. These characters are absent in other species.

Looking at the present day distributional records, one may expect *Simognathus* to be a Gondwanan genus with several founder species dispersing to the European, North American and Asian coastlines. More detailed analysis is necessary to elucidate these patterns.

#### ACKNOWLEDGEMENTS

The three species described were collected during two Marine Biological Workshops in Western Australia, on Rottneest Island and in Esperance. The workshops were organized by Dr F. Wells and colleagues. The travel expenses were covered by the Deutsche Forschungsgemeinschaft. To all my sincerest thanks.

#### REFERENCES

- Bartsch, I. (1978). Halacaridae (Acari) aus der Tiefsee des atlantischen Ozeans. *Cahiers de Biologie marine* 19: 47–62.
- Bartsch, I. (1979a). Halacaridae (Acari) aus der Subantarktis. *Cahiers de Biologie marine* 20: 325–339.
- Bartsch, I. (1979b). Halacaridae (Acari) von der Atlantikküste Nordamerikas. Beschreibung der Arten. *Mikrofauna Meeresboden* 79: 1–62.
- Bartsch, I. (1982). Weitere Halacaridae (Acari) aus dem Kanal von Moçambique. *Cahiers de Biologie marine* 23: 435–457.
- Bartsch, I. (1988a). Deep-sea halacarids (Acari) and description of a new species. *Journal of Natural History* 22: 811–821.
- Bartsch, I. (1988b). *Thalassophthirus auster* gen. et spec. nov., a halacarid mite (Acari) suspected of being a parasite. *Polar Research* 6: 181–184.
- Bartsch, I. (1989). *Rhombognathus auster*, a new rhombognathine mite (Acari, Halacaridae) from southern South America. *Zoologica Scripta* 18: 423–425.
- Bartsch, I. (1992). *Lohmannella dictyota* n. sp. und *L. africana* n. sp. von der Südhalbkugel (Arachnida: Acari: Halacaridae). *Senckenbergiana biologica* 72: 457–464.
- Bartsch, I. (1993a). Synopsis of the Antarctic Halacaroida (Acari), In J.W. Wägele and J. Sieg (eds) *Synopses of the Antarctic Benthos* 4: 1–176. Koeltz Scientific Books, Koenigstein.
- Bartsch, I. (1993b). Arenicolous Halacaridae (Acari) from south-western Australia, In F.E. Wells, D.I. Walker, H. Kirkman & R. Lethbridge (eds), *The Marine Flora and Fauna of Rottneest Island, Western Australia*: 45–71. Western Australian Museum, Perth.
- Bartsch, I. (1994). The genus *Simognathus* (Acari : Halacaridae), description of six new species from southern Australia and a tabular key to all species. *Acarologia* 35: 135–152.
- Bartsch, I. (1996). Halacarids (Halacaroida, Acari) in freshwater. Multiple invasions from the Paleozoic onwards. *Journal of Natural History* 30: 67–99.
- Bartsch, I. (2003a). Halacarids (Acari) from coral reefs off Norway, Northern Atlantic: Description of a new *Agauopsis* species. *Entomologische Mitteilungen aus dem Zoologischen Museum Hamburg* 14: 117–123.
- Bartsch, I. (2003b). Psammophilous halacarids (Halacaridae, Acari) from Dampier, Western Australia. Description of species and faunal comparison of the mesopsammal halacarid fauna of western and eastern Australia. *Records of the Western Australian Museum* 22: 23–45.
- Bartsch, I. (2004). Halacaridae (Acari) from the Great Meteor Seamount (Northeastern Atlantic). Description of *Simognathus* species. *Mitteilungen aus dem hamburgischen zoologischen Museum und Institut* 101: 185–196.
- Chatterjee, T. and Chang, C.Y. (2004). *Simognathus corensis*, a new halacarid mite (Halacaridae: Acari) from Korea. *Hydrobiologia* 529: 271–274.
- Knox, G.A. (1963). The biogeography and intertidal ecology of the Australasian coast. *Oceanography and marine Biology. Annual Review* 1: 341–404.

- Lohmann H. (1907). Die Meeresmilben der Deutschen Südpolar-Expedition 1901–1903. *Deutsche Südpolar Expedition 1901–1903*, 9: 361–413.
- Makarova, N.G. (1977). Marine mites (Acarina, Halacaridae) of the intertidal zone of the Kurile Islands. *Fauna pribreznih zon kurilskih ostrovov* 125–143. (In Russian)
- Makarova, N.G. (1978). Marine mites (Acarina, Halacaridae) of the intertidal zone of the Gulf of Anadyr (Bering Sea). In O.G. Kussakin (ed.) *The intertidal zone of the Bering Sea and southeastern Kamchatka* 131–149. Publishing House Nauka, Moskau (In Russian, with English summary).
- Murray, A. (1877). *Economic Entomology. Aptera*. South Kensington Museum Handbooks, London, 433 pp.
- Newell, I.M. (1947). A systematic and ecological study of Halacaridae of eastern North America. *Bulletin of the Bingham oceanographic Collection* 10: 1–232.
- Newell, I.M. (1951). Further studies on Alaskan Halacaridae (Acari). *American Museum Novitates* 1536: 1–56.
- Newell I.M. (1984). Antarctic Halacaroida. *Antarctic Research Series* 40: 1–284.
- O'Hara, T.D. and Poore, G.C.B. (2000). Patterns of distribution for southern Australian marine echinoderms and decapods. *Journal of Biogeography* 27: 1321–1335.
- Otto, J.C. (1994). New species of Halacaridae from Australia (Acarina : Prostigmata). *Acarologia* 35: 31–48.
- Otto, J.C. (2000). Halacaridae from the Great Barrier Reef and Coral Sea: the genera *Lohmannella*, *Scaptognathides* and *Scaptognathus* (Acarina: Halacaridae: Lohmannellinae). *Memoirs of the Queensland Museum* 45: 535–555
- Pepato, A.R. and Tiago, C.G. (2004). The genera *Acaromantis* and *Simognathus* (Simognathinae, Halacaridae) on the north coast of São Paulo State, Brazil. *Zootaxa* 615: 1–16.
- Pesic, V. (2004). Three interesting halacarid mite species (Acari: Halacaroida) from Montenegro and Italy. *Lauterbornia* 49: 37–42.
- Ponder, W.F. and Wells, F.E. (1998). Distribution and relationships of marine and estuarine fauna. In P.L. Beesley, G.J.B. Ross and A. Wells (eds), *Mollusca: The Southern Synthesis. Fauna of Australia 5A*, 77–80. CSIR Publishing, Melbourne.
- Trouessart, E. (1889). Sur les acariens marins des côtes de France. *Compte rendu de l'Académie des Sciences, Paris* 108: 1178–1181.
- Trouessart, E. (1901). Note sur les acariens marins (Halacaridae) récoltés par M. Henri Gadeau de Kerville dans la région d'Omonville-la-Rouge (Manche) et dans la fosse de la Hague (Juin–Juillet 1899). *Bulletin de la Société des amis des sciences naturelles de Rouen (Série 4)* 14: 247–266.
- Viets, K. (1927). Die Halacaridae der Nordsee. *Zeitschrift für wissenschaftliche Zoologie* 130: 83–173.
- Wells, F.E. and Walker, D.I. (1993). Introduction. In F.E. Wells, D.I. Walker, H. Kirkman and R. Lethbridge (eds), *The Marine Flora and Fauna of Rottnest Island, Western Australia*: 1–10. Western Australian Museum, Perth.
- Wilson, B.R. and Allen, G.R. (1987). Major components and distribution of marine fauna. In G.R. Dyne, and D.W. Walton (eds), *Fauna of Australia 1A*, 43–68. Australian Government Publishing Service, Canberra.