

## Checklist of the birds of Western Australia

R.E. Johnstone

Western Australian Museum, Francis Street, Perth, Western Australia 6000, Australia

### INTRODUCTION

In 1948 H.M. Whittell and D.L. Serventy published *A Systematic List of the Birds of Western Australia* (Public Library, Museum and Art Gallery of Western Australia Special Publication No. 1). The next forty-eight years witnessed such a resurgence in fieldwork, especially in the tropical north, deserts, seas, coasts and wetlands, that a further 110 species and many subspecies have been added to the State list. At the same time the classification of the birds has been improved and the distribution of the species and subspecies better defined. The area covered by this list includes the seas and islands of the adjacent continental shelf, including Ashmore Reef.

Considering its large size Western Australia has a relatively small avifauna. The 547 species recorded for the State include 385 breeding species, 145 non-breeding visitors and 17 foreign or introduced species. Sixteen species are endemic to the State namely: Carnaby's Cockatoo *Calyptorhynchus latirostris*, Baudin's Cockatoo *Calyptorhynchus baudinii*, Western Long-billed Corella *Cacatua pastinator*, Red-capped Parrot *Platycercus spurius*, Western Rosella *Platycercus icterotis*, Noisy Scrubbird *Atrichornis clamosus*, Red-winged Fairy-wren *Malurus elegans*, Black Grasswren *Amytornis housei*, Western Bristlebird *Dasyornis longirostris*, Dusky Gerygone *Gerygone tenebrosa*, Western Thornbill *Acanthiza inornata*, Western White-naped Honeyeater *Melithreptus chloropsis*, Western Spinebill *Acanthorhynchus superciliosus*, Western Little Wattlebird *Anthochaera lunulata*, White-breasted Robin *Eopsaltria georgiana*, and Red-eared Firetail *Stagonopleura oculata*.

The principal sources of reference for this list are the *Check-list of Birds of the World* by J.L. Peters *et al.* (1931–79), *Checklist of the Birds of Australia Non-passerines* by H.T. Condon (1975), *Interim List of Australian Songbirds* by R. Schodde (1975), *Field Guide to the Birds of Western Australia* by G.M. Storr and R.E. Johnstone (1979), *The New Guide to the Birds of New Zealand* by R.A. Falla *et al.* (1982), *Birds of New Guinea* by B.M. Beehler *et al.* (1986), *The Birds of Wallacea* by C.M.N. White and M.D. Bruce (1986), *Distribution and Taxonomy of the Birds of the World* by

C.G. Sibley and B.C. Monroe (1990), *The Taxonomy and Species of Birds of Australia and its Territories* by L. Christidis and W.E. Boles (1994), *An Annotated Checklist of the Birds of the Oriental Region* by T. Inskip, N. Lindsey and W. Duckworth (1996), the *Zoological Catalogue of Australia*, volume 37.2 (Columbidae to Coraciidae) by R. Schodde and I.J. Mason (1997), *Handbook of Western Australian Birds Volume I – non-passerines (Emu to Dollarbird)* by R.E. Johnstone and G.M. Storr (1998) and *The Directory of Australian Birds, Passerines* by R. Schodde and I.J. Mason (1999).

Species and subspecies requiring confirmation are enclosed in square brackets.

Manuscript received 5 November 1997; accepted 13 October 1999.

### Class AVES

#### Order STRUTHIONIFORMES

##### Family CASUARIIDAE

*Dromaius novaehollandiae* (Latham, 1790) Emu

#### Order GALLIFORMES

##### Family MEGAPODIIDAE

*Megapodius reinwardt* Dumont, 1823 Orange-legged Scrubfowl (Orange-footed Scrubfowl)

*M. r. tumulus* Gould, 1842  
*Leipoa ocellata* Gould, 1840 Malleefowl

##### Family PHASIANIDAE

*Coturnix pectoralis* Gould, 1837 Stubble Quail  
*Coturnix ypsiloniphora* Bosc, 1792 Brown Quail

*C. y. cervina* Gould, 1865  
*C. y. australis* (Latham, 1802)

*Coturnix chinensis* (Linnaeus, 1766) King Quail  
*C. c. lineata* (Scopoli, 1786)

<sup>1</sup>*Phasianus colchicus* Linnaeus, 1758 Common Pheasant (Domestic Pheasant)  
*Pavo cristatus* Linnaeus, 1758 Common Peafowl (Indian Peafowl)

<sup>1</sup> According to Condon (1975), the Common Pheasants introduced into Australia are of mixed hybrid stock. Small feral colonies of Silver Pheasants *Lophura nycthemera*, Domestic Turkeys *Meleagris gallopavo* and Helmeted Guinea Fowl *Numida meleagris* occurred briefly in Western Australia.

## Order ANSERIFORMES

### Family ANATIDAE

<i>Anseranas semipalmata</i> (Latham, 1798)	
	Magpie Goose (Pied Goose)
<i>Dendrocygna eytoni</i> (Eyton, 1838)	
	Plumed Whistling Duck
<i>Dendrocygna arcuata</i> (Horsfield, 1824)	Wandering Whistling Duck (Chestnut Whistling Duck)
<i>Oxyura australis</i> Gould, 1836	Blue-billed Duck
<i>Biziura lobata</i> (Shaw, 1796)	Musk Duck
<i>Stictonetta naevosa</i> (Gould, 1841)	Freckled Duck
<i>Cygnus olor</i> (Gmelin, 1789)	Mute Swan
<i>Cygnus atratus</i> (Latham, 1790)	Black Swan
<i>Cereopsis novaehollandiae</i> Latham, 1802	Cape Barren Goose
<i>C. n. grisea</i> (Vieillot, 1818)	
<i>Tadorna tadornoides</i> (Jardine and Selby, 1828)	Australian Shelduck (Mountain Duck)
<i>Tadorna radjah</i> (Lesson, 1828)	Radjah Shelduck
<i>T. r. rufitergum</i> Harttert, 1905	Burdekin Duck
<i>Nettapus pulchellus</i> Gould, 1842	Green Pygmy Goose
<i>Chenonetta jubata</i> (Latham, 1802)	
	Australian Wood Duck (Wood Duck)
<i>Anas gracilis</i> Buller, 1869	Grey Teal
<i>Anas castanea</i> (Eyton, 1838)	Chestnut Teal
<i>Anas platyrhynchos</i> Linnaeus, 1758	Mallard
<i>Anas superciliosa</i> Gmelin, 1789	Pacific Black Duck
<i>Anas acuta</i> Linnaeus, 1758	Pintail
<i>A. a. acuta</i> Linnaeus, 1758	
<i>Anas querquedula</i> Linnaeus, 1758	Garganey
<i>Anas rhynchos</i> Latham, 1802	Australasian Shoveler
<i>Anas r. rhynchos</i> Latham, 1802	
<i>Anas clypeata</i> Linnaeus, 1758	Northern Shoveler
<i>Malacorhynchus membranaceus</i> (Latham, 1802)	
	Pink-eared Duck
<i>Aythya australis</i> (Eyton, 1838)	Hardhead

## Order PODICIPEDIFORMES

### Family PODICIPEDIDAE

<i>Tachybaptus novaehollandiae</i> (Stephens, 1826)	
	Australasian Grebe (Black-throated Grebe)
<i>T. n. novaehollandiae</i> (Stephens, 1826)	
<i>Poliocephalus poliocephalus</i> (Jardine and Selby, 1827)	Hoary-headed Grebe
<i>Podiceps cristatus</i> (Linnaeus, 1758)	
	Great Crested Grebe
<i>P. c. australis</i> Gould, 1844	

## Order SPHENISCIFORMES

### Family SPHENISCIDAE

<i>Aptenodytes patagonicus</i> Miller, 1778	King Penguin
<i>A. p. halli</i> Mathews, 1911	

## Eudyptes chrysocome (Forster, 1781)

Rockhopper Penguin

*E. c. filholi* Hutton, 1879

*E. c. moseleyi* Mathews and Iredale, 1921

## Eudyptes pachyrhynchus G. R. Gray, 1845

Fiordland Penguin

## Eudyptes sclateri Buller, 1888

Erect-crested Penguin

## Eudyptes chrysolophus (Brandt, 1837)

Macaroni Penguin

<sup>2</sup>*E. c. schlegeli* Finsch, 1876

Royal Penguin

## Eudyptula minor (Forster, 1781)

Little Penguin

*E. m. novaehollandiae* (Stephens, 1826)

## Order PROCELLARIIFORMES

### Family PROCELLARIIDAE

## Macronectes giganteus (Gmelin, 1789)

Southern Giant Petrel

## Macronectes halli Mathews, 1912

Northern Giant Petrel

## Fulmarus glacialis (A. Smith, 1840)

Southern Fulmar

## Thalassoica antarctica (Gmelin, 1789)

Antarctic Petrel

## Daption capense (Linnaeus, 1758)

Cape Petrel

## Pterodroma rostrata (Peale, 1848)

Tahiti Petrel

## Pterodroma macroptera (Smith, 1840)

Great-winged Petrel

*P. m. macroptera* (Smith, 1840)

*P. m. gouldi* (Hutton, 1869)

## Pterodroma lessonii (Garnot, 1826)

White-headed Petrel

## Pterodroma mollis (Gould, 1844)

Soft-plumaged Petrel

*P. m. mollis* (Gould, 1844)

*P. m. dubius* Mathews, 1924

## Pterodroma brevirostris (Lesson, 1831)

Kerguelen Petrel

## Pterodroma arminjoniana (Giglioni and Salvadori, 1869)

Herald Petrel

*P. a. arminjoniana* (Giglioni and Salvadori, 1869)

## Pterodroma baraui (Jouanin, 1964)

Barau's Petrel

## Pterodroma leucoptera (Gould, 1844)

Gould's Petrel

*P. l. leucoptera* (Gould, 1844)

## Halobaena caerulea (Gmelin, 1789)

Blue Petrel

## Pachyptila vittata (G. Forster, 1777)

Broad-billed Prion

## Pachyptila salvini (Mathews, 1912)

Salvin's Prion

*P. s. salvini* (Mathews, 1912)

*P. s. macgillivrayi* (Mathews, 1912)

## Pachyptila desolata (Gmelin, 1789)

Antarctic Prion

## Pachyptila belcheri (Mathews, 1912)

Slender-billed Prion

## Pachyptila turtur (Kuhl, 1820)

Fairy Prion

## Bulweria bulwerii (Jardine and Selby, 1828)

Bulwer's Petrel

<sup>2</sup> Often treated as a full species *E. schlegeli*.

<i>Procellaria aequinoctialis</i> Linnaeus, 1758		<i>Oceanites nereis</i> (Gould, 1841)
<i>P. a. aequinoctialis</i> Linnaeus, 1758	White-chinned Petrel	Grey-backed Storm Petrel
<i>Procellaria cinerea</i> Gmelin, 1789	Grey Petrel	<i>Oceanites marinus</i> (Latham, 1790)
<i>Calonectris leucomelas</i> (Temminck, 1835)		White-faced Storm Petrel
	Streaked Shearwater	<i>O. m. dulciae</i> (Mathews, 1912)
<i>Puffinus pacificus</i> (Gmelin, 1789)		<i>Oceanites tropicus</i> (Gould, 1844)
<i>P. p. pacificus</i> (Gmelin, 1789)	Wedge-tailed Shearwater	Black-bellied Storm Petrel
<i>Puffinus carneipes</i> Gould, 1844		<i>Oceanodroma leucorhoa</i> (Vieillot, 1818)
<i>P. c. carneipes</i> Gould, 1844	Fleshy-footed Shearwater	Leach's Storm Petrel
<i>Puffinus griseus</i> (Gmelin, 1789)	Sooty Shearwater	<i>O. l. leucorhoa</i> (Vieillot, 1818)
<i>Puffinus tenuirostris</i> (Temminck, 1835)		<sup>5</sup> <i>Oceanodroma monorhis</i> (Swinhoe, 1867)
<i>P. t. tenuirostris</i> (Temminck, 1835)	Short-tailed Shearwater	Swinhoe's Storm Petrel
<i>Puffinus gavia</i> (J. R. Forster, 1844)		<i>Oceanodroma matsudariae</i> Kuroda, 1922
<i>P. g. gavia</i> (J. R. Forster, 1844)	Fluttering Shearwater	Matsudaira's Storm Petrel
<i>Puffinus huttoni</i> Mathews, 1912	Hutton's Shearwater	
<i>Puffinus assimilis</i> Gould, 1838	Little Shearwater	
<i>P. a. assimilis</i> Gould, 1838		
<b>Family PELECANOIDIDAE</b>		
<i>Pelecanoides urinatrix</i> (Gmelin, 1789)		<b>Order PELECANIFORMES</b>
<i>P. u. urinatrix</i> (Gmelin, 1789)	Common Diving Petrel	<b>Family PHAETHONTIDAE</b>
<i>P. u. exsul</i> Salvin, 1896		<i>Phaethon rubricauda</i> Boddaert, 1783
Red-tailed Tropicbird		
<b>Family DIOMEDEIDAE</b>		
<i>Diomedea exulans</i> Linnaeus, 1758		<i>Phaethon lepturus</i> Daudin, 1802
<i>D. e. exulans</i> Linnaeus, 1758	Wandering Albatross	White-tailed Tropicbird
<i>D. e. chionoptera</i> Salvin, 1896		
<sup>3</sup> <i>D. e. gibsoni</i> Robertson and Warham 1992		
<i>Diomedea epomophora</i> Lesson, 1825		
<i>D. e. epomophora</i> Lesson, 1825	Royal Albatross (Southern Royal Albatross)	
<i>Diomedea melanophris</i> Temminck, 1828		
<sup>4</sup> <i>D. m. melanophris</i> Temminck, 1828	Black-browed Albatross	
<i>D. m. melanophris</i> Temminck, 1828		
<i>Diomedea cauta</i> Gould, 1841	Shy Albatross	
<i>D. c. cauta</i> Gould, 1841		
<i>D. c. salvini</i> (Rothschild, 1893)		
<i>Diomedea chrysostoma</i> J. R. Forster, 1785		
<i>D. c. chrysostoma</i> J. R. Forster, 1785	Grey-headed Albatross	
<i>Diomedea chlororhynchos</i> Gmelin, 1789		
<i>D. c. chlororhynchos</i> Gmelin, 1789	Yellow-nosed Albatross	
<i>D. c. chlororhynchos</i> Gmelin, 1789		
<i>D. c. bassi</i> Mathews, 1912		
<i>Phoebetria fusca</i> (Hilsenberg, 1822)	Sooty Albatross	
<i>Phoebetria palpebrata</i> (J. R. Forster, 1785)		
<i>P. p. fusca</i> (Hilsenberg, 1822)	Light-mantled Sooty Albatross	
<b>Family HYDROBATIDAE</b>		
<i>Oceanites oceanicus</i> (Kuhl, 1820)		
<i>O. o. oceanicus</i> (Kuhl, 1820)	Wilson's Storm Petrel	
<b>Family SULIDAE</b>		
<i>Sula abbotti</i> Ridgway, 1893		Abbott's Booby
<i>Sula (bassana) capensis</i> (Lichtenstein, 1823)		Cape Gannet
<i>Sula (bassana) serrator</i> (G. R. Gray, 1843)		Australasian Gannet
<i>Sula dactylatra</i> Lesson, 1831		Masked Booby
<i>S. d. personata</i> Gould, 1846		
<i>Sula sula</i> (Linnaeus, 1766)		Red-footed Booby
<i>Sula leucogaster</i> Boddaert, 1783		Brown Booby
<i>S. l. plotus</i> (J. R. Forster, 1844)		
<b>Family ANHINGIDAE</b>		
<i>Anhinga melanogaster</i> Pennant, 1769		Darter
<i>A. m. novaehollandiae</i> (Gould, 1847)		
<b>Family PHALACROCORACIDAE</b>		
<i>Phalacrocorax carbo</i> (Linnaeus, 1758)		Great Cormorant
<i>P. c. novaehollandiae</i> Stephens, 1826		
<i>Phalacrocorax varius</i> (Gmelin, 1789)	Pied Cormorant	
<i>P. v. hypoleucus</i> (Brandt, 1837)		
<i>Phalacrocorax sulcirostris</i> (Brandt, 1837)		Little Black Cormorant
<i>Phalacrocorax verrucosus</i> (Cabanis, 1875)		Kerguelen Cormorant

<sup>3</sup> Gibson's Albatross. *D. e. gibsoni*. Sometimes considered specifically distinct from *exulans*.

<sup>4</sup> *Diomedea melanophris*, *D. cauta*, *D. chrysostoma* and *D. chlororhynchos* were placed in the genus *Thalassarche* by Nunn et al. 1996 *Auk* 113(4): 784-801 and according to some recent authors *D. cauta salvini* (Salvin's Albatross) and *D. chlororhynchos bassi* (Indian Yellow-nosed Albatross) should be treated as full species. The composition of genera, species and subspecies within the Diomedidae is still far from resolved.

<sup>5</sup> Sometimes treated as conspecific with *O. leucorhoa*.

<i>Phalacrocorax fuscescens</i> (Vieillot, 1817)	
	Black-faced Cormorant
<i>Phalacrocorax melanoleucus</i> (Vieillot, 1817)	
	Little Pied Cormorant

*P. m. melanoleucus* (Vieillot, 1817)

#### Family PELECANIDAE

<i>Pelecanus conspicillatus</i> Temminck, 1824	
	Australian Pelican

#### Family FREGATIDAE

<i>Fregata minor</i> (Gmelin, 1789)	Greater Frigatebird
<i>Fregata ariel</i> (G. R. Gray, 1845)	Lesser Frigatebird
<i>Fregata andrewsi</i> Mathews, 1914	Christmas Island Frigatebird

#### Order CICONIIFORMES

##### Family ARDEIDAE

<i>Ardea sumatrana</i> Raffles, 1822	Great-billed Heron
<i>Ardea pacifica</i> Latham, 1802	White-necked Heron
<i>Ardea picata</i> Gould, 1845	Pied Heron
<i>Ardea novaehollandiae</i> Latham, 1790	White-faced Heron
<i>Ardea alba</i> Linnaeus, 1758	Great Egret
<i>A. a. modesta</i> J.E. Gray, 1831	
<i>Ardea intermedia</i> Wagler, 1829	Intermediate Egret
<i>A. i. intermedia</i> Wagler, 1829	
<i>Ardea garzetta</i> Linnaeus, 1766	Little Egret
<i>A. g. nigripes</i> Temminck, 1840	
<i>Ardea sacra</i> Gmelin, 1789	Eastern Reef Heron (Eastern Reef Egret)
<i>A. s. sacra</i> Gmelin, 1789	
<i>Ardea ibis</i> Linnaeus, 1758	Cattle Egret
<i>A. i. coromanda</i> (Boddaert, 1783)	
<i>Butorides striatus</i> Linnaeus, 1766	Striated Heron (Mangrove Heron)
<i>B. s. stagnatilis</i> (Gould, 1848)	
<i>Nycticorax nycticorax</i> (Linnaeus, 1758)	Black-crowned Night Heron
<i>N. n. nycticorax</i> (Linnaeus, 1758)	
<i>Nycticorax caledonicus</i> (Gmelin, 1789)	Rufous Night Heron
<i>N. c. hilli</i> Mathews, 1912	
<i>Ixobrychus minutus</i> (Linnaeus, 1766)	Little Bittern
<i>I. m. dubius</i> Mathews, 1912	
<i>Ixobrychus sinensis</i> (Gmelin, 1789)	Yellow Bittern (Chinese Little Bittern)
<i>Ixobrychus flavicollis</i> (Latham, 1790)	Black Bittern
<i>I. f. australis</i> (Lesson, 1831)	

<i>Botaurus poiciloptilus</i> (Wagler, 1827)	
	Australasian Bittern

#### Family THRESKIORNITHIDAE

<i>Plegadis falcinellus</i> (Linnaeus, 1766)	Glossy Ibis
<sup>6</sup> <i>Threskiornis molucca</i> (Cuvier, 1829)	Australian White Ibis
<i>Threskiornis spinicollis</i> (Jameson, 1835)	
<sup>7</sup> <i>Platalea regia</i> Gould, 1838	Straw-necked Ibis
<i>Platalea flavipes</i> Gould, 1838	Royal Spoonbill
	Yellow-billed Spoonbill

#### Family CICONIIDAE

<i>Ephippiorhynchus asiaticus</i> (Latham, 1790)	
	Black-necked Stork

*E. a. australis* (Shaw, 1800)

#### Order FALCONIFORMES

##### Family ACCIPITRIDAE

<i>Pandion haliaetus</i> (Linnaeus, 1758)	Osprey
<i>P. h. cristatus</i> (Vieillot, 1816)	
<i>Elanus caeruleus</i> (Desfontaines, 1789)	Black-shouldered Kite
<sup>8</sup> <i>E. c. axillaris</i> Latham, 1802	Australian Black-shouldered Kite
<i>Elanus scriptus</i> Gould, 1842	Letter-winged Kite
<i>Aviceda subcristata</i> (Gould, 1838)	Pacific Baza
<i>A. s. subcristata</i> (Gould, 1838)	
<sup>9</sup> <i>Hamirostra isura</i> (Gould, 1838)	Square-tailed Kite
<i>Hamirostra melanosternon</i> (Gould, 1841)	
	Black-breasted Buzzard
<i>Milvus migrans</i> (Boddaert, 1783)	Black Kite
<i>M. m. affinis</i> Gould, 1838	
<i>Haliastur sphenurus</i> (Vieillot, 1818)	Whistling Kite
<i>Haliastur indus</i> (Boddaert, 1783)	Brahminy Kite
<i>H. i. girrenera</i> (Vieillot, 1822)	
<i>Accipiter novaehollandiae</i> (Gmelin, 1788)	Grey Goshawk
<i>A. n. novaehollandiae</i> (Gmelin, 1788)	
<i>Accipiter fasciatus</i> (Vigors and Horsfield, 1827)	Brown Goshawk
<i>A. f. fasciatus</i> (Vigors and Horsfield, 1827)	
<i>A. f. didimus</i> (Mathews, 1912)	
<i>Accipiter cirrocephalus</i> (Vieillot, 1817)	Collared Sparrowhawk
<i>A. c. cirrocephalus</i> (Vieillot, 1817)	
<i>Erythrotriorchis radiatus</i> (Latham, 1802)	
	Red Goshawk
<i>Aquila morphnoides</i> Gould, 1841	Little Eagle
<i>A. m. morphnoides</i> Gould, 1841	

<sup>6</sup> Sometimes treated as a subspecies of the Sacred Ibis *T. aethiopicus*.

<sup>7</sup> Forming a superspecies with Eurasian Spoonbill *P. leucorodia* of Europe, Asia and Africa.

<sup>8</sup> Sometimes treated as a full species *E. axillaris*.

<sup>9</sup> Usually placed in monotypic genus *Lophoictinia* but along with its single congener *H. melanosternon* comprise an old Australian group without close relatives.

<i>Aquila audax</i> (Latham, 1802)	Wedge-tailed Eagle
<i>Haliaeetus leucogaster</i> (Gmelin, 1788)	White-bellied Sea-Eagle
<i>Circus assimilis</i> Jardine and Selby, 1828	Spotted Harrier
<i>Circus approximans</i> Peale, 1848	Swamp Harrier

<i>Gallinula ventralis</i> Gould, 1837	Black-tailed Native-hen
<i>Gallinula tenebrosa</i> Gould, 1846	Dusky Moorhen
<i>G. t. tenebrosa</i> Gould, 1846	
<i>Fulica atra</i> Linnaeus, 1758	Eurasian Coot
<i>F. a. australis</i> Gould, 1845	

### Family FALCONIDAE

<i>Falco berigora</i> Vigors and Horsfield, 1827	Brown Falcon
<i>F. b. berigora</i> Vigors and Horsfield, 1827	
<i>Falco cenchroides</i> Vigors and Horsfield, 1827	Australian Kestrel
<i>F. c. cenchroides</i>	
[ <i>Falco subbuteo</i> Linnaeus, 1758]	Eurasian Hobby]
<i>Falco longipennis</i> Swainson, 1837	Australian Hobby
<i>F. l. longipennis</i> Swainson, 1837	
<i>Falco hypoleucus</i> Gould, 1841	Grey Falcon
<i>Falco peregrinus</i> Tunstall, 1771	Peregrine Falcon
<i>F. p. macropus</i> Swainson, 1837	
<i>Falco subniger</i> G. R. Gray, 1843	Black Falcon

### Order GRUIFORMES

Family GRUIDAE	
<i>Grus antigone</i> (Linnaeus, 1758)	Sarus Crane
<i>G. a. gillae</i> Schodde, 1988	
<i>Grus rubicunda</i> (Perry, 1810)	Brolga

### Family RALLIDAE

<i>Rallina fasciata</i> (Raffles, 1822)	Red-legged Crake
<i>Gallirallus philippensis</i> (Linnaeus, 1766)	
	Buff-banded Rail
<i>G. p. mellori</i> (Mathews, 1912)	
<i>Crex crex</i> (Linnaeus, 1758)	Corncrake
<i>Rallus pectoralis</i> Temminck, 1831	Lewin's Rail
<i>R. p. clelandi</i> Mathews, 1911	
<i>Amaurornis olivaceus</i> (Meyen, 1834)	Bush-hen
<i>A. o. ruficrissus</i> (Gould, 1869)	
<i>Porzana pusilla</i> (Pallas, 1776)	Baillon's Crake
<i>P. p. palustris</i> Gould, 1843	
<i>Porzana fluminea</i> Gould, 1843	Australian Spotted Crake
<i>Porzana tabuensis</i> (Gmelin, 1789)	Spotless Crake
<i>Porzana cinerea</i> (Vieillot, 1819)	White-browed Crake
<i>Eulabeornis castaneoventris</i> Gould, 1844	Chestnut Rail
<i>E. c. castaneoventris</i> Gould, 1844	
<i>Porphyrio porphyrio</i> (Linnaeus, 1758)	Purple Swamphen
<i>P. p. melanotus</i> Temminck, 1820	
<i>P. p. bellus</i> Gould, 1841	

### Family OTIDIDAE

<sup>10</sup> <i>Ardeotis australis</i> (J. E. Gray, 1829)	Australian Bustard
--	--------------------

### Order TURNICIFORMES

#### Family TURNICIDAE

<i>Turnix maculosa</i> (Temminck, 1815)	Red-backed Button-quail
<i>T. m. melanota</i> (Gould, 1837)	
<i>Turnix varia</i> (Latham, 1802)	Painted Button-quail
<i>T. v. varia</i>	
<i>T. v. scintillans</i> (Gould, 1845)	
<i>Turnix (castanota) castanota</i> (Gould, 1840)	Chestnut-backed Button-quail
<i>Turnix pyrrhothorax</i> (Gould, 1841)	Red-chested Button-quail
<i>Turnix velox</i> (Gould, 1841)	Little Button-quail

### Order CHARADRIIFORMES

#### Family SCOLOPACIDAE

<i>Gallinago hardwickii</i> (J. E. Gray, 1831)	Latham's Snipe
<i>Gallinago stenura</i> (Bonaparte, 1830)	Pin-tailed Snipe
<i>Gallinago megala</i> Swinhoe, 1861	Swinhoe's Snipe
<i>Limosa limosa</i> (Linnaeus, 1758)	Black-tailed Godwit
<i>L. l. melanuroides</i> Gould, 1846	
<i>Limosa lapponica</i> (Linnaeus, 1758)	Bar-tailed Godwit
[ <i>L. l. lapponica</i> ]	
<i>L. l. menzbieri</i> Portenko, 1936	
<i>Numenius minutus</i> Gould, 1841	Little Curlew
<i>Numenius phaeopus</i> (Linnaeus, 1758)	Whimbrel
<i>N. p. variegatus</i> (Scopoli, 1786)	
<i>Numenius madagascariensis</i> (Linnaeus, 1766)	
	Eastern Curlew
<i>Tringa erythropus</i> (Pallas, 1764)	Spotted Redshank
<sup>11</sup> <i>Tringa totanus</i> (Linnaeus, 1758)	
	Common Redshank
<i>Tringa stagnatilis</i> (Bechstein, 1803)	Marsh Sandpiper
<i>Tringa nebularia</i> (Gunnerus, 1767)	
	Common Greenshank
<i>Tringa glareola</i> Linnaeus, 1758	Wood Sandpiper
<sup>12</sup> <i>Tringa cinerea</i> (Güldenstädt, 1774)	Terek Sandpiper

<sup>10</sup> Previously merged with genus *Otis*.

<sup>11</sup> Subspecific identity of birds visiting Western Australia unknown.

<sup>12</sup> Sometimes placed in genus *Xenus*. *Tringa terek* (Latham, 1970) is synonym.

*Tringa hypoleucus* Linnaeus, 1758

	Common Sandpiper
<i>Tringa brevipes</i> (Vieillot, 1816)	Grey-tailed Tattler
<i>Arenaria interpres</i> (Linnaeus, 1758)	Ruddy Turnstone
<i>A. i. interpres</i> (Linnaeus, 1758)	
<i>Limnodromus semipalmatus</i> (Blyth, 1848)	
	Asian Dowitcher
<i>Calidris canutus</i> (Linnaeus, 1758)	Red Knot
<i>C. c. rogersi</i> Mathews, 1913	
<i>Calidris tenuirostris</i> (Horsfield, 1821)	Great Knot
<i>Calidris alba</i> (Pallas, 1764)	Sanderling
<i>Calidris ruficollis</i> (Pallas, 1776)	Red-necked Stint
<i>Calidris minuta</i> (Leisler, 1812)	Little Stint
<i>Calidris subminuta</i> (Middendorff, 1853)	
	Long-toed Stint
<i>Calidris fuscicollis</i> (Vieillot, 1819)	White-rumped Sandpiper
<i>Calidris bairdii</i> (Coues, 1861)	Baird's Sandpiper
<i>Calidris melanotos</i> (Vieillot, 1819)	Pectoral Sandpiper
<i>Calidris acuminata</i> (Horsfield, 1821)	
	Sharp-tailed Sandpiper
<i>Calidris alpina</i> (Linnaeus, 1758)	Dunlin
<i>C. a. sakhalina</i> (Vieillot, 1816)	
<i>Calidris ferruginea</i> (Pontoppidan, 1763)	Curlew Sandpiper
<i>Limicola falcinellus</i> (Pontoppidan, 1763)	
	Broad-billed Sandpiper
<i>L. f. sibiricus</i> Dresser, 1876	
<i>Tryngites subruficollis</i> (Vieillot, 1819)	
	Buff-breasted Sandpiper
<i>Philomachus pugnax</i> Linnaeus, 1758	Ruff
<i>Phalaropus lobatus</i> (Linnaeus, 1758)	
	Red-necked Phalarope

## Family ROSTRATULIDAE

<i>Rostratula benghalensis</i> (Linnaeus, 1758)	
	Painted Snipe
<i>R. b. australis</i> (Gould, 1838)	

## Family JACANIDAE

<sup>13</sup> <i>Metopidius gallinaceus</i> (Temminck, 1828)	
	Comb-crested Jacana
<i>Hydrophasianus chirurgus</i> (Scopoli, 1786)	
	Pheasant-tailed Jacana

## Family BURHINIDAE

<i>Burhinus grallarius</i> (Latham, 1802)	
	Bush Stone-curlew
<i>Esacus neglectus</i> Mathews, 1912	Beach Stone-curlew

## Family HAEMATOPODIDAE

<i>Haematopus longirostris</i> Vieillot, 1817	
	Pied Oystercatcher
<i>Haematopus fuliginosus</i> Gould, 1845	
	Sooty Oystercatcher
<i>H. f. fuliginosus</i> Gould, 1845	
<i>H. f. ophthalmicus</i> Castelnau and Ramsay, 1877	

## Family RECURVIROSTRIDAE

<i>Himantopus himantopus</i> (Linnaeus, 1758)	
	Black-winged Stilt
<i>H. h. leucocephalus</i> Gould, 1837	
<i>Cladorhynchus leucocephalus</i> (Vieillot, 1816)	
	Banded Stilt
<i>Recurvirostra novaehollandiae</i> Vieillot, 1816	
	Red-necked Avocet

## Family CHARADRIIDAE

<i>Vanellus miles</i> (Boddaert, 1783)	Masked Lapwing
<i>V. m. miles</i> (Boddaert, 1783)	
<i>V. m. novaehollandiae</i> Stephens, 1819	
<i>Vanellus tricolor</i> (Vieillot, 1818)	Banded Lapwing
<i>Pluvialis squatarola</i> (Linnaeus, 1758)	Grey Plover
<i>Pluvialis fulva</i> (Gmelin, 1789)	Pacific Golden Plover
<i>Pluvialis dominica</i> (Müller, 1776)	American Golden Plover
<i>Charadrius dubius</i> Scopoli, 1786	Little Ringed Plover
<i>C. d. curonicus</i> Gmelin, 1789	
<i>Charadrius alexandrinus</i> Linnaeus, 1758	Kentish Plover
	[ <i>C. a. dealbatus</i> (Swinhoe, 1870)]
<i>Charadrius ruficollis</i> Temminck, 1822	
	Red-capped Plover
<i>Charadrius bicinctus</i> Jardine and Selby, 1827	
	Double-banded Plover
<i>C. b. bicinctus</i> Jardine and Selby, 1827	
<i>Charadrius mongolus</i> Pallas, 1776	Lesser Sand Plover
<i>C. m. mongolus</i> Pallas, 1776	
<i>C. m. atrifrons</i> Wagler, 1829	
<i>Charadrius leschenaultii</i> Lesson, 1826	Greater Sand Plover
	[ <i>C. l. leschenaultii</i> Lesson, 1826]
<i>Charadrius melanops</i> Vieillot, 1818	
	Black-fronted Dotterel
<i>Charadrius rubricollis</i> (Gmelin, 1789)	Hooded Plover
<i>Charadrius veredus</i> Gould, 1848	Oriental Plover
<i>Erythrogonyx cinctus</i> Gould, 1838	
	Red-kneed Dotterel
<i>Peltohyas australis</i> (Gould, 1841)	Inland Dotterel

<sup>13</sup> Sometimes placed in monotypic genus *Irediparra*.

## Family GLAREOLIDAE

*Stiltia isabella* (Vieillot, 1816) Australian Pratincole  
*Glareola maldivarum* J. R. Forster, 1795 Oriental Pratincole

## Family LARIDAE

<sup>14</sup> <i>Catharacta antarctica</i> (Lesson, 1831)	Brown Skua
<i>C. a. lonnbergi</i> (Mathews, 1912)	
<sup>14</sup> <i>Catharacta maccormicki</i> (Saunders, 1893)	South Polar Skua
<i>Stercorarius pomarinus</i> (Temminck, 1815)	Pomarine Skua
<i>Stercorarius parasiticus</i> (Linnaeus, 1758)	Arctic Skua
<i>Stercorarius longicaudus</i> Vieillot, 1819	Long-tailed Skua
<i>Larus pacificus</i> Latham, 1802	Pacific Gull
<i>L. p. georgii</i> King, 1826	
<i>Larus crassirostris</i> Vieillot, 1818	Black-tailed Gull
<i>Larus dominicanus</i> Lichtenstein, 1823	Kelp Gull
<i>Larus novaehollandiae</i> Stephens, 1826	Silver Gull
<i>L. n. novaehollandiae</i> Stephens, 1826	
<i>Larus ridibundus</i> Linnaeus, 1766	Black-headed Gull
<i>Larus atricilla</i> Linnaeus, 1758	Laughing Gull
[ <i>L. a. megalopterus</i> Bruch, 1855]	
<i>Larus pipixcan</i> Wagler, 1831	Franklin's Gull
<i>Larus sabini</i> Sabine, 1819	Sabine's Gull
<i>Sterna nilotica</i> Gmelin, 1789	Gull-billed Tern
[ <i>S. n. nilotica</i> ]	
<i>S. n. affinis</i> Horsfield, 1821	
<i>S. n. macrotarsa</i> Gould, 1837	
<i>Sterna caspia</i> Pallas, 1770	Caspian Tern
<i>Sterna bengalensis</i> Lesson, 1831	Lesser Crested Tern
<i>Sterna bergii</i> Lichtenstein, 1823	Crested Tern
<i>Sterna dougallii</i> Montagu, 1813	Roseate Tern
<sup>15</sup> <i>S. d. gracilis</i> Gould, 1845	
<i>Sterna striata</i> Gmelin, 1789	White-fronted Tern
<i>Sterna sumatrana</i> Raffles, 1822	Black-naped Tern
<i>Sterna hirundo</i> Linnaeus, 1758	Common Tern
<i>S. h. hirundo</i> Linnaeus, 1758	
<i>S. h. longipennis</i> Nordmann, 1835	
<i>Sterna paradisaea</i> Pontoppidan, 1763	Arctic Tern
<i>Sterna vittata</i> Gmelin, 1789	Antarctic Tern
<i>Sterna (albifrons) sinensis</i> Gmelin, 1789	(Little Tern)
	White-shafted Little Tern
<i>Sterna nereis</i> (Gould, 1843)	Fairy Tern
<i>S. n. nereis</i> (Gould, 1843)	
<i>Sterna anaethetus</i> Scopoli, 1786	Bridled Tern
<i>S. a. anaethetus</i> Scopoli, 1786	

<i>Sterna fuscata</i> Linnaeus, 1766	Sooty Tern
<i>S. f. nubilosa</i> Sparrman, 1788	
<i>Sterna hybrida</i> Pallas, 1811	Whiskered Tern
<i>S. h. javanica</i> Horsfield, 1821	
<i>Sterna leucoptera</i> Temminck, 1815	
	White-winged Black Tern
<i>Anous stolidus</i> (Linnaeus, 1758)	Common Noddy
<i>A. s. pileatus</i> (Scopoli, 1786)	
<i>Anous minutus</i> Boie, 1844	Black Noddy
[i. m. minutus]	
<i>Anous tenuirostris</i> (Temminck, 1823)	Lesser Noddy
<i>A. t. melanops</i> Gould, 1846	

## **Order COLUMBIFORMES**

## Family COLUMBIDAE

<i>Columba livia</i> Gmelin, 1789	Domestic Pigeon
<i>Streptopelia senegalensis</i> (Linnaeus, 1766)	
	Laughing Turtle-Dove
<i>S. s. senegalensis</i> (Linnaeus, 1766)	
<sup>16</sup> <i>Streptopelia chinensis</i> (Scopoli, 1786)	Spotted Turtle Dove
<i>S. c. tigrina</i> (Temminck, 1810)	
<i>Chalcophaps indica</i> (Linnaeus, 1758)	Emerald Dove
<i>C. i. longirostris</i> Gould, 1848	
<i>Phaps chalcoptera</i> (Latham, 1790)	Common Bronzewing
<i>Phaps elegans</i> (Temminck, 1810)	Brush Bronzewing
<i>Phaps histrio</i> (Gould, 1841)	Flock Pigeon
<i>Ocyphaps lophotes</i> (Temminck, 1822)	Crested Pigeon
<sup>17</sup> <i>Geophaps plumifera</i> Gould, 1842	Spinifex Pigeon
<i>Geophaps smithii</i> (Jardine and Selby, 1830)	Partridge Pigeon
<i>G. s. smithii</i> (Jardine and Selby, 1830)	
<i>G. s. blaaui</i> Mathews, 1912	
<sup>18</sup> <i>Petrophassa albipennis</i> Gould, 1841	White-quilled Rock Pigeon
<i>P. a. albipennis</i> Gould, 1841	
<i>Geopelia cuneata</i> (Latham, 1802)	Diamond Dove
<i>Geopelia striata</i> (Linnaeus, 1766)	Peaceful Dove
<i>G. s. placida</i> Gould, 1844	
<i>Geopelia humeralis</i> Temminck, 1821	Bar-shouldered Dove
<i>Caloenas nicobarica</i> (Linnaeus, 1758)	Nicobar Pigeon
<i>Ptilinopus regina</i> Swainson, 1825	Rose-crowned Fruit-Dove
<i>P. r. ewingii</i> Gould, 1842	
<i>Ducula bicolor</i> (Scopoli, 1786)	Pied Imperial Pigeon
<i>D. b. spilorrhoa</i> (G. R. Gray, 1858)	
	Torres Strait Pigeon

<sup>14</sup> C. antarctica often treated as conspecific with Great Skua C. skua and along with C. maccormicki merged into *Stercorarius*.

<sup>15</sup> Judging from measurements in *Handbook of Western Australian Birds* another subspecies may occur in far north-west.

<sup>16</sup> Specimens from Perth show characters of two subspecies: *S. c. chinensis* and *S. c. tigrina*.

<sup>17</sup> There is considerable geographic variation within this species with white-bellied, red-bellied and intermediate populations, making it difficult to recognise subspecies. See Johnstone (1981, *Rec. West. Aust. Mus.* 9: 49).

<sup>18</sup> Forming a superspecies with Chestnut-quilled Rock Pigeon *P. rufipennis* of Arnhem Land.

### Order PSITTACIFORMES

#### Family PSITTACIDAE

<i>Calyptorhynchus banksii</i> (Latham, 1790)	Red-tailed Black Cockatoo	<i>Polytelis alexandrae</i> Gould, 1863	Princess Parrot
<i>C. b. macrorhynchus</i> Gould, 1843		<sup>23</sup> <i>Platycercus zonarius</i> (Shaw, 1805)	Australian Ringneck (Ring-necked Parrot)
<i>C. b. samueli</i> Mathews, 1917		<i>P. z. zonarius</i> (Shaw, 1805)	Twenty-eight Parrot
<i>C. b. naso</i> Gould, 1837		<i>P. z. semitorquatus</i> (Quoy and Gaimard, 1830)	Twenty-eight Parrot
Forest Red-tailed Black Cockatoo		<sup>24</sup> <i>Platycercus spurius</i> (Kuhl, 1820)	Red-capped Parrot
<i>Calyptorhynchus latirostris</i> Carnaby, 1948	Carnaby's Cockatoo	<i>Platycercus venustus</i> (Kuhl, 1820)	Northern Rosella
<i>Calyptorhynchus baudinii</i> Lear, 1832	Baudin's Cockatoo	<i>Platycercus icterotis</i> (Kuhl, 1820)	Western Rosella
<sup>19</sup> <i>Cacatua roseicapilla</i>	Galah	<i>P. i. icterotis</i> (Kuhl, 1820)	
<i>C. r. roseicapilla</i> Vieillot, 1817		<i>P. i. xanthogenys</i> Salvadori, 1891	
<i>C. r. assimilis</i> (Mathews, 1912)		<i>Platycercus haematogaster</i> Gould, 1838	Blue Bonnet
<i>Cacatua tenuirostris</i> (Kuhl, 1820)	Eastern Long-billed Corella	<i>P. h. narethae</i> (H. L. White, 1921)	
<i>Cacatua pastinator</i> (Gould, 1841)	Western Long-billed Corella	<i>Platycercus varius</i> (Clark, 1910)	Mulga Parrot
<i>C. p. pastinator</i>	Muir's Corella	<i>Neophema elegans</i> (Gould, 1837)	Elegant Parrot
<sup>20</sup> <i>C. p. butleri</i> Ford, 1987	Butler's Corella	<i>Neophema petrophila</i> (Gould, 1841)	Rock Parrot
<i>Cacatua sanguinea</i> Gould, 1843	Little Corella	<i>Neophema splendida</i> (Gould, 1841)	
<i>C. s. sanguinea</i> Gould, 1843			Scarlet-chested Parrot
<i>C. s. westralsis</i> (Mathews, 1917)		<i>Neophema bourkii</i> (Gould, 1841)	Bourke's Parrot
<i>Cacatua leadbeateri</i> (Vigors, 1831)	Major Mitchell's Cockatoo	<i>Melopsittacus undulatus</i> (Shaw, 1805)	Budgerigar
<i>Cacatua galerita</i> (Latham, 1790)	Sulphur-crested Cockatoo	<i>Pezoporus wallicus</i> (Kerr, 1792)	Ground Parrot
<i>C. g. galerita</i> (Latham, 1790)		<i>P. w. flaviventris</i> North, 1911	
<i>C. g. fitzroyi</i> (Mathews, 1912)		<i>Pezoporus occidentalis</i> (Gould, 1861)	Night Parrot
<i>Nymphicus hollandicus</i> (Kerr, 1792)	Cockatiel		
<i>Trichoglossus haematocephalus</i> (Linnaeus, 1771)	Rainbow Lorikeet		
<i>T. h. rubritorquis</i> Vigors and Horsfield, 1827	Red-collared Lorikeet		
<i>T. h. moluccanus</i> (Gmelin, 1788)			
<i>Trichoglossus versicolor</i> Lear, 1831	Varied Lorikeet		
<i>Glossopsitta concinna</i> (Shaw, 1791)	Musk Lorikeet		
<i>Glossopsitta porphyrocephala</i> (Dietrichsen, 1837)	Purple-crowned Lorikeet		
<sup>21</sup> <i>Aprosmictus erythropterus</i> (Gmelin, 1788)	Red-winged Parrot		
<sup>22</sup> <i>Polytelis anthopeplus</i> (Lear, 1831)	Regent Parrot		
<i>P. a. anthopeplus</i>			

<sup>19</sup> The type locality of the Galah remains an enigma. The species was described by Vieillot in 1817 as *Cacatua roseicapilla* from a specimen collected during the Baudin Expedition 1800–1803 labelled 'dans les Indes'. Mathews (1912) *Novit. Zool.* 18: determined this location to be New South Wales, however Schodde (1988) *Canberra Ed Notes* 13.4 believed the type to be from Western Australia probably from the Shark Bay area. I doubt however that this species occurred in the Shark Bay region prior to the 1920s, furthermore I cannot match photographs of the type with any Western Australian specimen. It is possible the type was collected in South Australia by De Freycinet however until the type locality is resolved *assimilis* is referable to the mid Western population.

<sup>20</sup> The provenance and subspecific identity of the type specimen of *Licmetis tenuirostris derbyi* Mathews 1916, are in doubt (most probably an immature *C. p. pastinator*) and until resolved the most appropriate name for the mid Western Australian (wheatbelt) population is *Cacatua pastinator butleri* Ford 1987.

<sup>21</sup> Forming a super species with *A. jonquilleus* of Wetar and Timor.

<sup>22</sup> The type locality of the Regent Parrot is uncertain and the type specimens appear to have been lost. The names *Palaeornis anthopeplus* Lear 1831 and *P. melanura* Lear 1832 were based on a female and male respectively, published on plates in his *Illustrations of the family Psittacidae, or Parrots* 1830–1832, without description or source. Mathews (1912) presumed the type locality was New South Wales and named *Polytelis anthopeplus westralis* from south-west Australia. Schodde 1993 (*Bull. Br. Ornithol. Club* 113: 44–47) noted that Lear's plate matched best with the western form and designated a neotype from Bolgart, Western Australia and provided the name *P. a. monachoides* for the eastern Australian population.

<sup>23</sup> The sole member of the subgenus *Barnardius*.

<sup>24</sup> Sole member of the subgenus *Purpureicephalus*.

<sup>25</sup> Sometimes treated as a full species.

### Order CUCULIFORMES

#### Family CUCULIDAE

<i>Cuculus saturatus</i> Blyth, 1843	Oriental Cuckoo
<sup>25</sup> <i>C. s. optatus</i> Gould, 1845	
<i>Cuculus pallidus</i> (Latham, 1802)	Pallid Cuckoo
<i>Cacomantis flabelliformis</i> (Latham, 1802)	Fan-tailed Cuckoo
<i>C. f. flabelliformis</i> (Latham, 1802)	
<i>Cacomantis variolosus</i> (Vigors and Horsfield, 1827)	Brush Cuckoo
<i>C. v. variolosus</i> (Vigors and Horsfield, 1827)	
<i>Chrysococcyx osculans</i> (Gould, 1847)	Black-eared Cuckoo
<i>Chrysococcyx basalis</i> (Horsfield, 1821)	Horsfield's Bronze Cuckoo
<i>Chrysococcyx lucidus</i> (Gmelin, 1788)	Shining Bronze Cuckoo
<i>C. l. plagus</i> (Latham, 1802)	

<i>Chrysococcyx minutillus</i> Gould, 1859	Little Bronze Cuckoo
<i>C. m. minutillus</i> Gould, 1859	
<i>Eudynamys scolopacea</i> (Linnaeus, 1758)	Common Koel
<i>E. s. cyanocephala</i> (Latham, 1802)	
<i>Scythrops novaehollandiae</i> Latham, 1790	Channel-billed Cuckoo
<i>S. n. novaehollandiae</i> Latham, 1790	

### Family CENTROPODIDAE

<i>Centropus phasianinus</i> (Latham, 1802)	Pheasant Coucal
<i>C. p. melanurus</i> Gould, 1847	
<i>C. p. highami</i> (Mathews, 1922)	

### Order STRIGIFORMES

#### Family STRIGIDAE

<i>Ninox rufa</i> (Gould, 1846)	Rufous Owl
<i>N. r. rufa</i> (Gould, 1846)	
<i>Ninox connivens</i> (Latham, 1802)	Barking Owl
<i>N. c. connivens</i> (Latham, 1802)	
<i>Ninox novaeseelandiae</i> (Gmelin, 1788)	Boobook Owl
<i>N. n. boobook</i> (Latham, 1802)	
<i>Ninox scutulata</i> (Raffles, 1822)	Brown Hawk-Owl
<i>N. s. japonica</i> (Temminck and Schlegel, 1844)	

#### Family TYTONIDAE

<i>Tyto novaehollandiae</i> (Stephens, 1826)	Masked Owl
<i>T. n. novaehollandiae</i> (Stephens, 1826)	
<i>T. n. kimberli</i> Mathews, 1912	
<i>Tyto alba</i> (Scopoli, 1796)	Barn Owl
<i>T. a. delicatula</i> (Gould, 1837)	
<i>Tyto capensis</i> (Smith, 1834)	Grass Owl
<i>T. c. longimembris</i> (Jerdon, 1839)	

### Order CAPRIMULGIFORMES

#### Family PODARGIDAE

<i>Podargus strigoides</i> (Latham, 1802)	Tawny Frogmouth
<i>P. s. phalaenoides</i> Gould, 1840	
<i>P. s. brachypterus</i> Gould, 1841	

#### Family CAPRIMULGIDAE

<i>Eurostopodus argus</i> Hartert, 1892	Spotted Nightjar
---	------------------

### Family AEGOTHELIDAE

<i>Aegotheles cristatus</i> (Shaw, 1790)	Australian Owlet-nightjar
<i>A. c. cristatus</i> (Shaw, 1790)	
<i>A. c. leucogaster</i> Gould, 1844	

### Order APODIFORMES

#### Family APODIDAE

<i>Hirundapus caudacutus</i> (Latham, 1802)	White-throated Needletail
<i>H. c. caudacutus</i> (Latham, 1802)	
<i>Apus pacificus</i> (Latham, 1802)	Fork-tailed Swift
<i>A. p. pacificus</i> (Latham, 1802)	
<sup>26</sup> <i>Apus nipalensis</i> (Hodgson, 1836)	House Swift

### Order CORACIIFORMES

#### Family ALCEDINIDAE

<i>Alcedo azurea</i> Latham, 1802	Azure Kingfisher
<i>A. a. ruficollaris</i> (Bankier, 1841)	

#### Family HALCYONIDAE

<i>Dacelo novaeguineae</i> (Hermann, 1783)	Laughing Kookaburra
<i>D. n. novaeguineae</i> (Hermann, 1783)	
<i>Dacelo leachii</i> Vigors and Horsfield, 1826	Blue-winged Kookaburra
<i>D. l. leachii</i> Vigors and Horsfield, 1826	
<i>Todiramphus pyrrhopygia</i> (Gould, 1841)	Red-backed Kingfisher
<i>Todiramphus sanctus</i> (Vigors and Horsfield, 1827)	Sacred Kingfisher
<i>T. s. sanctus</i> (Vigors and Horsfield, 1827)	
<i>Todiramphus chloris</i> (Boddaert, 1783)	Collared Kingfisher
<i>T. c. sordida</i> Gould, 1842	
<i>T. c. pilbara</i> Johnstone, 1983	

#### Family MEROPIDAE

<i>Merops ornatus</i> Latham, 1802	Rainbow Bee-eater
------------------------------------	-------------------

#### Family CORACIIDAE

<i>Eurystomus orientalis</i> (Linnaeus, 1766)	Dollarbird
<i>E. o. pacificus</i> (Latham, 1802)	

<sup>26</sup> Sometimes treated as conspecific with the Little Swift *Apus affinis*.

## Order PASSERIFORMES

### Family PITTIDAE

*Pitta moluccensis* (Müller, 1776) Blue-winged Pitta  
<sup>27</sup>*Pitta iris* Gould, 1842 Rainbow Pitta  
*P. i. johnstoneiana* Schodde and Mason 1999

### Family ATRICORNITHIDAE

*Atrichornis clamosus* (Gould, 1844) Noisy Scrub-bird

### Family CLIMACTERIDAE

*Climacteris affinis* Blyth, 1864  
     White-browed Treecreeper  
     *C. a. superciliosa* North, 1895  
*Climacteris melanura* Gould, 1843  
     Black-tailed Treecreeper  
     *C. m. melanura* Gould, 1843  
     *C. m. wellsi* Ogilvie-Grant, 1909  
<sup>28</sup>*Climacteris rufa* Gould, 1841 Rufous Treecreeper

### Family MALURIDAE

*Malurus coronatus* Gould, 1858  
     Purple-crowned Fairy-wren  
     *M. c. coronatus* Gould, 1858  
*Malurus splendens* (Quoy and Gaimard, 1830)  
     Splendid Fairy-wren  
     *M. s. splendens* (Quoy and Gaimard, 1830)  
     *M. s. musgravi* Mathews, 1922  
*Malurus lamberti* Vigors and Horsfield, 1827  
     Variegated Fairy-wren  
     *M. l. assimilis* North, 1901  
     *M. l. rogersi* Mathews, 1912  
     *M. l. bernieri* Ogilvie-Grant, 1909  
*Malurus pulcherrimus* Gould, 1844  
     Blue-breasted Fairy-wren  
*Malurus elegans* Gould, 1837 Red-winged Fairy-wren  
*Malurus leucopterus* Dumont, 1824  
     White-winged Fairy-wren  
     *M. l. leucopterus* Dumont, 1824  
     *M. l. leuconotus* Gould, 1865  
     *M. l. edouardi* Campbell, 1901  
*Malurus melanocephalus* (Latham, 1802)  
     Red-backed Fairy-wren  
     *M. m. cruentatus* Gould, 1839  
*Stipiturus malachurus* (Shaw, 1798)  
     Southern Emu-wren  
     *S. m. westernensis* Campbell, 1912  
     *S. m. hartogi* Carter, 1916  
*Stipiturus ruficeps* Campbell, 1899  
     Rufous-crowned Emu-wren

*S. r. ruficeps* Campbell, 1899  
*Amytornis striatus* (Gould, 1840) Striated Grasswren  
     *A. s. striatus* (Gould, 1840)  
     *A. s. whitei* Mathews, 1910  
*Amytornis textilis* (Dumont, 1824)  
     Thick-billed Grasswren  
     *A. t. textilis* (Dumont, 1824)  
*Amytornis purnelli* (Mathews, 1914)  
     Dusky Grasswren  
     *A. p. purnelli* (Mathews, 1914)  
*Amytornis housei* (Milligan, 1902) Black Grasswren

### Family PARDALOTIDAE

*Pardalotus punctatus* (Shaw, 1792) Spotted Pardalote  
     *P. p. punctatus* (Shaw, 1792)  
<sup>29</sup>*P. p. xanthopyge* McCoy, 1866  
     Yellow-rumped Pardalote  
*Pardalotus rubricatus* Gould, 1838  
     Red-browed Pardalote  
*Pardalotus striatus* (Gmelin, 1789) Striated Pardalote  
     *P. s. uropygialis* Gould, 1840  
     *P. s. murchisoni* Mathews, 1912  
     *P. s. westralsiensis* Mathews, 1912

### Family DASYORNITHIDAE

*Dasyornis longirostris* Gould, 1841  
     Western Bristlebird  
*Dasyornis broadbenti* (McCoy, 1867)  
     Rufous Bristlebird  
     *D. b. litoralis* (Milligan, 1902)

### Family ACANTHIZIDAE

*Sericornis frontalis* (Vigors and Horsfield, 1827)  
     White-browed Scrubwren  
     *S. f. maculatus* Gould, 1847  
     *S. f. balstoni* Ogilvie-Grant, 1909  
*Hylacola cauta* Gould, 1843  
     Shy Groundwren (Shy Heathwren)  
<sup>30</sup>*Calamanthus campestris* (Gould, 1841)  
     Rufous Fieldwren  
*Pyrrholaemus brunneus* Gould, 1841  
     Redthroat  
*Smicromis brevirostris* (Gould, 1838)  
     Weebill  
*Gerygone olivacea* (Gould, 1838)  
     White-throated Gerygone  
     *G. o. rogersi* Mathews, 1911  
*Gerygone fusca* (Gould, 1838)  
     Western Gerygone  
     *G. f. fusca* (Gould, 1838)  
<sup>31</sup>*Gerygone (fusca) mungi* Mathews, 1912  
     Desert Gerygone

<sup>27</sup> Forming a superspecies with *P. versicolor* of eastern Australia and *P. elegans* of Indonesia.

<sup>28</sup> Forming a superspecies with the Brown Treecreeper *C. picumnus* of eastern Australia.

<sup>29</sup> Tentatively treated as conspecific with *P. punctatus* based on evidence of hybridisation in southeast Australia (Schodde and Mason, 1999).

<sup>30</sup> Previously treated as a subspecies of Striated Fieldwren *Calamanthus fuliginosus*.

<sup>31</sup> Closely related to the Western Gerygone and possibly only a well marked subspecies.

<i>Gerygone levigaster</i> Gould, 1843	Mangrove Gerygone	<i>Lichenostomus ornatus</i> (Gould, 1838)
<i>G. l. levigaster</i> Gould, 1843		Yellow-plumed Honeyeater
<i>Gerygone tenebrosa</i> (Hall, 1901)	Dusky Gerygone	<i>Lichenostomus penicillatus</i> Gould, 1837
<i>Gerygone magnirostris</i> Gould, 1843	Large-billed Gerygone	White-plumed Honeyeater
<i>G. m. magnirostris</i> Gould, 1843		<i>Lichenostomus flavescens</i> (Gould, 1840)
<i>Gerygone chloronotus</i> Gould, 1843	Green-backed Gerygone	Yellow-tinted Honeyeater
<i>G. c. chloronotus</i> Gould, 1843		<i>L. f. flavescentis</i> (Gould, 1840)
<sup>32</sup> <i>Acanthiza apicalis</i> Gould, 1847	Broad-tailed Thornbill (Inland Thornbill)	<i>Lichenostomus cratitius</i> (Gould, 1841)
<i>Acanthiza robustirostris</i> Milligan, 1903	Slaty-backed Thornbill	<i>Lichenostomus leucotis</i> (Latham, 1802)
<i>Acanthiza uropygialis</i> Gould, 1838	Chestnut-rumped Thornbill	White-eared Honeyeater
<i>Acanthiza iredalei</i> Mathews, 1911	Samphire Thornbill (Slender-billed Thornbill)	<i>L. l. novaenorciae</i> (Milligan, 1904)
<i>A. i. iredalei</i> Mathews, 1911		<i>Lichenostomus unicolor</i> (Gould, 1843)
<i>Acanthiza inornata</i> Gould, 1841	Western Thornbill	White-gaped Honeyeater
<i>Acanthiza chrysorrhoa</i> (Quoy and Gaimard, 1830)	Yellow-rumped Thornbill	<i>L. u. unicolor</i> (Gould, 1843)
<i>Aphelocephala leucopsis</i> (Gould, 1841)	Southern Whiteface	<i>Melithreptus brevirostris</i> (Vigors and Horsfield, 1827)
<i>A. l. leucopsis</i> (Gould, 1841)		Brown-headed Honeyeater
<i>A. l. castaneiventris</i> (Milligan, 1903)		<i>M. b. leucogenys</i> Milligan, 1903
<i>Aphelocephala nigricincta</i> (North, 1895)	Banded Whiteface	<sup>37</sup> <i>Melithreptus chloropsis</i> Gould, 1848
		Western White-naped Honeyeater
		<i>Melithreptus albogularis</i> Gould, 1848
		White-throated Honeyeater
		<i>Melithreptus gularis</i> (Gould, 1837)
		Black-chinned Honeyeater
		<i>M. g. laetior</i> Gould, 1875
		<i>Melithreptus cyanotis</i> (Latham, 1802)
		Blue-faced Honeyeater
		<i>M. c. albipennis</i> (Gould, 1840)
		<i>Philemon citreogularis</i> (Gould, 1837) Little Friarbird
		<i>P. c. citreogularis</i> (Gould, 1837)
		<i>Philemon argenticeps</i> (Gould, 1840)
		Silver-crowned Friarbird
		<i>P. a. argenticeps</i> (Gould, 1840)
		<sup>38</sup> <i>Phylidonyris novaehollandiae</i> (Latham, 1790)
		New Holland Honeyeater
		<i>Phylidonyris nigra</i> (Bechstein, 1811)
		White-cheeked Honeyeater
		<i>P. n. gouldii</i> (Schlegel, 1872)
		<i>Phylidonyris albifrons</i> (Gould, 1841)
		White-fronted Honeyeater
		<i>Phylidonyris melanops</i> (Latham, 1802)
		Tawny-crowned Honeyeater
		<i>Acanthorhynchus superciliosus</i> Gould, 1837
		Western Spinebill
		<i>Cissomela pectoralis</i> (Gould, 1841)
		Banded Honeyeater
		<i>Ramsayornis fasciatus</i> (Gould, 1843)
		Bar-breasted Honeyeater

<sup>32</sup> Forming a superspecies with the Brown Thornbill *A. pusilla* of Tasmania and wetter parts of eastern Australia.

<sup>33</sup> Schodde (1975) split *Meliphaga* into three genera namely: *Lichenostomus* with 18 species in Australia, 4 in New Guinea (2 shared), 9 species occurring in Western Australia – *virescens*, *keartlandi*, *plumulus*, *ornatus*, *penicillatus*, *flavescens*, *cratitius*, *leucotis* and *unicolor*; *Meliphaga* with 9 species in New Guinea, 4 in Australia (1 shared), only one *albilineata* occurring in Western Australia; and *Xanthotis* with 2 species in Australia and 2 in New Guinea (1 shared). While this arrangement is followed here the composition of genera and species within the Meliphagidae are I believe, still far from resolved.

<sup>34</sup> Forming a superspecies with *M. dammermani*, *M. kuehni* and possibly *M. vulnerata*, all of the Lesser Sundas.

<sup>35</sup> An isolated species, not certainly close to Pied Honeyeater.

<sup>36</sup> Forming a superspecies with the Mangrove Honeyeater *L. versicolor* of eastern Australia and New Guinea.

<sup>37</sup> Forming a superspecies with *Melithreptus lunatus* of eastern Australia.

<sup>38</sup> Comprising with the White-cheeked Honeyeater *P. nigra* the subgenus *Meliornis*.

<i>Conopophila rufogularis</i> (Gould, 1843)	Rufous-throated Honeyeater
<i>Lacustoica whitei</i> North, 1910	Grey Honeyeater
<i>Manorina flavigula</i> (Gould, 1840)	Yellow-throated Miner
<i>Acanthagenys rufogularis</i> Gould, 1838	Spiny-cheeked Honeyeater
<sup>39</sup> <i>Anthochaera lunulata</i> Gould, 1838	Western Little Wattlebird
<i>Anthochaera carunculata</i> (White, 1790)	Red Wattlebird
<i>Epthianura albifrons</i> (Jardine and Selby, 1828)	White-fronted Chat
<i>Epthianura crocea</i> Castelnau and Ramsay, 1877	Yellow Chat
<i>Epthianura aurifrons</i> Gould, 1838	Orange Chat
<i>Epthianura tricolor</i> Gould, 1841	Crimson Chat

#### Family PETROICIDAE

<i>Microeca flavigaster</i> Gould, 1843	Lemon-breasted Flycatcher
<i>M. f. flavigaster</i> Gould, 1843	
<i>M. f. tormenti</i> Mathews, 1916	Kimberley Flycatcher
<i>Microeca fascinans</i> (Latham, 1802)	Jacky Winter
<i>M. f. fascinans</i> (Latham, 1802)	
<i>M. f. assimilis</i> Gould, 1841	
<i>Petroica multicolor</i> (Gmelin, 1789)	Scarlet Robin
<i>P. m. campbelli</i> Sharpe, 1898	
<i>Petroica goodenovii</i> (Vigors and Horsfield, 1827)	Red-capped Robin
<i>Petroica cucullata</i> (Latham, 1802)	Hooded Robin
<i>Eopsaltria australis</i> (Shaw, 1790)	Yellow Robin
<sup>40</sup> <i>E. a. griseogularis</i> Gould, 1838	Western Yellow Robin
<i>Eopsaltria georgiana</i> (Quoy and Gaimard, 1830)	White-breasted Robin
<i>Eopsaltria pulverulenta</i> (Bonaparte, 1850)	Mangrove Robin
<i>Poecilodryas superciliosa</i> (Gould, 1847)	White-browed Robin
<i>P. s. cerviniventris</i> (Gould, 1858)	Buff-sided Robin
<i>Drymodes brunneopygia</i> Gould, 1841	Southern Scrub-robin

#### Family POMATOSTOMIDAE

<i>Pomatostomus temporalis</i> (Vigors and Horsfield, 1827)	Grey-crowned Babbler
<i>P. t. rubeculus</i> Gould, 1839	
<i>Pomatostomus superciliosus</i> (Vigors and Horsfield, 1827)	White-browed Babbler

#### Family CINCLOSOMATIDAE

<i>Psophodes nigrogularis</i> Gould, 1844	Western Whipbird
<i>P. n. nigrogularis</i> Gould, 1844	
<i>P. n. oberon</i> Schodde and Mason, 1991	
<i>Psophodes occidentalis</i> (Mathews, 1912)	Western Wedgebill (Chiming Wedgebill)
<i>Cinclosoma castanotus</i> Gould, 1840	Chestnut Quail-thrush
<sup>41</sup> <i>Cinclosoma alisteri</i> Mathews, 1910	Nullarbor Quail-thrush
<i>Cinclosoma castaneothorax</i> Gould, 1848	Chestnut-breasted Quail-thrush
<i>C. c. marginatum</i> Sharpe, 1883	

#### Family NEOSITTIDAE

<i>Daphoenositta chrysoptera</i> (Latham, 1802)	Varied Sittella
<i>D. c. pileata</i> (Gould, 1838)	
<i>D. c. leucoptera</i> (Gould, 1840)	

#### Family PACHYCEPHALIDAE

<i>Falcunculus frontatus</i> (Latham, 1802)	Crested Shrike-tit
<i>F. f. leucogaster</i> Gould, 1838	
<i>F. f. whitei</i> Campbell, 1910	Northern Shrike-tit
<i>Oreoica gutturalis</i> (Vigors and Horsfield, 1827)	Crested Bellbird
<i>Pachycephala melanura</i> Gould, 1843	Mangrove Golden Whistler
<i>P. m. melanura</i> Gould, 1843	
<i>P. m. spinicauda</i> (Pucheran, 1853)	
<i>Pachycephala pectoralis</i> (Latham, 1802)	Golden Whistler
<i>P. p. fuliginosa</i> Vigors and Horsfield, 1827	
<i>Pachycephala rufiventris</i> (Latham, 1802)	Rufous Whistler
<i>P. r. rufiventris</i> (Latham, 1802)	
<i>Pachycephala lanioides</i> Gould, 1840	White-breasted Whistler
<i>Pachycephala inornata</i> Gould, 1841	Gilbert's Whistler
<i>Colluricincla megarhyncha</i> (Quoy and Gaimard, 1830)	Little Shrike-thrush
<i>C. m. parvula</i> Gould, 1845	
<i>Colluricincla woodwardi</i> Hartert, 1905	Sandstone Shrike-thrush
<i>Colluricincla harmonica</i> (Latham, 1802)	Grey Shrike-thrush
<i>C. h. brunnea</i> Gould, 1841	
<i>C. h. kolichisi</i> Ford, 1987	
<i>C. h. rufiventris</i> Gould, 1841	

<sup>39</sup> The eastern and western populations of *Anthochaera chrysoptera* (*chrysoptera* and *lunulata* respectively) were previously treated as conspecific.

<sup>40</sup> Sometimes treated as a full species *E. griseogularis*.

<sup>41</sup> Sometimes treated as conspecific with Cinnamon Quail-thrush *C. cinnamomeum* but the two forms are best treated as allospecies.

### Family DICRURIDAE

<i>Monarcha melanopsis</i> (Vieillot, 1818)	
	Black-faced Monarch
<i>Myiagra ruficollis</i> (Vieillot, 1818)	
	Broad-billed Flycatcher
<i>M. r. mimikae</i> Ogilvie-Grant, 1911	
<i>Myiagra rubecula</i> (Latham, 1802)	Leaden Flycatcher
<i>M. r. concinna</i> Gould, 1848	
<i>Myiagra cyanoleuca</i> (Vieillot, 1818)	Satin Flycatcher
<i>Myiagra inquieta</i> (Latham, 1802)	Restless Flycatcher
<i>M. i. inquieta</i> (Latham, 1802)	
<i>M. i. nana</i> Gould, 1870	
<i>Myiagra alecto</i> (Temminck, 1827)	Shining Flycatcher
<i>M. a. melvillensis</i> (Mathews, 1912)	
<sup>42</sup> <i>Rhipidura dryas</i> Gould, 1842	
	Wood Fantail (Rufous Faintail)
<i>R. d. dryas</i> Gould, 1842	
<sup>43</sup> <i>Rhipidura phasiana</i> De Vis, 1884	
	Mangrove Grey Fantail
<i>Rhipidura fuliginosa</i> Sparrman, 1787	Grey Fantail
<i>R. f. preissi</i> Cabanis, 1851	
<i>R. f. alisteri</i> Mathews, 1911	
<sup>44</sup> <i>Rhipidura (fuliginosa) albicauda</i> North, 1895	
	White-tailed Fantail
<i>Rhipidura leucophrys</i> (Latham, 1802)	Willie Wagtail
<i>R. l. leucophrys</i> (Latham, 1802)	
<i>Rhipidura rufiventris</i> (Vieillot, 1818)	Northern Fantail
<i>R. r. isura</i> Gould, 1841	
<i>Grallina cyanoleuca</i> (Latham, 1802)	Magpie-lark
<i>Dicrurus bracteatus</i> (Gould, 1842)	Spangled Drongo
<i>D. b. bracteatus</i> (Gould, 1842)	

### Family CAMPEPHAGIDAE

<i>Coracina novaehollandiae</i> (Gmelin, 1789)	
	Black-faced Cuckoo-shrike
<i>C. n. novaehollandiae</i> (Gmelin, 1789)	
<i>C. n. subpallida</i> Mathews, 1912	
<i>Coracina papuensis</i> (Gmelin, 1788)	Little Cuckoo-shrike (White-bellied Cuckoo-shrike)
<i>C. p. hypoleuca</i> (Gould, 1848)	
	White-breasted Cuckoo-shrike
<i>Coracina tenuirostris</i> (Jardine, 1831)	Cicadabird
<i>C. t. melvillensis</i> Mathews, 1912	
<i>Coracina maxima</i> (Rüppell, 1839)	Ground Cuckoo-shrike
<sup>45</sup> <i>Lalage tricolor</i> (Swainson, 1825)	White-winged Triller
<i>Lalage leucomela</i> (Vigors and Horsfield, 1827)	Varied Triller
<i>L. l. rufiventris</i> (Gray, 1846)	

### Family ORIOLIDAE

<i>Oriolus flavocinctus</i> (King, 1826)	Yellow Oriole
<i>Oriolus sagittatus</i> (Latham, 1802)	Olive-backed Oriole
<i>Sphecotheres viridis</i> Vieillot, 1816	Figbird
<i>S. v. flaviventris</i> Gould, 1850	Yellow Figbird

### Family ARTAMIDAE

<i>Artamus leucorynchus</i> (Linnaeus, 1771)	
	White-breasted Woodswallow
<i>A. l. leucopygialis</i> Gould, 1842	
<i>Artamus personatus</i> (Gould, 1841)	
	Masked Woodswallow
<i>Artamus superciliosus</i> (Gould, 1837)	
	White-browed Woodswallow
<i>Artamus cinereus</i> Vieillot, 1817	
	Black-faced Woodswallow
<i>A. c. melanops</i> Gould, 1865	
<i>Artamus cyanopterus</i> (Latham, 1802)	
	Dusky Woodswallow
<i>Artamus minor</i> Vieillot, 1817	
	Little Woodswallow

### Family CRACTICIDAE

<i>Cracticus quoyi</i> (Lesson and Garnot, 1827)	
	Black Butcherbird
<i>C. q. spaldingi</i> Ramsay, 1878	
<i>Cracticus torquatus</i> (Latham, 1802)	Grey Butcherbird
<i>C. t. torquatus</i> (Latham, 1802)	
<i>C. t. latens</i> Ford, 1979	
	Silver-backed Butcherbird
<i>Cracticus nigrogularis</i> (Gould, 1837)	Pied Butcherbird
<sup>46</sup> <i>Cracticus tibicen</i> (Latham, 1802)	Australian Magpie
<i>C. t. tibicen</i> (Latham, 1802)	
<i>C. t. dorsalis</i> (Campbell, 1894)	
<i>Strepera versicolor</i> (Latham, 1802)	Grey Currawong
<i>S. v. plumbea</i> Gould, 1846	

### Family CORVIDAE

<i>Corvus splendens</i> Vieillot, 1817	House Crow]
<i>Corvus orru</i> Bonaparte, 1850	Torresian Crow
<i>C. o. ceciliae</i> Mathews, 1912	Western Crow
<i>Corvus bennetti</i> North, 1901	Little Crow
<i>Corvus coronoides</i> Vigors and Horsfield, 1827	Australian Raven
<i>C. c. perplexus</i> Mathews, 1912	

<sup>42</sup> Most recent workers have treated *Rhipidura dryas* as a subspecies of the Rufous Faintail *R. rufifrons*. However *rufifrons* and *dryas* do not intergrade in north Queensland and are treated here as allospecies in view of their marked differences in habitat and morphology. Both *R. dryas* and *R. rufifrons* forming a superspecies with several east Indonesian species.

<sup>43</sup> Forming a superspecies with the Grey Fantail and possibly the White-tailed Fantail.

<sup>44</sup> Closely related to the Grey Fantail and possibly only a well marked subspecies.

<sup>45</sup> Forming a superspecies with *L. sueurii* of Indonesia.

<sup>46</sup> Formerly in genus *Gymnorhina*.

### Family PTILONORHYNCHIDAE

- Ptilonorhynchus maculatus* (Gould, 1837) Spotted Bowerbird  
<sup>47</sup>*P. m. guttatus* (Gould, 1862) Western Bowerbird  
*Ptilonorhynchus nuchalis* Jardine and Selby, 1830 Great Bowerbird  
*P. n. nuchalis* Jardine and Selby, 1830

### Family MUSCICAPIDAE

- Turdus merula* (Linnaeus, 1758) Eurasian Blackbird (Common Blackbird)  
[T. m. merula (Linnaeus, 1758)]  
*Muscicapa sibirica* Gmelin, 1789 Sooty Flycatcher (Dark-sided Flycatcher)  
[M. s. sibirica Gmelin, 1789]  
*Ficedula narcissina* (Temminck, 1835) Narcissus Flycatcher  
F. n. narcissina (Temminck, 1835)  
*Ficedula cyanomelana* (Temminck, 1829) Blue and White Flycatcher  
*F. c. cyanomelana* (Temminck, 1829)

### Family STURNIDAE

- Sturnus vulgaris* Linnaeus, 1758 Common Starling  
*S. v. vulgaris* Linnaeus, 1758  
*Acridotheres tristis* (Linnaeus, 1766) Common Myna  
[A. t. tristis (Linnaeus, 1766)]

### Family HIRUNDINIDAE

- Cheramoeca leucosternus* (Gould, 1841) White-backed Swallow  
*Hirundo rustica* Linnaeus, 1758 Barn Swallow  
*H. r. gutturalis* Scopoli, 1786  
*Hirundo tahitica* Gmelin, 1789 Pacific Swallow  
*Hirundo neoxena* Gould, 1842 Welcome Swallow  
*Hirundo daurica* Laxmann, 1769 Red-rumped Swallow  
*H. d. japonica* Temminck and Schlegel 1847  
*Hirundo nigricans* Vieillot, 1817 Tree Martin  
*H. n. nigricans* Vieillot, 1817  
*Hirundo ariel* (Gould, 1842) Fairy Martin

### Family PYCNONOTIDAE

- Pycnonotus jocosus* (Linnaeus, 1758) Red-whiskered Bulbul  
[P. j. jocosus (Linnaeus, 1758)]

### Family ZOSTEROPIDAE

- Zosterops luteus* Gould, 1843 Yellow White-eye  
*Zosterops lateralis* (Latham, 1802) Grey-breasted White-eye (Silvereye)  
*Z. l. gouldi* Bonaparte, 1850

### Family SYLVIIDAE

- Acrocephalus australis* (Gould, 1838) Australian Reed Warbler  
*A. a. gouldi* (Dubois, 1901)  
*Acrocephalus orientalis* (Temminck and Schlegel, 1847) Oriental Reed Warbler  
*Phylloscopus borealis* (Blasius, 1858) Arctic Warbler  
*Megalurus timoriensis* Wallace, 1864 Tawny Grassbird  
*M. t. alisteri* Mathews, 1912  
*Megalurus gramineus* (Gould, 1845) Little Grassbird  
*M. g. gramineus* (Gould, 1845)  
*Eremiornis carteri* North, 1900 Spinifex-bird  
*Cincloramphus mathewsi* Iredale, 1911 Rufous Songlark  
*Cincloramphus cruralis* (Vigors and Horsfield, 1827) Brown Songlark  
*Cisticola exilis* (Vigors and Horsfield, 1827) Golden-headed Cisticola  
*C. e. exilis* (Vigors and Horsfield, 1827)  
*Cisticola juncidis* (Rafinesque, 1810) Zitting Cisticola  
*C. j. fuscicapilla* Wallace, 1864

### Family ALAUDIDAE

- Mirafra javanica* Horsfield, 1821 Singing Bushlark  
*M. j. horsfieldii* Gould, 1847

### Family DICAEIDAE

- Dicaeum hirundinaceum* (Shaw, 1792) Mistletoebird  
*D. h. hirundinaceum* (Shaw, 1792)

### Family PASSERIDAE

- Passer domesticus* (Linnaeus, 1758) House Sparrow  
*P. d. domesticus* (Linnaeus, 1758)  
*Passer montanus* (Linnaeus, 1758) Eurasian Tree Sparrow  
*Taeniopygia guttata* (Vieillot, 1817) Zebra Finch  
*T. g. castanotis* (Gould, 1837)  
*Taeniopygia bichenovii* (Vigors and Horsfield, 1827) Double-barred Finch

<sup>47</sup> Sometimes treated as a full species *P. guttatus*.

<i>T. b. annulosa</i> (Gould, 1840)	
<i>Poephila acuticauda</i> (Gould, 1840)	Long-tailed Finch
<i>Poephila personata</i> Gould, 1842	Masked Finch
<i>P. p. personata</i> Gould, 1842	
<i>Neochmia phaeton</i> (Hombron and Jacquinot, 1841)	
	Crimson Finch
<i>N. p. phaeton</i> (Hombron and Jacquinot, 1841)	
<i>Neochmia ruficauda</i> (Gould, 1837)	Star Finch
<i>N. r. clarescens</i> (Hartert, 1899)	
<i>Neochmia temporalis</i> (Latham, 1802)	
	Red-browed Finch
<i>N. t. temporalis</i> (Latham, 1802)	
<i>Stagonopleura oculata</i> (Quoy and Gaimard, 1830)	
	Red-eared Firetail
<i>Emblema pictum</i> Gould, 1842	Painted Finch
<i>Lonchura pallida</i> (Wallace, 1863)	Pale-headed Munia
<i>Lonchura flavigrymnina</i> (Gould, 1845)	
	Yellow-rumped Mannikin
<i>Lonchura castaneothorax</i> (Gould, 1837)	
	Chestnut-breasted Mannikin
<i>L. c. castaneothorax</i> (Gould, 1837)	

<i>Heteromunia pectoralis</i> (Gould, 1841)	
	Pictorella Mannikin
<i>Erythrura gouldiae</i> (Gould, 1844)	Gouldian Finch

### Family MOTACILLIDAE

<i>Anthus cervinus</i> (Pallas, 1811)	Red-throated Pipit
* <i>Anthus australis</i> Vieillot, 1818	Australian Pipit
<i>A. a. australis</i> Vieillot, 1818	
<i>Motacilla flava</i> Linnaeus, 1758	Yellow Wagtail
<i>M. f. simillima</i> Hartert, 1905	
<i>M. f. taivana</i> (Swinhoe, 1863)	
<i>Motacilla alba</i> Linnaeus, 1758	White Wagtail
<i>M. a. ocularis</i> Swinhoe, 1860	
<i>M. a. lugens</i> Cloger, 1829	

### Family FRINGILLIDAE

<i>Carduelis carduelis</i> (Linnaeus, 1758)	Goldfinch
	(European Goldfinch)
[ <i>C. c. britannica</i> (Hartert, 1903)]	

\* Previously included in the superspecies *Anthus novaezeelandiae* (Gmelin, 1789).

## INDEX OF FAMILIES

Acanthizidae .....	84
Accipitridae .....	78
Aegothelidae .....	83
Alcedinidae .....	83
Alaudidae .....	88
Anatidae .....	76
Anhingidae .....	77
Apodidae .....	83
Ardeidae .....	78
Artamidae .....	87
Atrichornithidae .....	84
Burhinidae .....	80
Campephagidae .....	87
Caprimulgidae .....	83
Casuariidae .....	75
Centropodidae .....	83
Charadriidae .....	80
Cinclosomatidae .....	86
Ciconiidae .....	78
Climacteridae .....	84
Columbidae .....	81
Coraciidae .....	83
Corvidae .....	87
Cracticidae .....	87
Cuculidae .....	82
Dasyornithidae .....	84
Dicaeidae .....	88
Dicruridae .....	87
Diomedeidae .....	77
Falconidae .....	79
Fregatidae .....	78
Fringillidae .....	89
Glareolidae .....	81
Gruidae .....	79
Haematopodidae .....	80
Halcyonidae .....	83
Hirundinidae .....	88
Hydrobatidae .....	77
Jacanidae .....	80
Laridae .....	81
Maluridae .....	84
Megapodiidae .....	75
Meliphagidae .....	85
Meropidae .....	83
Motacillidae .....	89
Muscicapidae .....	88
Neosittidae .....	86
Oriolidae .....	87
Otididae .....	79
Pachycephalidae .....	86
Pardalotidae .....	84
Passeridae .....	88
Pelecanidae .....	78
Pelecanoididae .....	77
Petroicidae .....	86
Phaethontidae .....	77
Phalacrocoracidae .....	77
Phasianidae .....	75
Pittidae .....	84
Podargidae .....	83
Podicipedidae .....	76
Pomatostomidae .....	86
Procellariidae .....	76
Psittacidae .....	82
Ptilonorhynchidae .....	88
Pycnonotidae .....	88
Rallidae .....	79
Recorvirostridae .....	80
Rostratulidae .....	80
Scolopacidae .....	79
Spheniscidae .....	76
Strigidae .....	83
Sturnidae .....	88
Sulidae .....	77
Sylviidae .....	88
Threskiornithidae .....	78
Turnicidae .....	79
Tytonidae .....	83
Zosteropidae .....	88