

A replacement name for *Lerista concolor* Storr, 1990 (Squamata: Scincidae).

G. M. Shea*

In a recent note, Storr (1990) described as new *Lerista concolor*, from a single specimen from the Maitland River. Unfortunately, *Lerista concolor* Storr is a junior subjective homonym of *Lygosoma (Rhodona) bipes concolor* Werner, 1910, itself a junior subjective synonym (Storr 1971) of *Lerista lineopunctulata* (Duméril and Bibron, 1839), and consequently a replacement name is required while both species remain in *Lerista* (Article 59 of the Code). I propose *Lerista quadrivincula* as a replacement name for *Lerista concolor* Storr. The name, from the Latin *quadr-* (four) and *vinculum* (a chain), alludes to the four lines of dark streaks on the dorsum, one of the diagnostic characters of the species (Storr 1990).

References

- Duméril, A.M.C. and Bibron, G. (1839). *Erpétologie Générale ou Histoire Complète des Reptiles*. Vol. 5. Roret, Paris.
- Storr, G.M. (1971). The genus *Lerista* (Lacertilia: Scincidae) in Western Australia. *J.R. Soc. West. Aust.* **54**: 59-75.
- Storr, G.M. (1990). A new *Lerista* (Lacertilia: Scincidae) from the Pilbara. *Rec. West. Aust. Mus.* **14**: 669-670.
- Werner, F. (1910). Reptilia (Geckonidae und Scincidae). pp. 451-493 in Michaelsen W. and Hartmeyer R. (eds). *Die Fauna Südwest-Australiens*. Vol. 2. Gustav Fischer, Jena.

*Dept. of Veterinary Anatomy, University of Sydney, NSW, 2006.