

Checklist of the Birds of Western Australia

R.E. Johnstone and J.C. Darnell

Western Australian Museum, Perth, Western Australia 6000

August 2017

The area covered by this Western Australian Checklist includes the seas and islands of the adjacent continental shelf, including Ashmore Reef. Refer to a separate Checklist for Christmas and Cocos (Keeling) Islands. Criterion for inclusion of a species or subspecies on the list is, in most cases, supported by tangible evidence i.e. a museum specimen, an archived or published photograph or detailed description, video tape or sound recording.

Amendments to the previous Checklist have been carried out with reference to both global and regional publications/checklists. The prime reference material for global coverage has been the International Ornithological Committee (IOC) World Bird List, *The Clements Checklist of Birds of the World*, the *Illustrated Checklist of the Birds of the World* Volume, 1 (Lynx Edicions, Barcelona), A Checklist of the Birds of Britain, 8th edition, the Checklist of North American Birds and, for regional coverage, *Zoological Catalogue of Australia* volume 37.2 (Columbidae to Coraciidae), *The Directory of Australian Birds*, Passerines and the Working List of Australian Birds (Birdlife Australia).

The advent of molecular investigation into avian taxonomy has required, and still requires, extensive and ongoing revision at all levels - family, generic and specific. This revision to the 'Checklist of the Birds of Western Australia' is a collation of the most recent information/research emanating from such studies, together with the inclusion of newly recorded species.

As a result of the constant stream of publication of new research in many scientific journals, delays of its incorporation into the prime sources listed above, together with the fact that these are upgraded/re-issued at differing intervals and that their authors may hold varying opinions, these prime references, do on occasion differ. Where considered appropriate these latest research findings have been included as has some pre-published material in order to make this Checklist as up to date as possible.

From these comments it can be seen that avian taxonomy is a 'work in progress' and that it is currently a long way from having a definitive world list. Overall, departures from the prime reference material in both global and regional lists (as above) have been minimised. Subject to the general consensus/opinion of other authors, the IOC comments have been accepted relating to the higher taxonomic levels and taxonomic status beyond Australia, but not where the Western Australian Museum collection material, distributional data base etc. shows otherwise.

Changes should only come when there is certainty and consensus and the Western Australian Museum's objective is to maintain the status quo until claimed revisions have been thoroughly analysed and accepted.

A further element in this current revision to the 'Checklist of the Birds of Western Australia' is the number of additional species contained/recorded. Increased public interest in ornithology combined with improved identification knowledge, the increasing number of "bird watchers" having high-technology photographic equipment, and more extensive field excursions (often to remote areas e.g. Ashmore Reef) all have contributed here. Whether all of these newly recorded species are in fact migratory "strays" which have been missed earlier, or whether other factors such as habitat destruction (along migratory routes) or climatic change is involved is of course unknown.

The aim of this checklist is to summarise all available data on systematics and taxonomy, provide brief explanations for changes, incorporate new higher level classifications and recent new generic, specific and subspecific changes. Where necessary, reasons for departing from the current above-named checklists (IOC, Birds Australia etc.) are also given. Where there is doubt on the identification of a subspecies it is included in square brackets. An asterisk * denotes an introduced or naturalised species, not originally native.

We thank Kim Sarti for the development and layout of this checklist document.

Class AVES**Order CASUARIIFORMES****Family DROMAIIDAE Emu**

- ¹*Dromaius novaehollandiae* (Latham, 1790) Emu
Dromaius novaehollandiae novaehollandiae (Latham, 1790)

Order ANSERIFORMES**Family ANSERANATIDAE Magpie Goose**

- ²*Anseranas semipalmata* (Latham, 1798) Magpie Goose (Pied Goose)

Family ANATIDAE Ducks, geese and swans

- Dendrocygna eytoni* (Eyton, 1838) Plumed Whistling Duck
³*Dendrocygna arcuata* (Horsfield, 1824) Wandering Whistling Duck (Chestnut Whistling Duck)
Cereopsis novaehollandiae Latham, 1802 Cape Barren Goose
Cereopsis novaehollandiae grisea (Vieillot, 1818) Recherche Cape Barren Goose
Cygnus atratus (Latham, 1790) Black Swan
^{*}*Cygnus olor* (Gmelin, 1789) Mute Swan
Stictonetta naevosa (Gould, 1841) Freckled Duck
Tadorna radjah (Lesson, 1828) Radjah Shelduck (Raja Shelduck)
Tadorna radjah rufitergum Hartert, 1905 Burdekin Duck
Tadorna tadornoides (Jardine and Selby, 1828) Australian Shelduck (Mountain Duck)
Malacorhynchus membranaceus (Latham, 1802) Pink-eared Duck
Chenonetta jubata (Latham, 1802) Australian Wood Duck (Wood Duck, Maned Duck)
Nettapus pulchellus Gould, 1842 Green Pygmy Goose
⁴*Anas penelope* Linnaeus, 1758 Eurasian Wigeon
^{*}*Anas platyrhynchos* Linnaeus, 1758 Mallard
⁵*Anas superciliosa* Gmelin, 1789 Pacific Black Duck
Anas rhynchos Latham, 1802 Australasian Shoveler
Anas rhynchos rhynchos Latham, 1802
Anas clypeata Linnaeus, 1758 Northern Shoveler
Anas gracilis Buller, 1869 Grey Teal
Anas castanea (Eyton, 1838) Chestnut Teal
⁶*Anas acuta* Linnaeus, 1758 Northern Pintail
Anas querquedula Linnaeus, 1758 Garganey
Aythya australis (Eyton, 1838) Hardhead
Oxyura australis Gould, 1837 Blue-billed Duck
Biziura lobata (Shaw, 1796) Musk Duck

Order GALLIFORMES**Family MEGAPODIIDAE Megapodes**

- Leipoa ocellata* Gould, 1840 Malleefowl
Megapodius reinwardt Dumont, 1823 Orange-footed Scrubfowl (Orange-legged Scrubfowl)
Megapodius reinwardt tumulus Gould, 1842

Family PHASIANIDAE Pheasants

- Coturnix pectoralis* Gould, 1837 Stubble Quail
Coturnix ypsilophora Bosc, 1792 Brown Quail
⁷*Coturnix ypsilophora cervina* Gould, 1865
Coturnix ypsilophora australis (Latham, 1802)

¹ Formerly in Casuariidae. *D. n. rothschildi* is not accepted as a subspecies.

² Now relocated in monotypic family Anseranatidae, formerly in Anatidae.

³ *D. arcuata* is treated as monotypic with no subspecies.

⁴ Carnarvon 2013.

⁵ *A. superciliosa* is treated as monotypic with no subspecies.

⁶ Now treated as monotypic species following elevation of sub-Antarctic forms as full species.

⁷ The small northern ssp. *C. y. cervina* Gould is separable from other Australian populations.

⁸ <i>Excalfactoria chinensis</i> (Linnaeus, 1766)	King Quail
⁹ <i>Excalfactoria chinensis lineata</i> (Scopoli, 1786)	
^{10*} <i>Phasianus colchicus</i> Linnaeus, 1758	Common Pheasant (Domestic Pheasant)
[*] <i>Pavo cristatus</i> Linnaeus, 1758	Common Peafowl (Indian Peafowl)

Order SPHENISCIFORMES

Family SPHENISCIDAE Penguins

<i>Aptenodytes patagonicus</i> Miller, 1778	King Penguin
¹¹ <i>Aptenodytes patagonicus halli</i> Mathews, 1911	
<i>Eudyptes pachyrhynchus</i> G.R. Gray, 1845	Fiordland Penguin
<i>Eudyptes robustus</i> Oliver, 1953	Snares Penguin
<i>Eudyptes sclateri</i> Buller, 1888	Erect-crested Penguin
<i>Eudyptes chrysocome</i> (Forster, 1781)	Southern Rockhopper Penguin
<i>Eudyptes chrysocome filholi</i> Hutton, 1879	Eastern Rockhopper Penguin
¹² <i>Eudyptes moseleyi</i> Mathews and Iredale, 1921	Northern Rockhopper Penguin
<i>Eudyptes chrysolophus</i> (von Brandt, 1837)	Macaroni Penguin
¹³ <i>Eudyptes chrysolophus schlegeli</i> Finsch, 1876	Royal Penguin
<i>Eudyptula minor</i> (Forster, 1781)	Little Penguin
<i>Eudyptula minor novaehollandiae</i> (Stephens, 1826)	

Order PROCELLARIIFORMES

Family DIOMEDEIDAE Albatrosses

<i>Diomedea exulans</i> Linnaeus, 1758	Wandering Albatross
<i>Diomedea exulans exulans</i> Linnaeus, 1758	Snowy Albatross
¹⁴ <i>Diomedea exulans gibsoni</i> C.J.R. Robertson and Warham, 1992	Gibson's Albatross
¹⁵ <i>Diomedea exulans amsterdamensis</i> Roux, Jouventin, Mougin, Stahl and Weimerskirch, 1983	Amsterdam Albatross
¹⁶ <i>Diomedea exulans dabbenena</i> Mathews, 1929	Tristan Albatross
¹⁷ <i>Diomedea epomophora</i> Lesson, 1825	Southern Royal Albatross
¹⁸ <i>Diomedea sanfordi</i> Murphy, 1917	Northern Royal Albatross
<i>Phoebetria fusca</i> (Hilsenberg, 1822)	Sooty Albatross
<i>Phoebetria palpebrata</i> (J.R. Forster, 1785)	Light-mantled Sooty Albatross
¹⁹ <i>Thalassarche melanophrys</i> (Temminck, 1828)	Black-browed Albatross
<i>Thalassarche melanophrys melanophrys</i> (Temminck, 1828)	
²⁰ <i>Thalassarche melanophrys impavida</i> Mathews, 1912	Campbell Island Albatross
²¹ <i>Thalassarche cauta</i> (Gould, 1841)	Shy Albatross
<i>Thalassarche cauta cauta</i> (Gould, 1841)	
<i>Thalassarche cauta steadi</i> Falla, 1933	White-capped Albatross

⁸ Formerly in *Coturnix*.

⁹ *E. c. lineata* occurs in northern Australia, SE Asia and the Lesser Sundas.

¹⁰ Common Pheasants introduced into Australia are of mixed hybrid stock.

¹¹ Subspecies *halli* recognised on genetic differences. Breeds on Macquarie, Kerguelen, Crozet and Marion islands

¹² Sometimes treated as a ssp. of *E. chrysocome*.

¹³ Often treated as a full species.

¹⁴ Sometimes treated as a full species.

¹⁵ Sometimes treated as a full species. Tracking of birds from breeding islands shows that this subspecies occurs in Western Australian seas, however, no positive records.

¹⁶ Sometimes treated as a full species. Tracking of birds from breeding islands shows that this subspecies occurs in Western Australian seas, however, no positive records.

¹⁷ Sometimes treated as conspecific with *D. sanfordi*.

¹⁸ Sometimes treated as conspecific with *D. epomophora*. Tracking of birds from breeding islands shows that this species occurs in Western Australian seas, however, no positive records.

¹⁹ Formerly in *Diomedea*.

²⁰ Sometimes treated as a full species *T. impavida*.

²¹ Formerly in *Diomedea*.

- ²²*Thalassarche salvini* (Rothschild, 1893) Salvin's Albatross
²³*Thalassarche chrysostoma* (J.R. Forster, 1785) Grey-headed Albatross
²⁴*Thalassarche chlororhynchos* (Gmelin, 1789) Yellow-nosed Albatross
Thalassarche chlororhynchos chlororhynchos (Gmelin, 1789) Atlantic Yellow-nosed Albatross
²⁵*Thalassarche chlororhynchos carteri* (Rothschild, 1903) Indian Ocean Yellow-nosed Albatross

Family PROCELLARIIDAE Petrels and shearwaters

- Macronectes giganteus* (Gmelin, 1789) Southern Giant Petrel
Macronectes halli Mathews, 1912 Northern Giant Petrel
Fulmarus glacialisoides (A. Smith, 1840) Southern Fulmar
Thalassoica antarctica (Gmelin, 1789) Antarctic Petrel
²⁶*Daption capense* (Linnaeus, 1758) Cape Petrel
Daption capense capense (Linnaeus, 1758)
Halobaena caerulea (Gmelin, 1789) Blue Petrel
Pachyptila vittata (G. Forster, 1777) Broad-billed Prion
Pachyptila salvini (Mathews, 1912) Salvin's Prion
Pachyptila salvini salvini (Mathews, 1912)
Pachyptila salvini macgillivrayi (Mathews, 1912) Amsterdam Prion
Pachyptila desolata (Gmelin, 1789) Antarctic Prion
Pachyptila belcheri (Mathews, 1912) Slender-billed Prion
Pachyptila turtur (Kuhl, 1820) Fairy Prion
Pachyptila turtur turtur (Kuhl, 1820)
²⁷*Pterodroma brevirostris* (Lesson, 1831) Kerguelen Petrel
Pterodroma macroptera (Smith, 1840) Great-winged Petrel
Pterodroma macroptera macroptera (Smith, 1840)
²⁸*Pterodroma macroptera gouldi* (Hutton, 1869) Grey-faced Petrel
Pterodroma lessonii (Garnot, 1826) White-headed Petrel
Pterodroma mollis (Gould, 1844) Soft-plumaged Petrel
Pterodroma neglecta (Schlegel, 1863) Kermadec Petrel
Pterodroma heraldica (Salvin, 1888) Herald Petrel
Pterodroma baraui (Jouanin, 1964) Barau's Petrel
Pterodroma cervicalis (Salvin, 1891) White-necked Petrel
Pterodroma leucoptera (Gould, 1844) Gould's Petrel
²⁹*Pterodroma leucoptera leucoptera* (Gould, 1844)
³⁰*Pterodroma rostrata* (Peale, 1848) Tahiti Petrel
Procellaria cinerea Gmelin, 1789 Grey Petrel
Procellaria aequinoctialis Linnaeus, 1758 White-chinned Petrel
Calonectris leucomelas (Temminck, 1835) Streaked Shearwater
Puffinus pacificus (Gmelin, 1789) Wedge-tailed Shearwater
³¹*Puffinus griseus* (Gmelin, 1789) Sooty Shearwater
Puffinus tenuirostris (Temminck, 1835) Short-tailed Shearwater
³²*Puffinus carneipes* Gould, 1844 Fleshy-footed Shearwater
Puffinus gravis (O'Reilly, 1818) Great Shearwater

²² Sometimes treated as a subspecies of *T. cauta*.

²³ Formerly in *Diomedea*.

²⁴ Formerly in *Diomedea*.

²⁵ Sometimes treated as a full species *T. carteri*.

²⁶ Based on photos, individuals of New Zealand breeding population *D. c. australis* possibly also occur in West Aust.

²⁷ Sometimes placed in the genus *Aphrodroma* or *Lugensa*.

²⁸ Sometimes treated as a full species *P. gouldi*.

²⁹ Beach-washed bird found at Eyre 1990.

³⁰ Sometimes placed in the genus *Pseudobulweria*.

³¹ Off Albany 2004.

³² Sometimes placed in the genus *Ardenna*. Western Australian populations sometimes treated as ssp. *P. c. hullianus*.

<i>Puffinus gavia</i> (J.R. Forster, 1844)	Fluttering Shearwater
<i>Puffinus huttoni</i> Mathews, 1912	Hutton's Shearwater
³³ <i>Puffinus persicus</i> Hume, 1872	Persian Shearwater
³⁴ <i>Puffinus heinrothi</i> Reichenow, 1919	Heinroth's Shearwater
<i>Puffinus assimilis</i> Gould, 1838	Little Shearwater

Puffinus assimilis tunneyi Mathews, 1912

<i>Bulweria bulwerii</i> (Jardine and Selby, 1828)	Bulwer's Petrel
<i>Bulweria fallax</i> Jouanin, 1955	Jouanin's Petrel

Family HYDROBATIDAE Storm petrels

<i>Oceanites oceanicus</i> (Kuhl, 1820)	Wilson's Storm Petrel
<i>Oceanites oceanicus oceanicus</i> (Kuhl, 1820)	
³⁵ <i>Carrodia nereis</i> (Gould, 1841)	Grey-backed Storm Petrel
³⁶ <i>Pelagodroma marina</i> (Latham, 1790)	White-faced Storm Petrel
<i>Pelagodroma marina dulciae</i> Mathews, 1912	
³⁷ <i>Fregetta tropica</i> (Gould, 1844)	Black-bellied Storm Petrel
<i>Fregetta tropica tropica</i> (Gould, 1844)	
³⁸ <i>Oceanodroma monorhis</i> (Swinhoe, 1867)	Swinhoe's Storm Petrel
³⁹ <i>Oceanodroma leucorhoa</i> (Vieillot, 1818)	Leach's Storm Petrel
<i>Oceanodroma leucorhoa leucorhoa</i> (Vieillot, 1818)	
⁴⁰ <i>Oceanodroma matsudariae</i> Kuroda, 1922	Matsudaira's Storm Petrel

Family PELECANOIDIDAE Diving petrels

<i>Pelecanoides urinatrix</i> (Gmelin, 1789)	Common Diving Petrel
<i>Pelecanoides urinatrix exsul</i> Salvin, 1896	

Order PODICIPEDIFORMES

Family PODICIPEDIDAE Grebes

<i>Tachybaptus ruficollis</i> (Pallas, 1764)	Little Grebe (Red-throated Little Grebe)
⁴¹ <i>Tachybaptus ruficollis tricolor</i> (G.R. Gray, 1861)	
<i>Tachybaptus novaehollandiae</i> (Stephens, 1826)	Australasian Grebe (Black-throated Grebe)
<i>Tachybaptus novaehollandiae novaehollandiae</i> (Stephens, 1826)	
<i>Poliocephalus poliocephalus</i> (Jardine and Selby, 1827)	Hoary-headed Grebe
<i>Podiceps cristatus</i> (Linnaeus, 1758)	Great Crested Grebe
<i>Podiceps cristatus australis</i> Gould, 1844	

Order PHAETHONTIFORMES

Family PHAETHONTIDAE Tropicbirds

<i>Phaethon aethereus</i> Linnaeus, 1758	Red-billed Tropicbird
⁴² <i>Phaethon aethereus indicus</i> Hume, 1876	
⁴³ <i>Phaethon rubricauda</i> Boddaert, 1783	Red-tailed Tropicbird
⁴⁴ <i>Phaethon lepturus</i> Daudin, 1802	White-tailed Tropicbird

³³ Refer A. Pittie (*Indian Birds* Vol. 2. 2006) for confirmation of publication date. Region of origin of ssp. visiting Western Australia unresolved.

³⁴ Timor Sea 2014. Photographs only.

³⁵ Sometimes placed in the genus *Oceanites*.

³⁶ Sometimes placed in the genus *Oceanites*.

³⁷ Sometimes placed in the genus *Oceanites*.

³⁸ Sometimes treated as conspecific with *O. leucorhoa*.

³⁹ Sometimes placed in the genus *Hydrobates*.

⁴⁰ Mostly listed as *O. matsudariae*, but Kuroda latinised Y. Matsudaira's Japanese name as *matsudariae* and this has not been validly amended.

⁴¹ Sometimes treated as a full species *T. tricolor*.

⁴² Ashmore Reef 2014.

⁴³ Treated as monotypic with no subspecies recognised.

⁴⁴ Christmas Is. *P. l. fulvus* is not accepted as a ssp. due to the complete gradation between the two colour morphs.

Order CICONIIFORMES**Family CICONIIDAE Storks**

- Ephippiorhynchus asiaticus* (Latham, 1790) Black-necked Stork
Ephippiorhynchus asiaticus australis (Shaw, 1800)

Order PELECANIFORMES**Family THRESKIORNITHIDAE Ibises and spoonbills**

- ⁴⁵*Threskiornis moluccus* (Cuvier, 1829) Australian White Ibis
Threskiornis spinicollis (Jameson, 1835) Straw-necked Ibis
Plegadis falcinellus (Linnaeus, 1766) Glossy Ibis
⁴⁶*Platalea regia* Gould, 1838 Royal Spoonbill
Platalea flavipes Gould, 1838 Yellow-billed Spoonbill

Family ARDEIDAE Herons and bitterns

- Botaurus poiciloptilus* (Wagler, 1827) Australasian Bittern
Ixobrychus minutus (Linnaeus, 1766) Little Bittern
⁴⁷*Ixobrychus minutus dubius* Mathews, 1912
⁴⁸*Ixobrychus sinensis* (Gmelin, 1789) Yellow Bittern (Chinese Little Bittern)
⁴⁹*Ixobrychus flavicollis* (Latham, 1790) Black Bittern
Ixobrychus flavicollis australis (Lesson, 1831)
Nycticorax nycticorax (Linneaus, 1758) Black-crowned Night Heron
⁵⁰*Nycticorax nycticorax nycticorax* (Linneaus, 1758)
Nycticorax caledonicus (Gmelin, 1789) Nankeen Night Heron (Rufous Night Heron)
⁵¹*Nycticorax caledonicus australasiae* (Vieillot, 1823)
Butorides striata (Linnaeus, 1758) Striated Heron (Mangrove Heron)
⁵²*Butorides striata stagnatilis* (Gould, 1848)
⁵³*Ardeola speciosa* (Horsfield, 1821) Javan Pond Heron
⁵⁴*Ardea ibis* Linnaeus, 1758 Cattle Egret
⁵⁵*Ardea ibis coromanda* (Boddaert, 1783)
⁵⁶*Ardea cinerea* Linneaus, 1758 Grey Heron
Ardea pacifica Latham, 1802 White-necked Heron
Ardea sumatrana Raffles, 1822 Great-billed Heron
⁵⁷*Ardea purpurea* Linnaeus, 1766 Purple Heron
⁵⁸*Ardea modesta* J.E. Gray, 1831 Eastern Great Egret
⁵⁹*Ardea intermedia* Wagler, 1829 Intermediate Egret
Ardea intermedia intermedia Wagler, 1829
⁶⁰*Ardea intermedia plumifera* Gould, 1848

⁴⁵ Sometimes treated as a ssp. of the Sacred Ibis *T. aethiopicus*.

⁴⁶ Forming a superspecies with Eurasian Spoonbill *P. leucorodia* of Europe, Asia and Africa.

⁴⁷ Sometimes treated as a full species.

⁴⁸ Kalgoorlie 1967.

⁴⁹ Sometimes placed in the genus *Dupetor*.

⁵⁰ Cocos Islands 1941, Ashmore Reef 1994.

⁵¹ Formerly *N. c. hilli*.

⁵² Only one ssp. *B. s. stagnatilis* recognised in Western Australia (Johnstone 1990).

⁵³ No specimens extant; photographs only which indicate most probably the nominate ssp.

⁵⁴ Sometimes placed in the genus *Bubulcus*.

⁵⁵ Sometimes treated as a full species Eastern Cattle Egret.

⁵⁶ Busselton 2002. Most likely ssp. *jouyi* of eastern Asia.

⁵⁷ Herdsman Lake 2013. Most likely ssp. *manilensis* of Asia.

⁵⁸ Sometimes treated as a ssp. of *A. alba*.

⁵⁹ Sometimes placed in the genus *Egretta*.

⁶⁰ Sometimes treated as a full species.

⁶¹ <i>Ardea picata</i> Gould, 1845	Pied Heron
⁶² <i>Ardea novaehollandiae</i> Latham, 1790	White-faced Heron
⁶³ <i>Ardea garzetta</i> Linnaeus, 1766	Little Egret
⁶⁴ <i>Ardea garzetta immaculata</i> (Gould, 1846)	
⁶⁵ <i>Ardea sacra</i> Gmelin, 1789	Eastern Reef Heron (Eastern Reef Egret)
<i>Ardea sacra sacra</i> Gmelin, 1789	

Family PELECANIDAE Pelicans

<i>Pelecanus conspicillatus</i> Temminck, 1824	Australian Pelican
--	--------------------

Order SULIFORMES**Family FREGATIDAE Frigatebirds**

<i>Fregata andrewsi</i> Mathews, 1914	Christmas Island Frigatebird
<i>Fregata minor</i> (Gmelin, 1789)	Greater Frigatebird
<i>Fregata ariel</i> (G.R. Gray, 1845)	Lesser Frigatebird
<i>Fregata ariel ariel</i> (G.R. Gray, 1845)	

Family SULIDAE Gannets and boobies

⁶⁶ <i>Morus capensis</i> (Lichtenstein, 1823)	Cape Gannet
<i>Morus serrator</i> (G.R. Gray, 1843)	Australasian Gannet
<i>Papasula abbotti</i> (Ridgway, 1893)	Abbott's Booby
<i>Sula dactylatra</i> Lesson, 1831	Masked Booby
<i>Sula dactylatra personata</i> Gould, 1846	
<i>Sula sula</i> (Linnaeus, 1766)	Red-footed Booby
<i>Sula leucogaster</i> (Boddaert, 1783)	Brown Booby
<i>Sula leucogaster plotus</i> (J.R. Forster, 1844)	

Family PHALACROCORACIDAE Cormorants

⁶⁷ <i>Phalacrocorax melanoleucus</i> (Vieillot, 1817)	Little Pied Cormorant
<i>Phalacrocorax melanoleucus melanoleucus</i> (Vieillot, 1817)	
<i>Phalacrocorax fuscescens</i> (Vieillot, 1817)	Black-faced Cormorant
<i>Phalacrocorax sulcirostris</i> (von Brandt, 1837)	Little Black Cormorant
<i>Phalacrocorax varius</i> (Gmelin, 1789)	Pied Cormorant (Australian Pied Cormorant)
<i>Phalacrocorax varius hypoleucus</i> (von Brandt, 1837)	
<i>Phalacrocorax carbo</i> (Linnaeus, 1758)	Great Cormorant
<i>Phalacrocorax carbo novaehollandiae</i> Stephens, 1826	
⁶⁸ <i>Phalacrocorax verrucosus</i> (Cabanis, 1875)	Kerguelen Cormorant (Kerguelen Shag)

Family ANHINGIDAE Darters

⁶⁹ <i>Anhinga novaehollandiae</i> (Gould, 1847)	Australasian Darter
<i>Anhinga novaehollandiae novaehollandiae</i> (Gould, 1847)	

Order ACCIPITRIFORMES**Family PANDIONIDAE Osprey**

<i>Pandion haliaetus</i> (Linnaeus, 1758)	Osprey
⁷⁰ <i>Pandion haliaetus cristatus</i> (Vieillot, 1816)	(Eastern Osprey)

⁶¹ Sometimes placed in the genus *Egretta*.

⁶² Sometimes placed in the genus *Egretta*.

⁶³ Sometimes placed in the genus *Egretta*.

⁶⁴ Sometimes synonymised with *A. g. nigripes*.

⁶⁵ Sometimes placed in the genus *Egretta*.

⁶⁶ *Morus capensis* and *Morus serrator* sometimes treated as conspecific with Northern Gannet (*Morus bassana*) or alternatively all three treated as a superspecies and sometimes partitioned off in the genus *Sula*.

⁶⁷ Sometimes placed in the genus *Microcarbo*.

⁶⁸ Rottnest Island 1986. Sometimes placed in the genus *Leucocarbo*.

⁶⁹ Previously treated as a ssp. of *A. melanogaster*.

⁷⁰ Sometimes treated as a full species *P. cristatus* (Eastern Osprey).

Family ACCIPITRIDAE Hawks, eagles, kites and harriers

<i>Elanus caeruleus</i> (Desfontaines, 1789)	Black-shouldered Kite
⁷¹ <i>Elanus caeruleus axillaris</i> (Latham, 1802)	Australian Black-shouldered Kite
<i>Elanus scriptus</i> Gould, 1842	Letter-winged Kite
<i>Pernis ptilorhyncus</i> (Temminck, 1821)	Oriental Honey-Buzzard (Eastern or Crested Honey-Buzzard)
⁷² <i>Pernis ptilorhyncus orientalis</i> Taczanowski, 1891	
⁷³ <i>Hamirostra isura</i> (Gould, 1838)	Square-tailed Kite
<i>Hamirostra melanosternon</i> (Gould, 1841)	Black-breasted Buzzard
<i>Aviceda subcristata</i> (Gould, 1838)	Pacific Baza
<i>Aviceda subcristata subcristata</i> (Gould, 1838)	
⁷⁴ <i>Hieraetus morphnoides</i> (Gould, 1841).....	Little Eagle
<i>Aquila audax</i> (Latham, 1802)	Wedge-tailed Eagle
<i>Erythrotriorchis radiatus</i> (Latham, 1802)	Red Goshawk
⁷⁵ <i>Accipiter soloensis</i> (Horsfield, 1821)	Chinese Sparrowhawk (Chinese Goshawk)
⁷⁶ <i>Accipiter novaehollandiae</i> (Gmelin, 1788)	Grey Goshawk (Variable Goshawk)
<i>Accipiter novaehollandiae novaehollandiae</i> (Gmelin, 1788)	
<i>Accipiter fasciatus</i> (Vigors and Horsfield, 1827)	Brown Goshawk
<i>Accipiter fasciatus fasciatus</i> (Vigors and Horsfield, 1827)	
<i>Accipiter fasciatus didimus</i> (Mathews, 1912)	
⁷⁷ <i>Accipiter gularis</i> (Temminck and Schlegel, 1844)	Japanese Sparrowhawk
<i>Accipiter cirrocephalus</i> (Vieillot, 1817)	Collared Sparrowhawk
<i>Accipiter cirrocephalus cirrocephalus</i> (Vieillot, 1817)	
<i>Circus approximans</i> Peale, 1848	Swamp Harrier
<i>Circus assimilis</i> Jardine and Selby, 1828	Spotted Harrier
<i>Milvus migrans</i> (Boddaert, 1783)	Black Kite
<i>Milvus migrans affinis</i> Gould, 1838	
<i>Haliastur sphenurus</i> (Vieillot, 1818)	Whistling Kite
<i>Haliastur indus</i> (Boddaert, 1783)	Brahminy Kite
<i>Haliastur indus girrenera</i> (Vieillot, 1822)	
<i>Haliaeetus leucogaster</i> (Gmelin, 1788)	White-bellied Sea-Eagle

Order OTIDIFORMES**Family OTIDIDAE Bustards**

⁷⁸ <i>Ardeotis australis</i> (J.E. Gray, 1829)	Australian Bustard
---	--------------------

Order GRUIFORMES**Family RALLIDAE Rails, crakes, waterhens and coots**

⁷⁹ <i>Rallina fasciata</i> (Raffles, 1822)	Red-legged Crake
<i>Gallirallus philippensis</i> (Linnaeus, 1766)	Buff-banded Rail
<i>Gallirallus philippensis mellori</i> (Mathews, 1912)	
<i>Lewinia pectoralis</i> (Temminck, 1831)	Lewin's Rail
⁸⁰ <i>Lewinia pectoralis clelandi</i> (Mathews, 1911)	
<i>Crex crex</i> (Linnaeus, 1758)	Corncrake (Corn Crake)

⁷¹ Sometimes treated as a full species *E. axillaris* Black-winged Kite (Australian Kite).

⁷² Leinster, Joondalup.

⁷³ Usually placed in monotypic genus *Lophoictinia*, but along with its single congener *H. melanosternon* comprise an old Australian group without close relatives.

⁷⁴ Sometimes placed in the genus *Aquila*.

⁷⁵ Ashmore Reef 2013.

⁷⁶ *A. novaehollandiae* is sometimes treated as a full species; the status of other ssp. of the Variable Goshawk e.g. *A. n. hiogaster* and *A. n. griseogularis* is uncertain.

⁷⁷ Ashmore Reef 2011.

⁷⁸ Previously merged with the genus *Otis*.

⁷⁹ Broome 1958 and Whim Creek 2007.

⁸⁰ Possibly extinct, last report from south-western WA in 1932.

<i>Amaurornis moluccana</i> (Wallace, 1865)	Bush-hen
<i>Amaurornis moluccana ruficrissa</i> (Gould, 1869)	Pale-vented Bush-hen
<i>Amaurornis phoenicurus</i> (Pennant, 1769)	White-breasted Waterhen
⁸¹ <i>Amaurornis phoenicurus leucomelanus</i> (S. Müller, 1842)	
<i>Porzana pusilla</i> (Pallas, 1776)	Baillon's Crake
<i>Porzana pusilla palustris</i> Gould, 1843	
<i>Porzana fluminea</i> Gould, 1843	Australian Spotted Crake
<i>Porzana tabuensis</i> (Gmelin, 1789)	Spotless Crake
<i>Porzana cinerea</i> (Vieillot, 1819)	White-browed Crake
<i>Eulabeornis castaneoventris</i> Gould, 1844	Chestnut Rail
<i>Eulabeornis castaneoventris castaneoventris</i> Gould, 1844	
⁸² <i>Gallicrex cinerea</i> (Gmelin, 1789)	Watercock
<i>Porphyrio porphyrio</i> (Linnaeus, 1758)	Purple Swamphen
<i>Porphyrio porphyrio melanotus</i> Temminck, 1820	
<i>Porphyrio porphyrio bellus</i> Gould, 1841	
<i>Gallinula tenebrosa</i> Gould, 1846	Dusky Moorhen
<i>Gallinula tenebrosa tenebrosa</i> Gould, 1846	
⁸³ <i>Tribonyx ventralis</i> (Gould, 1837)	Black-tailed Native-hen
<i>Fulica atra</i> Linnaeus, 1758	Eurasian Coot
<i>Fulica atra australis</i> Gould, 1845	

Family GRUIDAE Cranes

<i>Grus antigone</i> (Linnaeus, 1758)	Sarus Crane
<i>Grus antigone gillae</i> Schodde, Blackman and Haffenden, 1988	
<i>Grus rubicunda</i> (Perry, 1810)	Brolga

Order CHARADRIIFORMES**Family TURNICIDAE Button-quails**

<i>Turnix maculosus</i> (Temminck, 1815)	Red-backed Button-quail
<i>Turnix maculosus melanotus</i> (Gould, 1837)	
<i>Turnix castanotus</i> (Gould, 1840)	Chestnut-backed Button-quail
<i>Turnix varius</i> (Latham, 1802)	Painted Button-quail
<i>Turnix varius varius</i> (Latham, 1802)	
<i>Turnix varius scintillans</i> (Gould, 1845)	Abrolhos Painted Button-quail
<i>Turnix pyrrhothorax</i> (Gould, 1841)	Red-chested Button-quail
<i>Turnix velox</i> (Gould, 1841)	Little Button-quail

Family BURHINIDAE Stone-curlews

<i>Burhinus grallarius</i> (Latham, 1802)	Bush Stone-curlew (Bush Thick-knee)
<i>Esacus magnirostris</i> (Vieillot, 1818)	Beach Stone-curlew (Beach Thick-knee)

Family HAEMATOPODIDAE Oystercatchers

<i>Haematopus longirostris</i> Vieillot, 1817	Pied Oystercatcher
<i>Haematopus fuliginosus</i> Gould, 1845	Sooty Oystercatcher
<i>Haematopus fuliginosus fuliginosus</i> Gould, 1845	
<i>Haematopus fuliginosus ophthalmicus</i> Castelnau and Ramsay, 1877	

Family RECURVIROSTRIDAE Stilts and avocets

<i>Himantopus himantopus</i> (Linnaeus, 1758)	Black-winged Stilt
⁸⁴ <i>Himantopus himantopus leucocephalus</i> Gould, 1837	
<i>Cladorhynchus leucocephalus</i> (Vieillot, 1816)	Banded Stilt
<i>Recurvirostra novaehollandiae</i> Vieillot, 1816	Red-necked Avocet

⁸¹ Ashmore Reef 2003.⁸² Ashmore Reef 2012.⁸³ Sometimes placed in the genus *Gallinula*.⁸⁴ Sometimes treated as a full species *H. leucocephalus*.

Family CHARADRIIDAE Plovers, dotterels and lapwings

<i>Vanellus tricolor</i> (Vieillot, 1818)	Banded Lapwing
<i>Vanellus miles</i> (Boddaert, 1783)	Masked Lapwing
<i>Vanellus miles miles</i> (Boddaert, 1783)	
<i>Vanellus miles novaehollandiae</i> Stephens, 1819	
<i>Erythrogonys cinctus</i> Gould, 1838	Red-kneed Dotterel
<i>Peltohyas australis</i> (Gould, 1841)	Inland Dotterel
85 <i>Pluvialis fulva</i> (Gmelin, 1789)	Pacific Golden Plover
86 <i>Pluvialis dominica</i> (Müller, 1776)	American Golden Plover
<i>Pluvialis squatarola</i> (Linnaeus, 1758)	Grey Plover
87 <i>Pluvialis squatarola squatarola</i> (Linnaeus, 1758)	
88 <i>Charadrius hiaticula</i> Linnaeus, 1758	Common Ringed Plover
89 <i>Charadrius semipalmatus</i> Bonaparte, 1825	Semipalmated Plover
<i>Charadrius dubius</i> Scopoli, 1786	Little Ringed Plover
90 <i>Charadrius dubius curonicus</i> Gmelin, 1789	
91 <i>Charadrius alexandrinus</i> Linnaeus, 1758	Kentish Plover
<i>Charadrius alexandrinus [dealbatus]</i> (Swinhoe, 1870)	
<i>Charadrius ruficollis</i> Temminck, 1822	Red-capped Plover
92 <i>Charadrius bicinctus</i> Jardine and Selby, 1827	Double-banded Plover
93 <i>Charadrius bicinctus bicinctus</i> Jardine and Selby, 1827	
94 <i>Charadrius mongolus</i> Pallas, 1776	Lesser Sand Plover
<i>Charadrius mongolus mongolus</i> Pallas, 1776	
95 <i>Charadrius mongolus atrifrons</i> Wagler, 1829	
<i>Charadrius leschenaultii</i> Lesson, 1826	Greater Sand Plover
<i>Charadrius leschenaultii leschenaultii</i> Lesson, 1826	
96 <i>Charadrius veredus</i> Gould, 1848	Oriental Plover
97 <i>Thinornis cucullatus</i> (Vieillot, 1818)	Hooded Plover (Hooded Dotterel)
<i>Elseyornis melanops</i> (Vieillot, 1818)	Black-fronted Dotterel

Family ROSTRATULIDAE Painted snipes

<i>Rostratula benghalensis</i> (Linnaeus, 1758)	Painted Snipe (Greater Painted Snipe)
98 <i>Rostratula benghalensis australis</i> (Gould, 1838)	Australian Painted Snipe
Family JACANIDAE Jacanas	
99 <i>Irediparra gallinacea</i> (Temminck, 1828)	Comb-crested Jacana
100 <i>Hydrophasianus chirurgus</i> (Scopoli, 1786)	Pheasant-tailed Jacana

85 Non-breeding migrant.

86 Broome 2009-2010.

87 Non-breeding migrant.

88 Non-breeding migrant. No specimens available, but ssp. probably *tundrae*.

89 Martin's Tank 2007.

90 Leslie Saltworks 1993.

91 Non-breeding migrant.

92 Non-breeding migrant.

93 Non-breeding migrant.

94 Non-breeding migrant.

95 Non-breeding migrant.

96 Non-breeding migrant.

97 Sometimes placed in the genus *Charadrius*.98 Sometimes treated as a full species *R. australis*.99 Sometimes placed in monotypic genus *Metopidius*. The retention of the eight species of Jacana into six genera cannot be justified. A better reflection of generic relationships may well be to retain *Hydrophasianus* as monotypic and combine all others in *Jacana*.

100 Paraburdoo 1974.

Family SCOLOPACIDAE Sandpipers and snipes

¹⁰¹ <i>Gallinago hardwickii</i> (J.E. Gray, 1831)	Latham's Snipe
¹⁰² <i>Gallinago stenura</i> (Bonaparte, 1830)	Pin-tailed Snipe
<i>Gallinago megala</i> Swinhoe, 1861	Swinhoe's Snipe
¹⁰³ <i>Limnodromus semipalmatus</i> (Blyth, 1848)	Asian Dowitcher
¹⁰⁴ <i>Limosa limosa</i> (Linnaeus, 1758)	Black-tailed Godwit
<i>Limosa limosa melanuroides</i> Gould, 1846	
¹⁰⁵ <i>Limosa haemastica</i> (Linnaeus, 1758)	Hudsonian Godwit
<i>Limosa lapponica</i> (Linnaeus, 1758)	Bar-tailed Godwit
<i>Limosa lapponica [lapponica]</i> (Linnaeus, 1758)	
¹⁰⁶ <i>Limosa lapponica menzbieri</i> Portenko, 1936	
¹⁰⁷ <i>Limosa lapponica [bauerii]</i> Naumann, 1836	
<i>Numenius minutus</i> Gould, 1841	Little Curlew
¹⁰⁸ <i>Numenius phaeopus</i> (Linnaeus, 1758)	Whimbrel
<i>Numenius phaeopus variegatus</i> (Scopoli, 1786)	
<i>Numenius arquata</i> (Linnaeus, 1758)	Eurasian Curlew
<i>Numenius arquata orientalis</i> C.L. Brehm, 1831	
¹⁰⁹ <i>Numenius madagascariensis</i> (Linnaeus, 1766)	Far Eastern Curlew (Eastern Curlew)
¹¹⁰ <i>Tringa erythropus</i> (Pallas, 1764)	Spotted Redshank
¹¹¹ <i>Tringa totanus</i> (Linnaeus, 1758)	Common Redshank
¹¹² <i>Tringa stagnatilis</i> (Bechstein, 1803)	Marsh Sandpiper
¹¹³ <i>Tringa nebularia</i> (Gunnerus, 1767)	Common Greenshank
¹¹⁴ <i>Tringa guttifer</i> (Nordmann, 1835)	Nordmann's Greenshank
¹¹⁵ <i>Tringa glareola</i> Linnaeus, 1758	Wood Sandpiper
¹¹⁶ <i>Tringa brevipes</i> (Vieillot, 1816)	Grey-tailed Tattler
¹¹⁷ <i>Tringa cinerea</i> (Güldenstädt, 1775)	Terek Sandpiper
¹¹⁸ <i>Tringa hypoleucos</i> Linnaeus, 1758	Common Sandpiper
<i>Arenaria interpres</i> (Linnaeus, 1758)	Ruddy Turnstone
¹¹⁹ <i>Arenaria interpres interpres</i> (Linnaeus, 1758)	
¹²⁰ <i>Calidris tenuirostris</i> (Horsfield, 1821)	Great Knot

¹⁰¹ Non-breeding migrant.

¹⁰² Non-breeding migrant.

¹⁰³ Non-breeding migrant.

¹⁰⁴ Non-breeding migrant.

¹⁰⁵ Joondalup 2012.

¹⁰⁶ Non-breeding migrant.

¹⁰⁷ Non-breeding migrant.

¹⁰⁸ Non-breeding migrant.

¹⁰⁹ Non-breeding migrant.

¹¹⁰ Lake Eda 2006.

¹¹¹ Subspecific identity of birds visiting Western Australia unknown.

¹¹² Non-breeding migrant.

¹¹³ Non-breeding migrant.

¹¹⁴ Eighty Mile Beach 2006 and 2009.

¹¹⁵ Non-breeding migrant.

¹¹⁶ Non-breeding migrant.

¹¹⁷ Sometimes placed in the genus *Xenus*. *Tringa terek* is a synonym. Non-breeding migrant.

¹¹⁸ Sometimes placed in the genus *Actitis*. Non-breeding migrant.

¹¹⁹ Non-breeding migrant.

¹²⁰ Non-breeding migrant.

¹²¹ <i>Calidris canutus</i> (Linnaeus, 1758)	Red Knot
¹²² <i>Calidris canutus piersmai</i> Tomkovich, 2001	
<i>Calidris canutus rogersi</i> (Mathews, 1913)	
¹²³ <i>Calidris alba</i> (Pallas, 1764)	Sanderling
¹²⁴ <i>Calidris ruficollis</i> (Pallas, 1776)	Red-necked Stint
¹²⁵ <i>Calidris minuta</i> (Leisler, 1812)	Little Stint
<i>Calidris temminckii</i> (Leisler, 1812)	Temminck's Stint
¹²⁶ <i>Calidris subminuta</i> (Middendorff, 1853)	Long-toed Stint
¹²⁷ <i>Calidris fuscicollis</i> (Vieillot, 1819)	White-rumped Sandpiper
¹²⁸ <i>Calidris bairdii</i> (Coues, 1861)	Baird's Sandpiper
¹²⁹ <i>Calidris melanotos</i> (Vieillot, 1819)	Pectoral Sandpiper
¹³⁰ <i>Calidris acuminata</i> (Horsfield, 1821)	Sharp-tailed Sandpiper
¹³¹ <i>Calidris ferruginea</i> (Pontoppidan, 1763)	Curlew Sandpiper
¹³² <i>Calidris alpina</i> (Linnaeus, 1758)	Dunlin
<i>Calidris alpina sakhalina</i> (Vieillot, 1816)	
¹³³ <i>Limicola falcinellus</i> (Pontoppidan, 1763)	Broad-billed Sandpiper
<i>Limicola falcinellus falcinellus</i> (Pontoppidan, 1763)	
<i>Limicola falcinellus sibirica</i> Dresser, 1876	
¹³⁴ <i>Tryngites subruficollis</i> (Vieillot, 1819)	Buff-breasted Sandpiper
<i>Philomachus pugnax</i> (Linnaeus, 1758)	Ruff
¹³⁵ <i>Phalaropus lobatus</i> (Linnaeus, 1758)	Red-necked Phalarope
¹³⁶ <i>Phalaropus fulicarius</i> (Linnaeus, 1758)	Red Phalarope
Family GLAREOLIDAE Pratincoles	
<i>Stiltia isabella</i> (Vieillot, 1816)	Australian Pratincole
¹³⁷ <i>Glareola maldivarum</i> J.R. Forster, 1795	Oriental Pratincole
Family LARIDAE Gulls and terns	
<i>Anous stolidus</i> (Linnaeus, 1758)	Common Noddy (Brown Noddy)
<i>Anous stolidus pileatus</i> (Scopoli, 1786)	
<i>Anous tenuirostris</i> (Temminck, 1823)	Lesser Noddy
¹³⁸ <i>Anous tenuirostris melanops</i> Gould, 1846	Australian Lesser Noddy
<i>Anous minutus</i> Boie, 1844	Black Noddy
<i>Anous minutus minutus</i> Boie, 1844	
¹³⁹ <i>Gygis alba</i> (Sparrman, 1786)	White Tern (White Noddy)

¹²¹ Non-breeding migrant.

¹²² Non-breeding migrant.

¹²³ Non-breeding migrant.

¹²⁴ Non-breeding migrant.

¹²⁵ Broome 2009.

¹²⁶ Non-breeding migrant.

¹²⁷ Forrestdale Lake 1981.

¹²⁸ Eyre 1979.

¹²⁹ Non-breeding migrant.

¹³⁰ Non-breeding migrant.

¹³¹ Non-breeding migrant.

¹³² Port Hedland Saltworks 1977.

¹³³ Non-breeding migrant.

¹³⁴ Eyre 1982.

¹³⁵ Non-breeding migrant.

¹³⁶ Greenough River 2012.

¹³⁷ Non-breeding migrant.

¹³⁸ Newly established population on Ashmore Reef not ascribed to subspecies.

¹³⁹ Probably referable to *G. a. candida*.

¹⁴⁰ <i>Larus sabini</i> Sabine, 1819	Sabine's Gull
¹⁴¹ <i>Larus novaehollandiae</i> Stephens, 1826	Silver Gull
<i>Larus novaehollandiae novaehollandiae</i> Stephens, 1826	
<i>Larus ridibundus</i> Linnaeus, 1766	Black-headed Gull
¹⁴² <i>Larus ridibundus sibiricus</i> Buturlin, 1911	
¹⁴³ <i>Larus atricilla</i> Linnaeus, 1758	Laughing Gull
¹⁴⁴ <i>Larus atricilla [megalopterus]</i> Bruch, 1855	
¹⁴⁵ <i>Larus pipixcan</i> Wagler, 1831	Franklin's Gull
<i>Larus pacificus</i> Latham, 1802	Pacific Gull
<i>Larus pacificus georgii</i> King, 1826	
<i>Larus crassirostris</i> Vieillot, 1818	Black-tailed Gull
<i>Larus dominicanus</i> Lichtenstein, 1823	Kelp Gull
<i>Larus dominicanus dominicanus</i> Lichtenstein, 1823	
<i>Larus fuscus</i> Linnaeus, 1758	Lesser Black-backed Gull (Baltic Gull)
<i>Larus fuscus fuscus</i> Linnaeus, 1758	
¹⁴⁶ <i>Sterna nilotica</i> Gmelin, 1789	Gull-billed Tern
<i>Sterna nilotica [nilotica]</i> Gmelin, 1790	
<i>Sterna nilotica affinis</i> Horsfield, 1821	Asian Gull-billed Tern
<i>Sterna nilotica macrotarsa</i> Gould, 1837	Australian Gull-billed Tern
¹⁴⁷ <i>Sterna caspia</i> Pallas, 1770	Caspian Tern
¹⁴⁸ <i>Sterna bergii</i> Lichtenstein, 1823	Crested Tern (Greater Crested Tern)
¹⁴⁹ <i>Sterna bergii cristata</i> Stephens, 1826	
¹⁵⁰ <i>Sterna bengalensis</i> Lesson, 1831	Lesser Crested Tern
¹⁵¹ <i>Sterna albifrons</i> (Pallas, 1764)	White-shafted Little Tern (Little Tern)
<i>Sterna albifrons sinensis</i> Gmelin, 1789	
¹⁵² <i>Sterna nereis</i> (Gould, 1843)	Fairy Tern
¹⁵³ <i>Sterna nereis nereis</i> (Gould, 1843)	
¹⁵⁴ <i>Sterna anaethetus</i> Scopoli, 1786	Bridled Tern
<i>Sterna anaethetus anaethetus</i> Scopoli, 1786	
¹⁵⁵ <i>Sterna fuscata</i> Linnaeus, 1766	Sooty Tern
<i>Sterna fuscata nubilosa</i> Sparrman, 1788	
<i>Sterna dougallii</i> Montagu, 1813	Roseate Tern
<i>Sterna dougallii bangsi</i> Mathews, 1912	
<i>Sterna dougallii gracilis</i> Gould, 1845	
<i>Sterna striata</i> Gmelin, 1789	White-fronted Tern
¹⁵⁶ <i>Sterna sumatrana</i> Raffles, 1822	Black-naped Tern

¹⁴⁰ Eighty Mile Beach 2004. Sometimes placed in the genus *Xema*.

¹⁴¹ Sometimes placed in the genus *Chroicocephalus*. All Australian forms treated as single ssp. (Johnstone 1982, *Rec. West. Aust. Mus.* **10**: 133–165).

¹⁴² Broome 1999. Sometimes placed in the genus *Chroicocephalus*.

¹⁴³ Eyre 1998-99. Sometimes placed in the genus *Leucophaeus*.

¹⁴⁴ Eyre 1998-99.

¹⁴⁵ Sometimes placed in the genus *Leucophaeus*.

¹⁴⁶ Sometimes placed in the genus *Gelochelidon*.

¹⁴⁷ Sometimes placed in the genus *Hydroprogne*.

¹⁴⁸ Sometimes placed in the genus *Thalasseus*.

¹⁴⁹ Only one ssp. *S. b. cristata* recognised for mainland Australia.

¹⁵⁰ Sometimes placed in the genus *Thalasseus*.

¹⁵¹ Sometimes placed in the genus *Sternula*.

¹⁵² Sometimes placed in the genus *Sternula*.

¹⁵³ Only nominate *nereis* recognised for mainland Australia.

¹⁵⁴ Sometimes placed in the genus *Onychoprion*.

¹⁵⁵ Sometimes placed in the genus *Onychoprion*.

¹⁵⁶ Non-breeding migrant.

¹⁵⁷ <i>Sterna hirundo</i> Linnaeus, 1758	Common Tern
¹⁵⁸ <i>Sterna hirundo hirundo</i> Linnaeus, 1758	
¹⁵⁹ <i>Sterna hirundo [minussensis]</i> Sushkin, 1925	
¹⁶⁰ <i>Sterna hirundo longipennis</i> Nordmann, 1835	
¹⁶¹ <i>Sterna paradisaea</i> Pontoppidan, 1763	Arctic Tern
<i>Sterna vittata</i> Gmelin, 1789	Antarctic Tern
¹⁶² <i>Sterna hybrida</i> Pallas, 1811	Whiskered Tern
<i>Sterna hybrida javanica</i> Horsfield, 1821	
¹⁶³ <i>Sterna leucoptera</i> Temminck, 1815	White-winged Black Tern
Family STERCORARIIDAE Skuas	
¹⁶⁴ <i>Stercorarius maccormicki</i> Saunders, 1893	South Polar Skua
<i>Stercorarius antarcticus</i> (Lesson, 1831)	Brown Skua
<i>Stercorarius antarcticus lonnbergi</i> (Mathews, 1912)	
<i>Stercorarius pomarinus</i> (Temminck, 1815)	Pomarine Skua (Pomarine Jaeger)
<i>Stercorarius parasiticus</i> (Linnaeus, 1758)	Arctic Skua (Parasitic Jaeger)
<i>Stercorarius longicaudus</i> Vieillot, 1819	Long-tailed Skua (Long-tailed Jaeger)

Order COLUMBIFORMES**Family COLUMBIDAE Pigeons and doves**

^{165*} <i>Columba livia</i> Gmelin, 1789	Domestic Pigeon (Rock Dove)
^{166*} <i>Spilopelia chinensis</i> (Scopoli, 1786)	Spotted Turtle-Dove
^{167*} <i>Spilopelia chinensis tigrina</i> (Temminck, 1810)	
^{168*} <i>Spilopelia senegalensis</i> (Linnaeus, 1766)	Laughing Turtle-Dove
* <i>Spilopelia senegalensis senegalensis</i> (Linnaeus, 1766)	
<i>Chalcophaps indica</i> (Linnaeus, 1758)	Emerald Dove
¹⁶⁹ <i>Chalcophaps indica longirostris</i> Gould, 1848	
<i>Phaps chalcoptera</i> (Latham, 1790)	Common Bronzewing
¹⁷⁰ <i>Phaps elegans</i> (Temminck, 1809)	Brush Bronzewing
<i>Phaps histrionica</i> (Gould, 1841)	Flock Bronzewing (Flock Pigeon)
¹⁷¹ <i>Ocyphaps lophotes</i> (Temminck, 1822)	Crested Pigeon
¹⁷² <i>Geophaps plumifera</i> Gould, 1842	Spinifex Pigeon
<i>Geophaps plumifera plumifera</i> Gould, 1842	
<i>Geophaps plumifera ferruginea</i> (Gould, 1865)	

157 Non-breeding migrant.

158 Non-breeding migrant.

159 Some sight records probably refer to this subspecies.

160 Non-breeding migrant.

161 Non-breeding migrant.

162 Sometimes placed in the genus *Chlidonias*.

163 Sometimes placed in the genus *Chlidonias*.

164 Sometimes placed in the genus *Catharacta*.

165 Introduced in late 1800s.

166 Formerly in *Streptopelia*.

167 Specimens from Perth show characters of two subspecies: *S. c. chinensis* and *S. c. tigrina*.

168 Formerly in *Streptopelia*.

169 *C. i. longirostris* treated by Rasmussen & Anderton (2005) as a full species. However, Johnstone *et al.* (2014) shows that *C. i. longirostris* intergrades with *C. i. indica* in eastern Indonesia and they should be treated as conspecific.

170 Validity of ssp. *occidentalis* from south-west Western Australia is doubtful. Its range is continuous across the Great Australian Bight and geographic variation between Western Australian and South Australian populations is slight.

171 Subspecies *O. l. whitlocki* is not recognised in this almost continually distributed species; geographic variation is slight and clinal.

172 There is considerable geographic variation within this species with white-bellied, red-bellied and intermediate populations e.g. in Cape, Edgar and Grant ranges. These populations are probably best treated as polytopic subspecies. (Johnstone 1981, *Rec. West. Aust. Mus.* 9: 49–64).

<i>Geophaps smithii</i> (Jardine and Selby, 1830)	Partridge Pigeon
<i>Geophaps smithii smithii</i> (Jardine and Selby, 1830)	Partridge Pigeon (Eastern form)
<i>Geophaps smithii blaauwi</i> Mathews, 1912	Partridge Pigeon (Western form)
¹⁷³ <i>Petrophassa albipennis</i> Gould, 1841	White-quilled Rock Pigeon
<i>Geopelia cuneata</i> (Latham, 1802)	Diamond Dove
<i>Geopelia striata</i> (Linnaeus, 1766)	Zebra Dove
¹⁷⁴ <i>Geopelia striata placida</i> Gould, 1844	Peaceful Dove
¹⁷⁵ <i>Geopelia humeralis</i> (Temminck, 1821)	Bar-shouldered Dove
¹⁷⁶ <i>Caloenas nicobarica</i> (Linnaeus, 1758)	Nicobar Pigeon
<i>Ptilinopus regina</i> Swainson, 1825	Rose-crowned Fruit-Dove
<i>Ptilinopus regina ewingii</i> Gould, 1842	
<i>Ducula bicolor</i> (Scopoli, 1786)	Pied Imperial Pigeon
¹⁷⁷ <i>Ducula bicolor spilorrhoa</i> (G.R. Gray, 1858)	Torres Strait Pigeon (Torresian Imperial Pigeon)

Order CUCULIFORMES

Family CUCULIDAE Cuckoos

<i>Centropus bengalensis</i> (Gmelin, 1788)	Lesser Coucal
¹⁷⁸ <i>Centropus bengalensis sarasinorum</i> Stresemann, 1912	
<i>Centropus phasianinus</i> (Latham, 1801)	Pheasant Coucal
¹⁷⁹ <i>Centropus phasianinus phasianinus</i> (Latham, 1801)	
¹⁸⁰ <i>Centropus phasianinus highami</i> (Mathews, 1922)	
<i>Eudynamys orientalis</i> (Linnaeus, 1766)	Pacific Koel (Australian Koel)
<i>Eudynamys orientalis subcyanocephalus</i> Mathews, 1912	
<i>Scythrops novaehollandiae</i> Latham, 1790	Channel-billed Cuckoo
<i>Scythrops novaehollandiae novaehollandiae</i> Latham, 1790	
<i>Chrysococcyx basalis</i> (Horsfield, 1821)	Horsfield's Bronze Cuckoo
<i>Chrysococcyx osculans</i> (Gould, 1847)	Black-eared Cuckoo
<i>Chrysococcyx lucidus</i> (Gmelin, 1788)	Shining Bronze Cuckoo
<i>Chrysococcyx lucidus plagosus</i> (Latham, 1802)	
<i>Chrysococcyx minutillus</i> Gould, 1859	Little Bronze Cuckoo
<i>Chrysococcyx minutillus minutillus</i> Gould, 1859	
<i>Cacomantis pallidus</i> (Latham, 1802)	Pallid Cuckoo
<i>Cacomantis flabelliformis</i> (Latham, 1802)	Fan-tailed Cuckoo
<i>Cacomantis flabelliformis flabelliformis</i> (Latham, 1802)	
<i>Cacomantis variolosus</i> (Vigors and Horsfield, 1827)	Brush Cuckoo
<i>Cacomantis variolosus variolosus</i> (Vigors and Horsfield, 1827)	
<i>Cuculus saturatus</i> Blyth, 1843	Oriental Cuckoo
¹⁸¹ <i>Cuculus saturatus optatus</i> Gould, 1845	Horsfield's Cuckoo

Order STRIGIFORMES

Family TYTONIDAE Barn owls

<i>Tyto novaehollandiae</i> (Stephens, 1826)	Australian Masked Owl (Masked Owl)
<i>Tyto novaehollandiae novaehollandiae</i> (Stephens, 1826)	
<i>Tyto novaehollandiae kimberli</i> Mathews, 1912	

¹⁷³ Forming a superspecies with Chestnut-quilled Rock Pigeon *P. rufipennis* of Arnhem Land. Subspecies *P. a. boothi* is not recognised; it is the end of a cline in size and colouration.

¹⁷⁴ Only one ssp. *G. s. placida* recognised in Australia (Johnstone 1992, *West. Aust. Nat.* **19**(1)).

¹⁷⁵ No ssp. recognised (Johnstone 1990, *Rec. West. Aust. Mus. Suppl.* **32**: 1–120).

¹⁷⁶ Timor Sea 1989.

¹⁷⁷ Sometimes treated as a full species *D. spilorrhoa*.

¹⁷⁸ Ashmore Reef 2005.

¹⁷⁹ The ssp. *melanurus* of northern and north-western Australia is not recognised.

¹⁸⁰ Tentatively recognised for Pilbara and Gascoyne populations.

¹⁸¹ Sometimes treated as a full species *C. optatus*.

<i>Tyto alba</i> (Scopoli, 1769)	Barn Owl
182 <i>Tyto alba delicatula</i> (Gould, 1837)	Eastern Barn Owl
<i>Tyto longimembris</i> (Jerdon, 1839)	Eastern Grass Owl
Family STRIGIDAE Typical owls	
<i>Otus sunia</i> (Hodgson, 1836)	Oriental Scops Owl
183 <i>Otus sunia sunia</i> (Hodgson, 1836)	
<i>Ninox rufa</i> (Gould, 1846)	Rufous Owl
<i>Ninox rufa rufa</i> (Gould, 1846)	
184 <i>Ninox connivens</i> (Latham, 1802)	Barking Owl
<i>Ninox connivens connivens</i> (Latham, 1802)	
185 <i>Ninox boobook</i> (Latham, 1801)	Boobook Owl
<i>Ninox boobook boobook</i> (Latham, 1801)	Southern Boobook
<i>Ninox japonica</i> (Temminck and Schlegel, 1844)	Northern Boobook (Brown Hawk-Owl)
<i>Ninox japonica japonica</i> (Temminck and Schlegel, 1844)	

Order CAPRIMULGIFORMES**Family PODARGIDAE Frogmouths**

<i>Podargus strigoides</i> (Latham, 1802)	Tawny Frogmouth
<i>Podargus strigoides brachypterus</i> Gould, 1841	
<i>Podargus strigoides phalaenoides</i> Gould, 1840	

Family CAPRIMULGIDAE Nightjars

<i>Eurostopodus argus</i> Hartert, 1892	Spotted Nightjar
186 <i>Caprimulgus jotaka</i> Temminck and Schlegel, 1847	Grey Nightjar
187 <i>Caprimulgus affinis</i> Horsfield, 1821	Savanna Nightjar

Order APODIFORMES**Family AEGOTHELIIDAE Owlet-nightjars**

<i>Aegotheles cristatus</i> (Shaw, 1790)	Australian Owlet-nightjar
<i>Aegotheles cristatus cristatus</i> (Shaw, 1790)	
188 <i>Aegotheles cristatus leucogaster</i> Gould, 1844	

Family APODIDAE Swifts

<i>Hirundapus caudacutus</i> (Latham, 1801)	White-throated Needletail
<i>Hirundapus caudacutus caudacutus</i> (Latham, 1801)	
189 <i>Apus pacificus</i> (Latham, 1801)	Pacific Swift (Fork-tailed Swift)
<i>Apus pacificus pacificus</i> (Latham, 1801)	
190 <i>Apus nipalensis</i> (Hodgson, 1837)	House Swift

Order CORACIIFORMES**Family CORACIIDAE Rollers**

<i>Eurystomus orientalis</i> (Linnaeus, 1766)	Dollarbird
<i>Eurystomus orientalis pacificus</i> (Latham, 1802)	

182 Sometimes treated as a full species *T. delicatula*.

183 Between Barrow I. and mainland 2013.

184 Most of this species variation in size and colouration is clinal and, as such, the recognition of *addenda* in the south-west and *occidentalis* in north Western Australia is not warranted.185 Only one ssp. in Western Australia is recognised. *N. b. ocellata* is treated as a synonym of *N. b. boobook*.186 Ashmore Reef 2003. No specimens available. Sometimes treated as a ssp. of *C. indicus*.

187 Ashmore Reef 2006. Subspecies not identified and migratory and local Indonesian ssp. are possible vagrants.

188 The ssp. *leucogaster* is recognised for the north-eastern and southern Kimberley population (Johnstone and Storr 1998).

189 Non-breeding migrant.

190 Timor Sea 1988 and Broome 1998. Currently expanding its range in Lesser Sundas, Indonesia (Johnstone *et al.* 2014). Sometimes treated as conspecific with the Little Swift *Apus affinis*.

Family ALCEDINIDAE Kingfishers

- **Dacelo novaeguineae* (Hermann, 1783) Laughing Kookaburra
 **Dacelo novaeguineae novaeguineae* (Hermann, 1783)
Dacelo leachii Vigors and Horsfield, 1827 Blue-winged Kookaburra
Dacelo leachii leachii Vigors and Horsfield, 1827
¹⁹¹*Halcyon pileata* (Boddaert, 1783) Black-capped Kingfisher
¹⁹²*Todiramphus macleayii* (Jardine and Selby, 1830) Forest Kingfisher
Todiramphus chloris (Boddaert, 1783) Collared Kingfisher
Todiramphus chloris sordidus (Gould, 1842)
Todiramphus chloris pilbara (Johnstone, 1983) Pilbara Collared Kingfisher
Todiramphus sanctus (Vigors and Horsfield, 1827) Sacred Kingfisher
Todiramphus sanctus sanctus (Vigors and Horsfield, 1827)
Todiramphus pyrrhopygus (Gould, 1841) Red-backed Kingfisher
¹⁹³*Ceyx azureus* (Latham, 1802) Azure Kingfisher
Ceyx azureus ruficollaris (Bankier, 1841)

Family MEROPIDAE Bee-eaters

- Merops ornatus* Latham, 1802 Rainbow Bee-eater

Order BUCEROTIFORMES**Family UPUPIDAE Hoopoe**

- Upupa epops* Linnaeus, 1758 Eurasian Hoopoe
¹⁹⁴*Upupa epops [saturata]* Lönberg, 1909

Order FALCONIFORMES**Family FALCONIDAE Falcons and kestrels**

- Falco cenchroides* Vigors and Horsfield, 1827 Australian Kestrel (Nankeen Kestrel)
Falco cenchroides cenchroides Vigors and Horsfield, 1827
¹⁹⁵*Falco subbuteo* Linnaeus, 1758 Eurasian Hobby
Falco longipennis Swainson, 1837 Australian Hobby
¹⁹⁶*Falco longipennis longipennis* Swainson, 1837
¹⁹⁷*Falco berigora* Vigors and Horsfield, 1827 Brown Falcon
Falco berigora berigora Vigors and Horsfield, 1827
Falco hypoleucus Gould, 1841 Grey Falcon
Falco subniger G.R. Gray, 1843 Black Falcon
Falco peregrinus Tunstall, 1771 Peregrine Falcon
¹⁹⁸*Falco peregrinus [japonensis]* Gmelin, 1788
¹⁹⁹*Falco peregrinus macropus* Swainson, 1837

Order PSITTACIFORMES**Family CACATUIDAE Cockatoos**

- Calyptorhynchus banksii* (Latham, 1790) Red-tailed Black Cockatoo
Calyptorhynchus banksii macrorhynchus Gould, 1843 Northern Red-tailed Black Cockatoo
Calyptorhynchus banksii samueli Mathews, 1917 Inland Red-tailed Black Cockatoo
Calyptorhynchus banksii naso Gould, 1837 Forest Red-tailed Black Cockatoo

¹⁹¹ North Burracoppin 2005.

¹⁹² One record, a bird photographed on Jabiru Venture, Timor Sea, April 1990.

¹⁹³ Sometimes placed in the genus *Alcedo*.

¹⁹⁴ Roebuck Plains 2011. No specimens available; ssp. not determined, but most likely *saturata*.

¹⁹⁵ Scott Reef – Rowley Shoals 1997. South Lake 2016. Most likely the nominate subspecies.

¹⁹⁶ Geographic variation in this continuously distributed species is slight and clinal; ssp. *murchisonianus* is not recognised.

¹⁹⁷ This species is semi-nomadic with continuous distribution; the various colour forms can only be grouped in arbitrary fashion and therefore only one ssp. recognised for Australia.

¹⁹⁸ No specimens examined.

¹⁹⁹ *F. p. submelanogenys* of south-western WA is not recognised.

²⁰⁰ <i>Calyptorhynchus latirostris</i> Carnaby, 1948	Carnaby's Cockatoo
²⁰¹ <i>Calyptorhynchus baudinii</i> Lear, 1832	Baudin's Cockatoo
²⁰² <i>Cacatua leadbeateri</i> (Vigors, 1831)	Major Mitchell's Cockatoo
²⁰³ <i>Cacatua roseicapilla</i> Vieillot, 1817	Galah
<i>Cacatua roseicapilla roseicapilla</i> Vieillot, 1817	
<i>Cacatua roseicapilla albiceps</i> (Schodde, 1989)	
* <i>Cacatua tenuirostris</i> (Kuhl, 1820)	Eastern Long-billed Corella
<i>Cacatua pastinator</i> (Gould, 1841)	Western Long-billed Corella
<i>Cacatua pastinator pastinator</i> (Gould, 1841)	Muir's Corella
²⁰⁴ <i>Cacatua pastinator butleri</i> Ford, 1987	Butler's Corella
<i>Cacatua sanguinea</i> Gould, 1843	Little Corella
<i>Cacatua sanguinea sanguinea</i> Gould, 1843	
<i>Cacatua sanguinea westralensis</i> (Mathews, 1917)	Western Little Corella
<i>Cacatua galerita</i> (Latham, 1790)	Sulphur-crested Cockatoo
²⁰⁵ * <i>Cacatua galerita galerita</i> (Latham, 1790)	
<i>Cacatua galerita fitzroyi</i> (Mathews, 1912)	
<i>Nymphicus hollandicus</i> (Kerr, 1792)	Cockatiel
Family PSITTACIDAE Parrots	
²⁰⁶ * <i>Trichoglossus moluccanus</i> (Gmelin, 1788)	Rainbow Lorikeet
<i>Trichoglossus rubritorquis</i> Vigors and Horsfield, 1827	Red-collared Lorikeet
²⁰⁷ <i>Trichoglossus versicolor</i> Lear, 1831	Varied Lorikeet
²⁰⁸ * <i>Glossopsitta concinna</i> (Shaw, 1791)	Musk Lorikeet
²⁰⁹ <i>Parvipsitta porphyrocephala</i> (Dietrichsen, 1837)	Purple-crowned Lorikeet
²¹⁰ <i>Platycercus spurius</i> (Kuhl, 1820)	Red-capped Parrot
²¹¹ <i>Platycercus zonarius</i> (Shaw, 1805)	Australian Ringneck
<i>Platycercus zonarius zonarius</i> (Shaw, 1805)	Port Lincoln Parrot
²¹² <i>Platycercus zonarius semitorquatus</i> (Quoy and Gaimard, 1830)	Twenty-eight Parrot
²¹³ <i>Platycercus venustus</i> (Kuhl, 1820)	Northern Rosella
<i>Platycercus icterotis</i> (Temminck and Kuhl, 1820)	Western Rosella
<i>Platycercus icterotis icterotis</i> (Temminck and Kuhl, 1820)	
<i>Platycercus icterotis xanthogenys</i> Salvadori, 1891	

²⁰⁰ Sometimes placed in the genus *Zanda*.

²⁰¹ Sometimes placed in the genus *Zanda*.

²⁰² Sometimes placed in the genus *Lophochroa*. Subspecies taxonomic status requires confirmation.

²⁰³ Type locality for the Galah remains an enigma; described by Vieillot (1817) as *Cacatua roseicapilla* from a specimen collected during Baudin Expedition 1800–1803 labelled 'dans les Indes'. Mathews (1912) Novit. Zool. 18: determined this location to be New South Wales, however Schodde (1988) *Canberra Bird Notes* 13:4 believed the type to be probably from Western Australia, Shark Bay area. Recent evidence confirms that a specimen was collected by the Baudin Expedition in Shark Bay in August/September 1801, thus the nominate ssp. is now assigned to populations south of the Kimberley and ssp. *albiceps* for northern and eastern Australia; ssp. *kuhli* is not recognised. Sometimes placed in the genus *Eolophus*.

²⁰⁴ The provenance and subspecific identity of the type specimen of *Licmetis tenuirostris derbyi* Mathews 1916, are in doubt (most probably an immature *C. p. pastinator*) and until resolved the most appropriate name for the mid Western Australian (wheatbelt) population is *Cacatua pastinator butleri* Ford 1987.

²⁰⁵ Small colonies established in southwest of Western Australia since 1955.

²⁰⁶ Exotic species established in Perth area since 1968.

²⁰⁷ Sometimes placed in the genus *Psitteuteles*.

²⁰⁸ Introduced into Perth area in 1970s.

²⁰⁹ Formerly in *Glossopsitta*. See Schweizer *et al.* 2015.

²¹⁰ The sole member of the subgenus *Purpureicephalus*. See Johnstone and Benken (2009) for hybridisation with *P. zonarius*.

²¹¹ The sole member of the subgenus *Barnardiuss*.

²¹² Known to occasionally hybridise with *P. spurius* on the Swan Coastal Plain.

²¹³ Distribution is continuous between Western Australia and Northern Territory populations; variation is slight and clinal (across Joseph Bonaparte Gulf region) and thus inadequate for recognition of subspecies *hilli*.

²¹⁴ <i>Platycercus narethae</i> (H.L. White, 1921)	Naretha Blue Bonnet
²¹⁵ <i>Platycercus varius</i> (Clark, 1910)	Mulga Parrot
²¹⁶ <i>Neophema bourkii</i> (Gould, 1841)	Bourke's Parrot
<i>Neophema elegans</i> (Gould, 1837)	Elegant Parrot
<i>Neophema petrophila</i> (Gould, 1841)	Rock Parrot
²¹⁷ <i>Neophema petrophila petrophila</i> (Gould, 1841)	
<i>Neophema splendida</i> (Gould, 1841)	Scarlet-chested Parrot
<i>Melopsittacus undulatus</i> (Shaw, 1805)	Budgerigar
<i>Pezoporus wallicus</i> (Kerr, 1792)	Ground Parrot
²¹⁸ <i>Pezoporus wallicus flaviventris</i> North, 1911	Western Ground Parrot
<i>Pezoporus occidentalis</i> (Gould, 1861)	Night Parrot
²¹⁹ <i>Aprosmictus erythropterus</i> (Gmelin, 1788)	Red-winged Parrot
²²⁰ <i>Polytelis anthopeplus</i> (Lear, 1831)	Regent Parrot
<i>Polytelis anthopeplus westralsis</i> Mathews, 1915	
<i>Polytelis alexandrae</i> Gould, 1863	Princess Parrot
^{221*} <i>Psittacula krameri</i> (Scopoli, 1769)	Rose-ringed Parakeet (Indian Ringnecked Parrot)

Order PASSERIFORMES

Family PITTIDAE Pittas

²²² <i>Pitta sordida</i> (P.L.S. Müller, 1776)	Hooded Pitta
²²³ <i>Pitta sordida cucullata</i> Hartlaub, 1843	
<i>Pitta nymphula</i> Temminck and Schlegel, 1850	Fairy Pitta
²²⁴ <i>Pitta moluccensis</i> (P.L.S. Müller, 1776)	Blue-winged Pitta
²²⁵ <i>Pitta iris</i> Gould, 1842	Rainbow Pitta
²²⁶ <i>Pitta iris johnstoneiana</i> Schodde and Mason, 1999	

Family ATRICHORNITHIDAE Scrub-birds

<i>Atrichornis clamosus</i> (Gould, 1844)	Noisy Scrub-bird
---	------------------

Family PTILONORHYNCHIDAE Bowerbirds

²²⁷ <i>Ptilonorhynchus nuchalis</i> Jardine and Selby, 1830	Great Bowerbird
<i>Ptilonorhynchus nuchalis nuchalis</i> Jardine and Selby, 1830	

²¹⁴ Sometimes placed in the genus *Northiella*.

²¹⁵ Sometimes placed in the genus *Psephotus*.

²¹⁶ Sometimes placed in the genus *Neopsephotus*.

²¹⁷ The ssp. *zietzi* from coastal South Australia is at best very weakly defined.

²¹⁸ Sometimes treated as a full species.

²¹⁹ Forming a superspecies with *A. jonquillaceus* of Wetar and Timor.

²²⁰ The type locality for the Regent Parrot is uncertain and the type specimens appear to have been lost. The names *Palaeornis anthopeplus* Lear 1831 and *P. melanura* Lear 1932 were based on a female and male respectively, published on plates in his *Illustrations of the family Psittacidae, or Parrots* 1830–1832, without description or source. Mathews (1912) presumed the type locality was New South Wales and named *Polytelis anthopeplus westralsis* from southwest Australia. Schodde 1993 (*Bull. Br. Ornithol. Club* 113: 44–47) noted that Lear's plate matched best with the western form and designated a neotype from Bolgart, Western Australia and provided the name *P. a. monarchoides* for the eastern Australian populations. Mees 2004 (*Zool. Med. Leiden* 78(10) 27 viii 2004: 205–208) on the other hand, argued that Schodde's action was invalid and that the type locality should remain as New South Wales.

²²¹ Aviary escapees.

²²² Barrow Island 2010.

²²³ Barrow Island 2010.

²²⁴ Non-breeding migrant.

²²⁵ Forming a superspecies with *P. versicolor* of eastern Australia and *P. elegans* of Indonesia.

²²⁶ Confined to west Kimberley.

²²⁷ Sometimes placed in the genus *Chlamydera*.

²²⁸ <i>Ptilonorhynchus maculatus</i> (Gould, 1837)	Spotted Bowerbird
²²⁹ <i>Ptilonorhynchus maculatus guttatus</i> (Gould, 1862)	Western Bowerbird
Family CLIMACTERIDAE Treecreepers	
<i>Climacteris affinis</i> Blyth, 1864	White-browed Treecreeper
<i>Climacteris affinis superciliosus</i> North, 1895	
²³⁰ <i>Climacteris rufus</i> Gould, 1841	Rufous Treecreeper
<i>Climacteris melanurus</i> Gould, 1843	Black-tailed Treecreeper
<i>Climacteris melanurus melanurus</i> Gould, 1843	
<i>Climacteris melanurus wellsi</i> Ogilvie-Grant, 1909	
Family MALURIDAE Grasswrens, fairy-wrens and emu-wrens	
<i>Malurus lamberti</i> Vigors and Horsfield, 1827	Variegated Fairy-wren
<i>Malurus lamberti rogersi</i> Mathews, 1912	
<i>Malurus lamberti assimilis</i> North, 1901	
²³¹ <i>Malurus lamberti bernieri</i> Ogilvie-Grant, 1909	Shark Bay Variegated Fairy-wren
<i>Malurus pulcherrimus</i> Gould, 1844	Blue-breasted Fairy-wren
<i>Malurus elegans</i> Gould, 1837	Red-winged Fairy-wren
<i>Malurus splendens</i> (Quoy and Gaimard, 1830)	Splendid Fairy-wren
<i>Malurus splendens splendens</i> (Quoy and Gaimard, 1830)	
²³² <i>Malurus splendens callainus</i> Gould, 1867	
<i>Malurus coronatus</i> Gould, 1858	Purple-crowned Fairy-wren
<i>Malurus coronatus coronatus</i> Gould, 1858	
<i>Malurus melanocephalus</i> (Latham, 1802)	Red-backed Fairy-wren
<i>Malurus melanocephalus cruentatus</i> Gould, 1840	
<i>Malurus leucopterus</i> Dumont, 1824	White-winged Fairy-wren
²³³ <i>Malurus leucopterus leucopterus</i> Dumont, 1824	Dirk Hartog Island Black and White Fairy-wren
²³⁴ <i>Malurus leucopterus edouardi</i> Campbell, 1901	Barrow Island Black and White Fairy-wren
<i>Malurus leucopterus leuconotus</i> Gould, 1865	
<i>Stipiturus malachurus</i> (Shaw, 1798)	Southern Emu-wren
<i>Stipiturus malachurus westernensis</i> Campbell, 1912	
²³⁵ <i>Stipiturus malachurus hartogi</i> Carter, 1916	Dirk Hartog Island Emu-wren
<i>Stipiturus ruficeps</i> Campbell, 1899	Rufous-crowned Emu-wren
²³⁶ <i>Stipiturus ruficeps ruficeps</i> Campbell, 1899	
<i>Amytornis housei</i> (Milligan, 1902)	Black Grasswren
<i>Amytornis striatus</i> (Gould, 1840)	Striated Grasswren
<i>Amytornis striatus striatus</i> (Gould, 1840)	
<i>Amytornis striatus whitei</i> Mathews, 1910	Rufous Grasswren
²³⁷ <i>Amytornis textilis</i> (Dumont, 1824)	Thick-billed Grasswren
<i>Amytornis textilis textilis</i> (Dumont, 1824)	Western Grasswren
<i>Amytornis purnelli</i> (Mathews, 1914)	Dusky Grasswren
Family MELIPHAGIDAE Honeyeaters and chats	
²³⁸ <i>Sugomel niger</i> (Gould, 1838)	Black Honeyeater
<i>Myzomela obscura</i> Gould, 1843	Dusky Honeyeater (Dusky Myzomela)
<i>Myzomela obscura obscura</i> Gould, 1843	

²²⁸ Sometimes placed in the genus *Chlamydera*.

²²⁹ Sometimes treated as a full species *P. guttatus*. Subspecies *P. guttatus carteri* of North-west Cape is not recognised.

²³⁰ Forming a superspecies with Brown Treecreeper *C. picumnus* of eastern Australia.

²³¹ Confined to Bernier and Dorre Is.

²³² Sometimes referred to as *M. s. musgravi* Mathews 1922.

²³³ Confined to Dirk Hartog I.

²³⁴ Confined to Barrow I.

²³⁵ Confined to Dirk Hartog I.

²³⁶ Treated as conspecific with *S. r. mallee* Campbell 1908.

²³⁷ South-western populations *A. t. macrourus* are extinct (Johnstone and Storr 2004).

²³⁸ Following leCroy (2011).

²³⁹ <i>Myzomela erythrocephala</i> Gould, 1840	Red-headed Honeyeater (Red-headed Myzomela)
<i>Myzomela erythrocephala erythrocephala</i> Gould, 1840	
²⁴⁰ <i>Glyciphila melanops</i> (Latham, 1802)	Tawny-crowned Honeyeater
<i>Glyciphila melanops melanops</i> (Latham, 1802)	
<i>Acanthorhynchus superciliosus</i> Gould, 1837	Western Spinebill
<i>Certhionyx variegatus</i> Lesson, 1830	Pied Honeyeater
<i>Cissomela pectoralis</i> (Gould, 1841)	Banded Honeyeater
²⁴¹ <i>Lichmera indistincta</i> (Vigors and Horsfield, 1827)	Brown Honeyeater
<i>Lichmera indistincta indistincta</i> (Vigors and Horsfield, 1827)	
²⁴² <i>Phylidonyris novaehollandiae</i> (Latham, 1790)	New Holland Honeyeater
<i>Phylidonyris novaehollandiae longirostris</i> (Gould, 1846)	
<i>Phylidonyris niger</i> (Bechstein, 1811)	White-cheeked Honeyeater
<i>Phylidonyris niger gouldii</i> (Schlegel, 1872)	
<i>Grantiella picta</i> (Gould, 1838)	Painted Honeyeater
<i>Philemon citreogularis</i> (Gould, 1837)	Little Friarbird
²⁴³ <i>Philemon citreogularis citreogularis</i> (Gould, 1837)	
<i>Philemon argenticeps</i> (Gould, 1840)	Silver-crowned Friarbird
<i>Philemon argenticeps argenticeps</i> (Gould, 1840)	
²⁴⁴ <i>Melithreptus cyanotis</i> (Latham, 1802)	Blue-faced Honeyeater
<i>Melithreptus cyanotis albipennis</i> (Gould, 1840)	
<i>Melithreptus gularis</i> (Gould, 1837)	Black-chinned Honeyeater
<i>Melithreptus gularis laetior</i> Gould, 1875	
<i>Melithreptus brevirostris</i> (Vigors and Horsfield, 1827)	Brown-headed Honeyeater
<i>Melithreptus brevirostris leucogenys</i> Milligan, 1903	
<i>Melithreptus albogularis</i> Gould, 1848	White-throated Honeyeater
<i>Melithreptus albogularis albogularis</i> Gould, 1848	
²⁴⁵ <i>Melithreptus chloropsis</i> Gould, 1848	Western White-naped Honeyeater
<i>Epthianura tricolor</i> Gould, 1841	Crimson Chat
<i>Epthianura aurifrons</i> Gould, 1838	Orange Chat
²⁴⁶ <i>Epthianura crocea</i> Castelnau and Ramsay, 1877	Yellow Chat
<i>Epthianura albifrons</i> (Jardine and Selby, 1828)	White-fronted Chat
<i>Conopophila rufogularis</i> (Gould, 1843)	Rufous-throated Honeyeater
²⁴⁷ <i>Lacustoica whitei</i> North, 1910	Grey Honeyeater
<i>Ramsayornis fasciatus</i> (Gould, 1843)	Bar-breasted Honeyeater
<i>Acanthagenys rufogularis</i> Gould, 1838	Spiny-cheeked Honeyeater
²⁴⁸ <i>Anthochaera lunulata</i> Gould, 1838	Western Little Wattlebird (Western Wattlebird)
<i>Anthochaera carunculata</i> (Shaw, 1790)	Red Wattlebird
²⁴⁹ <i>Lichenostomus leucotis</i> (Latham, 1802)	White-eared Honeyeater
<i>Lichenostomus leucotis novaenorciae</i> (Milligan, 1904)	Western White-eared Honeyeater

²³⁹ Forming a superspecies with *M. dammermani*, *M. kuehni* and possibly *M. vulnerata*, all of the Lesser Sundas.

²⁴⁰ Sometimes placed in the genus *Phylidonyris*.

²⁴¹ Indonesian Honeyeater *L. i. limbata* treated as the Wallacean ssp. of the Brown Honeyeater.

²⁴² Comprising with the White-cheeked Honeyeater *P. niger*, the subgenus *Meliornis*.

²⁴³ Only one ssp. recognised for mainland Australia. Geographic variation in this continuously distributed species is clinal, with the nominate subspecies intergrading with *sordidus* over most of north Queensland. While some of the variation is undoubtedly geographic, it hardly amounts to subspeciation.

²⁴⁴ Sometimes placed in the genus *Entomyzon*, but retained here in *Melithreptus* based on morphology, behaviour and eggs.

²⁴⁵ Forming a superspecies with *M. lunatus* and *M. affinis* of eastern Australia (Johnstone and Storr 2004).

²⁴⁶ No ssp. recognised (Ford and Parker 1974).

²⁴⁷ Sometimes placed in the genus *Conopophila*.

²⁴⁸ The eastern and western populations of *Anthochaera chrysoptera* (*chrysoptera* and *lunulata* respectively) were previously treated as conspecific.

²⁴⁹ Sometimes placed in the genus *Nesoptilotis*.

<i>Lichenostomus cratitius</i> (Gould, 1841)	Purple-gaped Honeyeater
<i>Lichenostomus cratitius occidentalis</i> Cabanis, 1851	
²⁵⁰ <i>Manorina flavigula</i> (Gould, 1840)	Yellow-throated Miner
²⁵¹ <i>Purnella albifrons</i> (Gould, 1841)	White-fronted Honeyeater
<i>Stomiopera unicolor</i> (Gould, 1843)	White-gaped Honeyeater
²⁵² <i>Gavicalis virescens</i> (Vieillot, 1817)	Singing Honeyeater
²⁵³ <i>Gavicalis virescens virescens</i> (Vieillot, 1817)	
<i>Ptilotula flavescens</i> (Gould, 1840)	Yellow-tinted Honeyeater
<i>Ptilotula flavescens flavescens</i> (Gould, 1840)	
<i>Ptilotula keartlandi</i> (North, 1895)	Grey-headed Honeyeater
²⁵⁴ <i>Ptilotula plumula</i> (Gould, 1841)	Grey-fronted Honeyeater
<i>Ptilotula ornata</i> (Gould, 1838)	Yellow-plumed Honeyeater
<i>Ptilotula penicillata</i> (Gould, 1837)	White-plumed Honeyeater
²⁵⁵ <i>Ptilotula penicillata carteri</i> (A.J. Campbell, 1899)	Carter's White-plumed Honeyeater
<i>Meliphaga albilineata</i> (White, 1917)	White-lined Honeyeater
²⁵⁶ <i>Meliphaga albilineata fordiana</i> Schodde, 1989	Kimberley White-lined Honeyeater
Family DASYORNITHIDAE Bristlebirds	
<i>Dasyornis longirostris</i> Gould, 1841	Western Bristlebird
<i>Dasyornis broadbenti</i> (McCoy, 1867)	Rufous Bristlebird
<i>Dasyornis broadbenti litoralis</i> (Milligan, 1902)	
Family PARDALOTIDAE Pardalotes	
<i>Pardalotus punctatus</i> (Shaw, 1792)	Spotted Pardalote
<i>Pardalotus punctatus punctatus</i> (Shaw, 1792)	
²⁵⁷ <i>Pardalotus punctatus xanthopyge</i> McCoy, 1866	Yellow-rumped Pardalote
<i>Pardalotus rubricatus</i> Gould, 1838	Red-browed Pardalote
<i>Pardalotus rubricatus rubricatus</i> Gould, 1838	
²⁵⁸ <i>Pardalotus striatus</i> (Gmelin, 1789)	Striated Pardalote
<i>Pardalotus striatus uropygialis</i> Gould, 1840	
²⁵⁹ <i>Pardalotus striatus murchisoni</i> Mathews, 1912	
²⁶⁰ <i>Pardalotus striatus westraliensis</i> Mathews, 1912	
Family ACANTHIZIDAE Gerygones, thornbills, scrubwrens and whitefaces	
²⁶¹ <i>Calamanthus cautus</i> (Gould, 1843)	Shy Groundwren (Shy Heathwren)
²⁶² <i>Calamanthus cautus whitlocki</i> (Mathews, 1912)	

²⁵⁰ Schodde and Mason (1999) recognised three ssp. in WA. The gradation from dark southern *M. f. "obscura"* type birds to paler *M. f. "wayensis"* is a smooth cline and thus inadequate for recognition of subspecies (Johnstone and Storr 2004).

²⁵¹ Previously placed in the genus *Phylidonyris*.

²⁵² Forming a superspecies with the Mangrove Honeyeater *L. versicolor* of eastern Australia and New Guinea.

²⁵³ Only the nominate ssp. recognised in Western Australia with a broad cline between southern *virescens* and northern "*forresti*" forms.

²⁵⁴ Northern Australian populations sometimes treated as a separate ssp. "*planasi*".

²⁵⁵ The ssp. *carteri* is recognised for the isolated mid-western population, however, there is a number of other disjunct populations in WA (Johnstone and Storr 2004). Those in south-west Kimberley differ only slightly from northern *carteri* and the name "*calconi*" could be used for them, whereas north-eastern WA populations are merely the northern end of a north-south cline beginning in eastern Australia.

²⁵⁶ Sometimes treated as a full species *M. fordiana*.

²⁵⁷ Tentatively treated as conspecific with *P. punctatus* based on evidence of hybridisation in southeast Australia (Schodde and Mason 1999), but little or no evidence of hybridisation in WA (Johnstone and Storr 2004).

²⁵⁸ For subspecies recognition see Johnstone and Storr 2004.

²⁵⁹ Subspecies *murchisoni* and *westraliensis* recognised (Johnstone and Storr 2004).

²⁶⁰ Subspecies *westraliensis* recognised for populations breeding in south-west of Western Australia (Johnstone and Storr 2004).

²⁶¹ Previously in the genus *Hylacola*.

²⁶² Tentatively recognised, but taxonomic status requires confirmation.

²⁶³ <i>Calamanthus campestris</i> (Gould, 1841)	Rufous Fieldwren
<i>Pyrrholaemus brunneus</i> Gould, 1841	Redthroat
<i>Sericornis frontalis</i> (Vigors and Horsfield, 1827)	White-browed Scrubwren
<i>Sericornis frontalis maculatus</i> Gould, 1847	
<i>Sericornis frontalis balstoni</i> Ogilvie-Grant, 1909	
²⁶⁴ <i>Smicrornis brevirostris</i> (Gould, 1838)	Weebill
<i>Gerygone levigaster</i> Gould, 1843	Mangrove Gerygone
<i>Gerygone levigaster levigaster</i> Gould, 1843	
<i>Gerygone fusca</i> (Gould, 1838)	Western Gerygone
<i>Gerygone fusca fusca</i> (Gould, 1838)	
²⁶⁵ <i>Gerygone fusca mungi</i> Mathews, 1912	Desert Gerygone
²⁶⁶ <i>Gerygone tenebrosa</i> (Hall, 1901)	Dusky Gerygone
<i>Gerygone magnirostris</i> Gould, 1843	Large-billed Gerygone
<i>Gerygone magnirostris magnirostris</i> Gould, 1843	
<i>Gerygone chloronota</i> Gould, 1843.....	Green-backed Gerygone
²⁶⁷ <i>Gerygone chloronota chloronota</i> Gould, 1843	
<i>Gerygone olivacea</i> (Gould, 1838)	White-throated Gerygone
<i>Gerygone olivacea rogersi</i> Mathews, 1911	
²⁶⁸ <i>Acanthiza apicalis</i> Gould, 1847	Broad-tailed Thornbill (Inland Thornbill)
²⁶⁹ <i>Acanthiza apicalis apicalis</i> Gould, 1847	
<i>Acanthiza uropygialis</i> Gould, 1838	Chestnut-rumped Thornbill
<i>Acanthiza inornata</i> Gould, 1841	Western Thornbill
<i>Acanthiza iredalei</i> Mathews, 1911	Samphire Thornbill (Slender-billed Thornbill)
<i>Acanthiza iredalei iredalei</i> Mathews, 1911	
<i>Acanthiza chrysorrhoa</i> (Quoy and Gaimard, 1830)	Yellow-rumped Thornbill
²⁷⁰ <i>Acanthiza chrysorrhoa chrysorrhoa</i> (Quoy and Gaimard, 1830)	
<i>Acanthiza robustirostris</i> Milligan, 1903	Slaty-backed Thornbill
²⁷¹ <i>Aphelocephala leucopsis</i> (Gould, 1841)	Southern Whiteface
<i>Aphelocephala leucopsis leucopsis</i> (Gould, 1841)	
<i>Aphelocephala leucopsis castaneiventris</i> (Milligan, 1903)	
<i>Aphelocephala nigricincta</i> (North, 1895)	Banded Whiteface
Family POMATOSTOMIDAE Australasian babblers	
<i>Pomatostomus temporalis</i> (Vigors and Horsfield, 1827)	Grey-crowned Babbler
<i>Pomatostomus temporalis rubeculus</i> (Gould, 1840)	
²⁷² <i>Pomatostomus superciliosus</i> (Vigors and Horsfield, 1827)	White-browed Babbler

²⁶³ Schodde and Mason (1999) separated the southwestern population as a distinct species, the Western Fieldwren. Its distribution is continuous in WA (Johnstone and Storr 2004) and there is a smooth cline in variation between the southern olive and northern rufous forms; furthermore the calls are identical. Previously treated as a subspecies of Striated Fieldwren *C. fuliginosus*.

²⁶⁴ Not unexpectedly this sedentary bird undergoes some geographic variation throughout its immense range, but this hardly amounts to subspeciation.

²⁶⁵ Closely related to the Western Gerygone and possibly only a well-marked subspecies.

²⁶⁶ Variation in WA populations is clinal and thus inadequate for recognition of subspecies (Johnstone and Storr 2004).

²⁶⁷ Geographic variation in WA populations is clinal without steps and inadequate for recognition of subspecies (Johnstone and Storr 2004).

²⁶⁸ Forming a superspecies with the Brown Thornbill *A. pusilla* of Tasmania and wetterparts of eastern Australia.

²⁶⁹ Only the nominate form recognised in Western Australia with a broad cline between southern nominate and northern "whitlocki" forms.

²⁷⁰ Only the nominate ssp. recognised in Western Australia.

²⁷¹ Subspecies *leucopsis* and *castaneiventris* intergrade in a narrow zone through the Gibson Desert in Western Australia (Johnstone and Storr 2004).

²⁷² A number of ssp. have been proposed for this species including "*ashbyi*" and "*centralis*" in Western Australia (Shodde and Mason 1999), but their descriptions are unconvincing, geographic variation is clinal and the distribution map is erroneous.

Family PSOPHODIDAE Quail-thrushes, whipbirds and wedgebills

- Psophodes nigrogularis* Gould, 1844 Western Whipbird
²⁷³*Psophodes nigrogularis nigrogularis* Gould, 1844
Psophodes occidentalis (Mathews, 1912) Western Wedgebill (Chiming Wedgebill)
²⁷⁴*Cinclosoma clarum* Morgan, 1926 Western Chestnut Quail-thrush (Copperback Quail-thrush)
²⁷⁵*Cinclosoma alisteri* Mathews, 1910 Nullarbor Quail-thrush
Cinclosoma marginatum Sharpe, 1883 Western Quail-thrush

Family ARTAMIDAE Woodswallows

- Artamus leucorynchus* (Linnaeus, 1771) White-breasted Woodswallow
Artamus leucorynchus leucopygialis Gould, 1842
Artamus personatus (Gould, 1841) Masked Woodswallow
Artamus superciliosus (Gould, 1837) White-browed Woodswallow
²⁷⁶*Artamus cinereus* Vieillot, 1817 Black-faced Woodswallow
Artamus cinereus melanops Gould, 1865
²⁷⁷*Artamus cyanopterus* (Latham, 1802) Dusky Woodswallow
²⁷⁸*Artamus minor* Vieillot, 1817 Little Woodswallow

Family CRACTICIDAE Butcherbirds, Australian magpies and currawongs

- Cracticus quoyi* (Lesson and Garnot, 1827) Black Butcherbird
Cracticus quoyi spaldingi Masters, 1878
Cracticus torquatus (Latham, 1802) Grey Butcherbird
Cracticus torquatus torquatus (Latham, 1802)
Cracticus torquatus latens Ford, 1979
²⁷⁹*Cracticus argenteus* Gould, 1841 Silver-backed Butcherbird
²⁸⁰*Cracticus argenteus latens* Ford, 1979
²⁸¹*Cracticus nigrogularis* (Gould, 1837) Pied Butcherbird
²⁸²*Cracticus tibicen* (Latham, 1802) Australian Magpie
Cracticus tibicen tibicen (Latham, 1802) Black-backed Magpie
Cracticus tibicen dorsalis (Campbell, 1895) White-backed Magpie
Strepera versicolor (Latham, 1802) Grey Currawong
Strepera versicolor plumbea Gould, 1846

Family CAMPEPHAGIDAE Cuckoo-shrikes and trillers

- Coracina maxima* (Rüppell, 1839) Ground Cuckoo-shrike
Coracina novaehollandiae (Gmelin, 1789) Black-faced Cuckoo-shrike
Coracina novaehollandiae subpallida Mathews, 1912
²⁸³*Coracina novaehollandiae melanops* (Latham, 1802)

²⁷³ *P. n. oberon* is not recognised (Johnstone and Storr 2004).

²⁷⁴ Formerly *C. castanotum clarum*.

²⁷⁵ Sometimes treated as conspecific with Cinnamon Quail-thrush *C. cinnamomeum*, but the two forms are best treated as allospecies.

²⁷⁶ There is little firm evidence that the type came from opposite the Archipelago of the Recherche in south-west Australia, but far more likely from Timor (Peters 1962 and contra Mees 1968; Schodde and Mason 1999).

²⁷⁷ The differences between eastern nominate "cyanopterus" and western "perthii" are extremely slight and insufficient to warrant recognition of subspecies.

²⁷⁸ Distribution in WA is disjunct with isolates in the Cape Range region and on islands, nevertheless no subspecies are recognised (Johnstone and Storr 2004).

²⁷⁹ Formerly treated as a ssp. of *C. torquatus* (see Schodde and Mason 1999 and Kearns *et al.* 2014).

²⁸⁰ Ford (1979 and 1987) treats Port Essington in the Northern Territory as the type locality of *C. argenteus* and thus *C. a. latens* Ford (1979) is used for the Kimberley population.

²⁸¹ Although northern birds are much smaller than those from the far south, the distribution is virtually continuous, size changes clinally suggesting that the distribution of ssp. 'picatus' would be difficult to define and therefore not recognised.

²⁸² Sometimes placed in the genus *Gymnorhina*, but included here in *Cracticus*. Only two ssp. are recognised – *C. t. tibicen* of northern two-thirds of WA and *C. t. dorsalis* of south-western WA (Johnstone and Storr 2004).

²⁸³ The supposed differences between nominate *novaehollandiae* from Tasmania and mainland *melanops* require further confirmation as there appears to be considerable overlap in measurements.

<i>Coracina papuensis</i> (Gmelin, 1788)	White-bellied Cuckoo-shrike
<i>Coracina papuensis hypoleuca</i> (Gould, 1848)	White-breasted Cuckoo-shrike
<i>Coracina tenuirostris</i> (Jardine, 1831).....	Cicadabird (Common Cicadabird)
<i>Coracina tenuirostris melvillensis</i> Mathews, 1912	
²⁸⁴ <i>Lalage tricolor</i> (Swainson, 1825)	White-winged Triller
<i>Lalage leucomela</i> (Vigors and Horsfield, 1827)	Varied Triller
²⁸⁵ <i>Lalage leucomela rufiventris</i> (Gray, 1846)	
Family NEOSITTIDAE Sittellas	
<i>Daphoenositta chrysoptera</i> (Latham, 1802)	Varied Sittella
<i>Daphoenositta chrysoptera leucoptera</i> (Gould, 1840)	Varied Sittella (White-winged Sittella)
<i>Daphoenositta chrysoptera pileata</i> (Gould, 1838)	Varied Sittella (Black-capped Sittella)
Family OREOICIDAE Australo-Papuan bellbirds	
²⁸⁶ <i>Oreoica gutturalis</i> (Vigors and Horsfield, 1827)	Crested Bellbird
Family PACHYCEPHALIDAE Whistlers, shrike-thrushes and shriketits	
<i>Falcunculus frontatus</i> (Latham, 1802)	Crested Shriketit
²⁸⁷ <i>Falcunculus frontatus whitei</i> Campbell, 1910	Northern Shriketit
²⁸⁸ <i>Falcunculus frontatus leucogaster</i> Gould, 1838	
²⁸⁹ <i>Pachycephala inornata</i> Gould, 1841	Gilbert's Whistler
²⁹⁰ <i>Pachycephala occidentalis</i> Ramsay, 1878	Western Golden Whistler (Western Whistler)
<i>Pachycephala melanura</i> Gould, 1843	Mangrove Golden Whistler
²⁹¹ <i>Pachycephala melanura melanura</i> Gould, 1843	
<i>Pachycephala melanura spinicaudus</i> (Pucheran, 1853)	
²⁹² <i>Pachycephala rufiventris</i> (Latham, 1802)	Rufous Whistler
<i>Pachycephala rufiventris rufiventris</i> (Latham, 1802)	
²⁹³ <i>Pachycephala laniooides</i> Gould, 1840	White-breasted Whistler
<i>Colluricincla megarhyncha</i> (Quoy and Gaimard, 1830)	Little Shrike-thrush
²⁹⁴ <i>Colluricincla megarhyncha parvula</i> Gould, 1845	
<i>Colluricincla harmonica</i> (Latham, 1802)	Grey Shrike-thrush
<i>Colluricincla harmonica brunnea</i> Gould, 1841	
<i>Colluricincla harmonica kolichisi</i> Ford, 1987	
<i>Colluricincla harmonica rufiventris</i> Gould, 1841	
<i>Colluricincla woodwardi</i> Hartert, 1905	Sandstone Shrike-thrush
Family LANIIDAE Shrikes	
²⁹⁵ <i>Lanius tigrinus</i> Drapiez, 1828	Tiger Shrike

²⁸⁴ Forming a superspecies with *L. sueurii* of Indonesia.

²⁸⁵ The Kimberley population was described as a separate subspecies *L. l. macrura* by Schodde (1989) and although there is some overlap in length of wing and tail between Kimberley and Northern Territory populations, the two forms are separated by dry country at the head of Joseph Bonaparte Gulf and further work is needed to clarify its status.

²⁸⁶ Variation is clinal towards north and interior of WA with birds becoming smaller and paler. Mayr (1953) and Schodde and Mason (1999) recognised a pale northern and central Australian "*pallescens*" which is merely the northern end of this north-south cline. The change from typical "*gutturalis*" birds to "*pallescens*" is so gradual that it is impossible to draw a line between the two subspecies.

²⁸⁷ Sometimes treated as a full species.

²⁸⁸ Sometimes treated as a full species.

²⁸⁹ Contra Schodde and Mason (1999), distribution is accurately mapped in Johnstone and Storr (2004).

²⁹⁰ Formerly *P. pectoralis fuliginosa* Golden Whistler (Joseph *et al.* 2014).

²⁹¹ Nominate ssp. *melanura* ranges from Exmouth Gulf to Port Warrender and *spinicauda* from Cambridge Gulf to Mackay, Queensland (including islands in Torres Strait).

²⁹² The geographic variation of this continuously distributed species is clinal; the ssp. "*falcata*" of northern Australia is not recognised (Johnstone and Storr 2004).

²⁹³ Disjunct populations occur in WA, however, there is no concordance of characters used to separate these into ssp. "*laniooides*", "*carnarvonii*" and "*fretorum*". As such, no ssp. are recognised (Johnstone 1990).

²⁹⁴ Contra Schodde and Mason (1999), distribution is accurately mapped in Johnstone and Storr (2004).

²⁹⁵ Fremantle wharf 1997.

<i>Lanius cristatus</i> Linnaeus, 1758	Brown Shrike
296 <i>Lanius cristatus cristatus</i> Linnaeus, 1758	
297 <i>Lanius cristatus lucionensis</i> Linnaeus, 1766	
Family ORIOLIDAE Orioles and figbirds	
<i>Sphecootheres vieilloti</i> Vigors and Horsfield, 1827	Australasian Figbird
<i>Sphecootheres vieilloti ashbyi</i> Mathews, 1912	Yellow Figbird
<i>Oriolus sagittatus</i> (Latham, 1802)	Olive-backed Oriole
<i>Oriolus sagittatus affinis</i> Gould, 1848	
<i>Oriolus flavocinctus</i> (King, 1826)	Yellow Oriole (Green Oriole)
<i>Oriolus flavocinctus flavocinctus</i> (King, 1826)	
Family DICRURIDAE Drongos	
298 <i>Dicrurus bracteatus</i> Gould, 1843	Spangled Drongo
<i>Dicrurus bracteatus bracteatus</i> Gould, 1843	
Family RHIPIDURIDAE Typical fantails	
299 <i>Rhipidura leucophrys</i> (Latham, 1802)	Willie Wagtail
<i>Rhipidura leucophrys leucophrys</i> (Latham, 1802)	
<i>Rhipidura rufiventris</i> (Vieillot, 1818)	Northern Fantail
<i>Rhipidura rufiventris isura</i> Gould, 1841	
300 <i>Rhipidura albiscapa</i> Gould, 1840	Grey Fantail
301 <i>Rhipidura albiscapa alisteri</i> Mathews, 1911	
<i>Rhipidura albiscapa preissi</i> Cabanis, 1850	
<i>Rhipidura albiscapa albicauda</i> North, 1895	White-tailed Fantail
302 <i>Rhipidura phasiana</i> De Vis, 1885	Mangrove Grey Fantail (Mangrove Fantail)
303 <i>Rhipidura dryas</i> Gould, 1843	Wood Fantail (Arafura Fantail, Rufous Fantail)
<i>Rhipidura dryas dryas</i> Gould, 1843	
<i>Rhipidura dryas semicollaris</i> Müller, 1843	
Family MONARCHIDAE Monarch flycatchers	
<i>Monarcha cinerascens</i> (Temminck, 1827)	Island Monarch
<i>Monarcha melanopsis</i> (Vieillot, 1818)	Black-faced Monarch
304 <i>Grallina cyanoleuca</i> (Latham, 1802)	Magpie-lark
<i>Myiagra rubecula</i> (Latham, 1802)	Leaden Flycatcher
<i>Myiagra rubecula concinna</i> Gould, 1848	
<i>Myiagra ruficollis</i> (Vieillot, 1818)	Broad-billed Flycatcher
<i>Myiagra ruficollis mimikae</i> Ogilvie-Grant, 1911	
<i>Myiagra cyanoleuca</i> (Vieillot, 1818)	Satin Flycatcher
<i>Myiagra alecto</i> (Temminck, 1827)	Shining Flycatcher
<i>Myiagra alecto melvillensis</i> (Mathews, 1912)	
<i>Myiagra inquieta</i> (Latham, 1802)	Restless Flycatcher
<i>Myiagra inquieta inquieta</i> (Latham, 1802)	
305 <i>Myiagra inquieta nana</i> (Gould, 1870)	Paperbark Flycatcher

296 Ashmore Reef. Photographs only.

297 Ashmore Reef. Photographs only.

298 The recognition of northern ssp. "baileyi" on the grounds of its larger size is perplexing as measurements show it is smaller than nominate *bracteatus*, and there appears to be considerable overlap in size between northern and southern birds.299 Only the nominate form *R. l. leucophrys* is recognised for mainland Australia (Mees 1961).300 Previously treated as ssp. of the New Zealand Fantail *R. fuliginosa*. Distribution of ssp. in WA mapped in Johnstone and Storr 2004.

301 Winter visitor.

302 Forming a superspecies with the Grey Fantail and possibly the White-tailed Fantail.

303 Most recently treated as a ssp. of the Rufous Fantail *R. rufifrons*. However, *rufifrons* and *dryas* do not intergrade in north Queensland and are treated here as allospecies in a view for their marked differences in habitat and morphology. Both *R. dryas* and *R. rufifrons* forming a superspecies with several east Indonesian species.

304 Distribution is continuous both north-south and east-west. No ssp. recognised in WA.

305 Sometimes treated as a full species, but retained here as ssp. (refer Andersen *et al.* 2015)

Family CORVIDAE Crows, ravens, rooks, jays and allies

- ³⁰⁶*Pica pica* (Linnaeus, 1758) Eurasian Magpie
Corvus splendens Vieillot, 1817 House Crow
Corvus orru Bonaparte, 1850 Torresian Crow
 Corvus orru cecilae Mathews, 1912 Western Crow
Corvus bennetti North, 1901 Little Crow
Corvus coronoides Vigors and Horsfield, 1827 Australian Raven
 Corvus coronoides perplexus Mathews, 1912

Family PETROICIDAE Australasian robins, scrub-robins and flycatchers

- ³⁰⁷*Poecilodryas cerviniventris* (Gould, 1858) Buff-sided Robin
³⁰⁸*Eopsaltria pulverulenta* (Bonaparte, 1850) Mangrove Robin
Eopsaltria australis (Shaw, 1790) Yellow Robin
³⁰⁹*Eopsaltria australis griseogularis* Gould, 1838 Western Yellow Robin
Eopsaltria georgiana (Quoy and Gaimard, 1830) White-breasted Robin
³¹⁰*Melanodryas cucullata* (Latham, 1802) Hooded Robin
Microeca flavigaster Gould, 1843 Lemon-breasted Flycatcher (Lemon-bellied Flyrobin)
 Microeca flavigaster flavigaster Gould, 1843
 Microeca flavigaster tormenti Mathews, 1916 Kimberley Flycatcher
³¹¹*Microeca fascinans* (Latham, 1802) Jacky Winter
 Microeca fascinans fascinans (Latham, 1802)
 Microeca fascinans assimilis Gould, 1841
³¹²*Petroica boodang* (Lesson, 1838) Scarlet Robin
 Petroica boodang campbelli Sharpe, 1898 Campbell's Scarlet Robin
Petroica goodenovii (Vigors and Horsfield, 1827) Red-capped Robin
Drymodes brunneopygia Gould, 1841 Southern Scrub-robin
Family ALAUDIDAE Larks
³¹³*Mirafra javanica* Horsfield, 1821 Horsfield's Bushlark
 Mirafra javanica horsfieldii Gould, 1847
Family PYCNONOTIDAE Bulbs
Pycnonotus jocosus (Linnaeus, 1758) Red-whiskered Bulbul
³¹⁴*Pycnonotus jocosus jocosus* (Linnaeus, 1758)
- Family HIRUNDINIDAE Swallows and martins**
Cheramoeca leucosterna (Gould, 1841) White-backed Swallow

³⁰⁶ No specimens examined, subspecies unknown. Probably ship-assisted.

³⁰⁷ Sometimes treated as a ssp. of *P. superciliosa*.

³⁰⁸ Sometimes placed in the genus *Peneoenanthe*. No subspecies recognised (Mees 1982; Ford 1983 and Johnstone 1990).

³⁰⁹ Sometimes treated as a full species *E. griseogularis*. Ford (1979) treated the western and eastern forms of the Yellow Robin as conspecific. The ssp. "rosinae" is not recognised as variation is clinal and it is impossible to draw a line between the two races.

³¹⁰ No ssp. are recognised in Western Australia. Subspecies "picata" in the north i.e. Kimberley and "westralensis" in the south are merely the ends of a north-south cline in increasing size (Johnstone and Storr 2004). The Melville Island population (*M. c. melvillensis*) may be taxonomically distinguishable in which case the mainland population would be treated as the nominate form.

³¹¹ The north-western Australian population in Kimberley, northern Northern Territory and western Queensland is sometimes treated as separate subspecies "pallida".

³¹² Australian populations now separated from Pacific populations *P. multicolor* as a separate species *P. boodang* (Shodde and Mason 1999).

³¹³ In Western Australia intergradation of colour morphs is complete between one local variant and another (birds varying locally according to colour of soil substrate). All Australian populations are best treated as components of a single variable ssp. (Johnstone and Storr 2004).

³¹⁴ Safety Bay 1993.

<i>Hirundo rustica</i> Linnaeus, 1758	Barn Swallow
³¹⁵ <i>Hirundo rustica gutturalis</i> Scopoli, 1786	
³¹⁶ <i>Hirundo rustica tytleri</i> Jerdon, 1864	
³¹⁷ <i>Hirundo tahitica</i> Gmelin, 1789	Pacific Swallow
<i>Hirundo neoxena</i> Gould, 1842	Welcome Swallow
³¹⁸ <i>Hirundo neoxena [carteri]</i> (Mathews, 1912)	
³¹⁹ <i>Cecropis daurica</i> (Laxmann, 1769)	Red-rumped Swallow
³²⁰ <i>Cecropis daurica japonica</i> (Temminck and Schlegel, 1845)	
³²¹ <i>Petrochelidon ariel</i> (Gould, 1842)	Fairy Martin
³²² <i>Petrochelidon nigricans</i> (Vieillot, 1817)	Tree Martin
<i>Petrochelidon nigricans nigricans</i> (Vieillot, 1817)	
Family CETTIIDAE Cettiid warblers	
³²³ <i>Urosphena squameiceps</i> (Swinhoe, 1863)	Asian Stubtail
Family PHYLLOSCOPIDAE Leaf warblers	
³²⁴ <i>Phylloscopus trochilus</i> (Linnaeus, 1758)	Willow Warbler
³²⁵ <i>Phylloscopus inornatus</i> (Blyth, 1842)	Yellow-browed Warbler
³²⁶ <i>Phylloscopus borealis</i> (Blasius, 1858)	Arctic Warbler
<i>Phylloscopus borealis borealis</i> (Blasius, 1858)	
³²⁷ <i>Phylloscopus examinandus</i> Stresemann, 1913	Kamchatka Leaf Warbler
³²⁸ <i>Phylloscopus coronatus</i> (Temminck and Schlegel, 1847)	Eastern Crowned Warbler
Family ACROCEPHALIDAE Reed warblers	
³²⁹ <i>Acrocephalus orientalis</i> (Temminck and Schlegel, 1847)	Oriental Reed Warbler
<i>Acrocephalus australis</i> (Gould, 1838)	Australian Reed Warbler
<i>Acrocephalus australis gouldi</i> Dubois, 1901	
Family LOCUSTELLIDAE Grass warblers and grassbirds	
³³⁰ <i>Locustella ochotensis</i> (Middendorff, 1853)	Middendorff's Grasshopper Warbler
³³¹ <i>Locustella certhiola</i> (Pallas, 1811)	Pallas's Grasshopper Warbler
<i>Locustella fasciolata</i> (Gray, 1861)	Gray's Grasshopper Warbler (Gray's Warbler)
³³² <i>Megalurus mathewsi</i> (Iredale, 1911)	Rufous Songlark
³³³ <i>Megalurus cruralis</i> Vigors and Horsfield, 1827	Brown Songlark

³¹⁵ Non-breeding migrant.³¹⁶ Non-breeding migrant.³¹⁷ No specimens available, but ssp. almost certainly *javanica*.³¹⁸ Tentatively recognised on measurements, but range is continuous between eastern and western populations in southern Australia and variation possibly clinal.³¹⁹ Non-breeding migrant. Sometimes placed in the genus *Hirundo*.³²⁰ Sometimes placed in the genus *Hirundo*.³²¹ Sometimes placed in the genus *Hirundo*.³²² Sometimes placed in the genus *Hirundo*. Australian and Timor populations treated under the nominate form contra Gill and Donsker (2013).³²³ Ashmore Reef 2012. Photographs only.³²⁴ Ashmore Reef 2013. No specimens, but most likely ssp. *yakutensis* of eastern Siberia.³²⁵ Ashmore Reef 2011.³²⁶ Non-breeding migrant.³²⁷ Ashmore Reef 2012.³²⁸ Ashmore Reef 2012. Photographs only.³²⁹ Ashmore Reef 2011.³³⁰ No specimens and ssp. unresolved.³³¹ Ashmore Reef 2011.³³² Sometimes placed in the genus *Cincloramphus*.³³³ Sometimes placed in the genus *Cincloramphus*.

<i>Megalurus gramineus</i> (Gould, 1845)	Little Grassbird
³³⁴ <i>Megalurus gramineus gramineus</i> (Gould, 1845)	
<i>Megalurus timoriensis</i> Wallace, 1864	Tawny Grassbird
<i>Megalurus timoriensis alisteri</i> Mathews, 1912	
³³⁵ <i>Eremiornis carteri</i> North, 1900	Spinifexbird
Family CISTICOLIDAE Cisticolas and allies	
<i>Cisticola juncidis</i> (Rafinesque, 1810)	Zitting Cisticola
³³⁶ <i>Cisticola juncidis fuscicapilla</i> Wallace, 1864	
<i>Cisticola exilis</i> (Vigors and Horsfield, 1827)	Golden-headed Cisticola
³³⁷ <i>Cisticola exilis lineocapilla</i> Gould, 1847	
Family ZOSTEROPIDAE White-eyes	
³³⁸ <i>Zosterops citrinella</i> Bonaparte, 1850	Ashy-bellied White-eye
³³⁹ <i>Zosterops luteus</i> Gould, 1843	Yellow White-eye (Canary White-eye)
<i>Zosterops lateralis</i> (Latham, 1802)	Grey-breasted White-eye (Silvereye)
³⁴⁰ <i>Zosterops lateralis chloronotus</i> Gould, 1841	
Family STURNIDAE Starlings	
<i>Acridotheres tristis</i> (Linnaeus, 1766)	Common Myna
³⁴¹ <i>Acridotheres tristis [tristis]</i> (Linnaeus, 1766)	
³⁴² <i>Agropsar philippensis</i> (Forster, 1781)	Chestnut-cheeked Starling
<i>Pastor roseus</i> (Linnaeus, 1758)	Rosy Starling (Rose-coloured Starling)
<i>Sturnus vulgaris</i> Linnaeus, 1758	Common Starling
<i>Sturnus vulgaris vulgaris</i> Linnaeus, 1758	
Family TURDIDAE Thrushes	
³⁴³ <i>Geokichla sibirica</i> (Pallas, 1776)	Siberian Thrush
<i>Turdus merula</i> Linnaeus, 1758	Common Blackbird (Eurasian Blackbird)
<i>Turdus merula merula</i> Linnaeus, 1758	
³⁴⁴ <i>Turdus obscurus</i> Gmelin, 1789	Eyebrowed Thrush
Family MUSCICAPIDAE Old World flycatchers	
³⁴⁵ <i>Muscicapa griseisticta</i> (Swinhoe, 1861)	Grey-streaked Flycatcher
<i>Muscicapa sibirica</i> Gmelin, 1789	Sooty Flycatcher (Dark-sided Flycatcher)
³⁴⁶ <i>Muscicapa sibirica [sibirica]</i> Gmelin, 1789	
<i>Muscicapa latirostris</i> Raffles, 1822	Asian Brown Flycatcher
³⁴⁷ <i>Muscicapa latirostris [latirostris]</i> Raffles, 1822	
<i>Cyanoptila cyanomelana</i> (Temminck, 1829)	Blue and White Flycatcher
<i>Cyanoptila cyanomelana cyanomelana</i> (Temminck, 1829)	
³⁴⁸ <i>Larvivora cyane</i> (Pallas, 1776)	Siberian Blue Robin

³³⁴ Numerous disjunct populations in WA, ssp. "thomasi" of south Western Australia is not recognised (Johnstone and Storr 2004).

³³⁵ Sometimes placed in the genus *Megalurus*.

³³⁶ The few Kimberley specimens match *fuscicapilla* of the eastern Lesser Sundas.

³³⁷ Only one ssp. *lineocapilla* is recognised for Western Australia and the Northern Territory.

³³⁸ Ashmore Reef. No specimens, but ssp. most likely to be nominate.

³³⁹ Distribution in WA is disjunct, but variation largely clinal. No subspecies are recognised (Johnstone 1990 and Ford 1983).

³⁴⁰ Previously known as ssp. *gouldii*, but *chloronotus* has priority.

³⁴¹ Possibly ship-assisted.

³⁴² Ashmore Reef 2011.

³⁴³ Esperance 2011 and Ashmore Reef 2012.

³⁴⁴ Port Hedland 2014.

³⁴⁵ Ashmore Reef 2010.

³⁴⁶ No specimens examined, but most likely ssp. the nominate form.

³⁴⁷ Ashmore Reef 2005. No specimens examined, but most likely ssp. the nominate form.

³⁴⁸ Ashmore Reef 2012. Photographs only.

<i>Ficedula narcissina</i> (Temminck, 1836)	Narcissus Flycatcher
<i>Ficedula narcissina narcissina</i> (Temminck, 1836)	
<i>Monticola solitarius</i> (Linnaeus, 1758)	Blue Rock Thrush
Family DICAEIDAE Mistletoebird	
<i>Dicaeum hirundinaceum</i> (Shaw, 1792)	Mistletoebird
<i>Dicaeum hirundinaceum hirundinaceum</i> (Shaw, 1792)	
Family PASSERIDAE True (Old World) sparrows	
^{349*} <i>Passer domesticus</i> (Linnaeus, 1758)	House Sparrow
^{350*} <i>Passer domesticus domesticus</i> (Linnaeus, 1758)	
* <i>Passer montanus</i> (Linnaeus, 1758)	Eurasian Tree Sparrow
Family ESTRILDIDAE Estrildid finches	
<i>Emblema pictum</i> Gould, 1842	Painted Finch
<i>Stagonopleura oculata</i> (Quoy and Gaimard, 1830)	Red-eared Firetail
^{351*} <i>Neochmia temporalis</i> (Latham, 1802)	Red-browed Finch
* <i>Neochmia temporalis temporalis</i> (Latham, 1802)	
<i>Neochmia phaeton</i> (Hombron and Jacquinot, 1841)	Crimson Finch
<i>Neochmia phaeton phaeton</i> (Hombron and Jacquinot, 1841)	
³⁵² <i>Neochmia ruficauda</i> (Gould, 1837)	Star Finch
<i>Neochmia ruficauda clarescens</i> (Hartert, 1899)	
<i>Poephila personata</i> Gould, 1842	Masked Finch
<i>Poephila personata personata</i> Gould, 1842	
³⁵³ <i>Poephila acuticauda</i> (Gould, 1840)	Long-tailed Finch
<i>Taeniopygia guttata</i> (Vieillot, 1817)	Zebra Finch
<i>Taeniopygia guttata castanotis</i> (Gould, 1837)	
<i>Taeniopygia bichenovii</i> (Vigors and Horsfield, 1827)	Double-barred Finch
<i>Taeniopygia bichenovii annulosa</i> (Gould, 1840)	
<i>Erythrura gouldiae</i> (Gould, 1844)	Gouldian Finch
<i>Lonchura punctulata</i> (Linnaeus, 1758)	Scaly-breasted Munia
³⁵⁴ <i>Lonchura pallida</i> (Wallace, 1864)	Pale-headed Munia
<i>Lonchura flaviprymna</i> (Gould, 1845)	Yellow-rumped Mannikin
<i>Lonchura castaneothorax</i> (Gould, 1837)	Chestnut-breasted Mannikin
<i>Lonchura castaneothorax castaneothorax</i> (Gould, 1837)	
<i>Heteromunia pectoralis</i> (Gould, 1841)	Pictorella Mannikin
Family MOTACILLIDAE Wagtails and pipits	
³⁵⁵ <i>Motacilla flava</i> Linnaeus, 1758	Yellow Wagtail
³⁵⁶ <i>Motacilla flava simillima</i> Hartert, 1905	
³⁵⁷ <i>Motacilla flava macronyx</i> (Stresemann, 1920)	
³⁵⁸ <i>Motacilla flava plexa</i> (Thayer and Bangs, 1914)	
³⁵⁹ <i>Motacilla flava taivana</i> (Swinhoe, 1863)	

³⁴⁹ Mainly ship-assisted vagrants.³⁵⁰ Mainly ship-assisted vagrants.³⁵¹ Established in northern Darling Range in 1950-60.³⁵² Western Australian, Northern Territory and Cape York populations treated as a single ssp. *clarescens* (Ford 1986).³⁵³ Subspecies "hecki" of Northern Territory is doubtfully distinct (Johnstone and Storr 2004).³⁵⁴ Ashmore Reef 2000.³⁵⁵ Sometimes Yellow Wagtail treated as two taxa - Eastern and Western Yellow Wagtail, but retained here as a single species *M. flava*.³⁵⁶ Non-breeding migrant. *M. f. simillima* herein considered distinct from *M. f. tschutschensis*.³⁵⁷ Non-breeding migrant.³⁵⁸ Non-breeding migrant.³⁵⁹ Non-breeding migrant.

³⁶⁰ <i>Motacilla cinerea</i> Tunstall, 1771	Grey Wagtail
³⁶¹ <i>Motacilla cinerea cinerea</i> Tunstall, 1771	
<i>Motacilla alba</i> Linnaeus, 1758	White Wagtail
<i>Motacilla alba ocularis</i> Swinhoe, 1860	
<i>Motacilla alba lugens</i> Gloger, 1829	
<i>Motacilla alba leucopsis</i> Gould, 1838	
³⁶² <i>Anthus australis</i> Vieillot, 1818	Australian Pipit
<i>Anthus australis australis</i> Vieillot, 1818	
³⁶³ <i>Anthus gustavi</i> Swinhoe, 1863	Pechora Pipit
<i>Anthus cervinus</i> (Pallas, 1811)	Red-throated Pipit
Family FRINGILLIDAE True finches	
^{364*} <i>Carduelis carduelis</i> (Linnaeus, 1758)	European Goldfinch
*i [†] <i>Carduelis carduelis [britannica]</i> (Hartert, 1903)	

³⁶⁰ Non-breeding migrant.

³⁶¹ *M. c. robusta* synonymised with *M. c. cinerea*.

³⁶² Previously included in the superspecies *Anthus novaeseelandiae* (Gmelin 1789). All Australian mainland populations treated as *A. a. australis*.

³⁶³ No specimens examined, but the nominate form the most likely ssp.

³⁶⁴ Known only from aviary escapees. Some recent records of large flocks on Swan Coastal Plain indicate that at times escapees can establish colonies.

LIST OF ORDERS AND FAMILIES

CASUARIIFORMES.....	2
DROMAIIDAE Emu.....	2
ANSERIFORMES	2
ANSERANATIDAE Magpie Goose.....	2
ANATIDAE Ducks, geese and swans	2
GALLIFORMES	2
MEGAPODIIDAE Megapodes	2
PHASIANIDAE Pheasants.....	2
SPHENISCIFORMES	3
SPHENISCIDAE Penguins.....	3
PROCELLARIIFORMES.....	3
DIOMEDEIDAE Albatrosses	3
PROCELLARIIDAE Petrels and shearwaters	4
HYDROBATIDAE Storm petrels.....	5
PELECANOIDIDAE Diving petrels.....	5
PODICIPEDIFORMES.....	5
PODICIPEDIDAE Grebes	5
PHAETHONTIFORMES	5
PHAETHONTIDAE Tropicbirds.....	5
CICONIIFORMES	6
CICONIIDAE Storks	6
PELECANIFORMES	6
THRESKIORNITHIDAE Ibises and spoonbills	6
ARDEIDAE Herons and bitterns	6
PELECANIDAE Pelicans	7
SULIFORMES	7
FREGATIDAE Frigatebirds	7
SULIDAE Gannets and boobies.....	7
PHALACROCORACIDAE Cormorants	7
ANHINGIDAE Darters.....	7
ACCIPITRIFORMES	7
PANDIONIDAE Osprey.....	7
ACCIPITRIDAE Hawks, eagles, kites and harriers.....	8
OTIDIIFORMES	8
OTIDIDAE Bustards	8
GRUIFORMES	8
RALLIDAE Rails, crakes, waterhens and coots	8
GRUIDAE Cranes.....	9
CHARADRIIFORMES	9
TURNICIDAE Button-quails	9
BURHINIDAE Stone-curlews.....	9
HAEMATOPODIDAE Oystercatchers.....	9
RECURVIROSTRIDAE Stilts and avocets	9
CHARADRIIDAE Plovers, dotterels and lapwings	10
ROSTRATULIDAE Painted snipes	10
JACANIDAE Jacanas.....	10
SCOLOPACIDAE Sandpipers and snipes	11
GLAREOLIDAE Pratincoles	12
LARIDAE Gulls and terns	12
STERCORARIIDAE Skuas.....	14
COLUMBIFORMES	14
COLUMBIDAE Pigeons and doves.....	14
CUCULIFORMES.....	15
CUCULIDAE Cuckoos	15
STRIGIFORMES	15
TYTONIDAE Barn owls.....	15
STRIGIDAE Typical owls.....	16

CAPRIMULGIFORMES	16
PODARGIDAE Frogmouths.....	16
CAPRIMULGIDAE Nightjars.....	16
APODIFORMES	16
AEGOTHELIDAE Owlet-nightjars	16
APODIDAE Swifts.....	16
CORACIIFORMES	16
CORACIIDAE Rollers.....	16
ALCEDINIDAE Kingfishers.....	17
MEROPIDAE Bee-eaters	17
BUCEROTIFORMES.....	17
UPUPIDAE Hoopoe	17
FALCONIFORMES	17
FALCONIDAE Falcons and kestrels.....	17
PSITTACIFORMES	17
CACATUIDAE Cockatoos	17
PSITTACIDAE Parrots.....	18
PASSERIFORMES	19
PITTIDAE Pittas.....	19
ATRICHORNITHIDAE Scrub-birds	19
PTILONORHYNCHIDAE Bowerbirds	19
CLIMacteridae Treecreepers.....	20
MALURIDAE Grasswrens, fairy-wrens and emu-wrens	20
MELIPHAGIDAE Honeyeaters and chats.....	20
DASYORNITHIDAE Bristlebirds	22
PARDALOTIDAE Pardalotes.....	22
ACANTHIZIDAE Gerygones, thornbills, scrubwrens and whitefaces.....	22
POMATOSTOMIDAE Australasian babblers.....	23
PSOPHODIDAE Quail-thrushes, whipbirds and wedgebills.....	24
ARTAMIDAE Woodswallows.....	24
CRACTICIDAE Butcherbirds, Australian magpies and currawongs	24
CAMPEPHAGIDAE Cuckoo-shrikes and trillers	24
NEOSITTIDAE Sittellas	25
OREOICIDAE Australo-Papuan bellbirds	25
PACHYCEPHALIDAE Whistlers, shrike-thrushes and shriketits	25
LANIIDAE Shrikes.....	25
ORIOLIDAE Orioles and figbirds	26
DICRURIDAE Drongos.....	26
RHIPIDURIDAE Typical fantails	26
MONARCHIDAE Monarch flycatchers.....	26
CORVIDAE Crows, ravens, rooks, jays and allies	27
PETROICIDAE Australasian robins, scrub-robins and flycatchers.....	27
ALAUDIDAE Larks.....	27
PYCNONOTIDAE Bulbuls	27
HIRUNDINIDAE Swallows and martins	27
CETTIIDAE Cettiid warblers	28
PHYLLOSCOPIDAE Leaf-Warblers.....	28
ACROCEPHALIDAE Reed warblers	28
LOCUSTELLIDAE Grass warblers and grassbirds	28
CISTICOLIDAE Cisticolas and allies	29
ZOSTEROPIDAE White-eyes	29
STURNIDAE Starlings	29
TURDIDAE Thrushes	29
MUSCICAPIDAE Old World flycatchers	29
DICAЕIDAE Mistletoebird	30
PASSERIDAE True (Old World) sparrows.....	30
ESTRILDIDAE Estrildid finches	30
MOTACILLIDAE Wagtails and pipits	30
FRINGILLIDAE True finches.....	31

INDEX OF ENGLISH FAMILY NAMES

- | | | |
|----------------------------|-------------------------|--------------------------|
| Albatrosses, 3 | Gannets, 7 | Pratincoles, 12 |
| Australasian babblers, 23 | Geese, 2 | Quails, 2 |
| Avocets, 9 | Gerygones, 22 | Quail-thrushes, 24 |
| Babblers, 23 | Goose, Magpie, 2 | Rails, 8 |
| Barn owls, 15 | Grass warblers, 28 | Ravens, 27 |
| Bee-eaters, 17 | Grassbirds, 28 | Reed warblers, 28 |
| Bellbirds, 25 | Grasswrens, 20 | Robins, Australasian, 27 |
| Bitterns, 6 | Grebes, 5 | Rollers, 16 |
| Boobies, 7 | Gulls, 12 | Rooks, 27 |
| Bowerbirds, 19 | Harriers, 8 | Sandpipers, 11 |
| Bristlebirds, 22 | Hawks, 8 | Scrub-birds, 19 |
| Bulbuls, 27 | Herons, 6 | Scrub-robbins, 27 |
| Bustards, 8 | Honeyeaters, 20 | Scrubwrens, 22 |
| Butcherbirds, 24 | Hoopoe, 17 | Shearwaters, 4 |
| Button-quails, 9 | Ibises, 6 | Shrikes, 25 |
| Cettiid warblers, 28 | Jacanas, 10 | Shrike-thrushes, 25 |
| Chats, 20 | Jays, 27 | Shriketits, 25 |
| Cisticolas, 29 | Kestrels, 17 | Sittellas, 25 |
| Cockatoos, 17 | Kingfishers, 17 | Skuas, 14 |
| Coots, 8 | Kites, 8 | Snipes, 11 |
| Cormorants, 7 | Lapwings, 10 | Sparrows, Old World, 30 |
| Crakes, 8 | Larks, 27 | Spoonbills, 6 |
| Cranes, 9 | Leaf-Warbblers, 28 | Starlings, 29 |
| Crows, 27 | Magpie Goose, 2 | Stilts, 9 |
| Cuckoos, 15 | Magpies, Australian, 24 | Stone-curlews, 9 |
| Cuckoo-shrikes, 24 | Martins, 27 | Storks, 6 |
| Currawongs, 24 | Megapodes, 2 | Storm petrels, 5 |
| Darters, 7 | Mistletoebird, 30 | Swallows, 27 |
| Diving petrels, 5 | Monarch flycatchers, 26 | Swans, 2 |
| Dotterels, 10 | Nightjars, 16 | Swifts, 16 |
| Doves, 14 | Orioles, 26 | Terns, 12 |
| Drongos, 26 | Osprey, 7 | Thornbills, 22 |
| Ducks, 2 | Owl-nightjars, 16 | Thrushes, 29 |
| Eagles, 8 | Owls, 16 | Treecreepers, 20 |
| Emu, 2 | Oystercatchers, 9 | Trillers, 24 |
| Emu-wrens, 20 | Painted snipes, 10 | Tropicbirds, 5 |
| Estrildid finches, 30 | Pardalotes, 22 | Wagtails, 30 |
| Fairy-wrens, 20 | Parrots, 18 | Warblers |
| Falcons, 17 | Pelicans, 7 | Cettiid, 28 |
| Fantails, 26 | Penguins, 3 | Grass, 28 |
| Figbirds, 26 | Petrels, 4 | Reed, 28 |
| Finches, 31 | Diving, 5 | Waterhens, 8 |
| Estrildid, 30 | Storm, 5 | Wedgebills, 24 |
| True, 31 | Pheasants, 2 | Whipbirds, 24 |
| Flycatchers, 27 | Pigeons, 14 | Whistlers, 25 |
| Flycatchers, Old World, 29 | Pipits, 30 | White-eyes, 29 |
| Frigatebirds, 7 | Pittas, 19 | Whitefaces, 22 |
| Frogmouths, 16 | Plovers, 10 | Woodswallows, 24 |