

VARIICHTHYS, A REPLACEMENT NAME FOR THE TERAPONIID FISH GENUS VARIA AND FIRST RECORD OF *V. LACUSTRIS* FROM AUSTRALIA

Gerald R. Allen*

Vari (1978) revised the family Terapontidae (spelled Theraponidae or Teraponidae by some authors). These perch-like fishes inhabit the Indo-West Pacific region with 33 of 45 known species being confined to fresh waters of Australia and New Guinea. Vari provisionally assigned *Therapon jamaerensis* Mees (1971) from southern New Guinea to the genus "*Terapon*". He stated that it belonged to a new genus, but failed to describe it because of its uncertain relationships. Although he did not examine specimens of *Therapon lacustris* Mees and Kailola (1977), Vari mentioned in the addendum of his 1978 work that it was possibly the sister species of *T. jamaerensis*.

Allen (1991) eventually placed the two species in a new genus *Varia*. However, Carl J. Ferraris, Jr. of the Department of Ichthyology at the American Museum of Natural History (New York) recently informed the author that this generic name is preoccupied. Walker (1867) described *Varia* as a new genus of moths (Lepidoptera). Therefore, its use for terapontid fishes is invalid. The replacement name *Variichthys* is proposed herein.

A brief diagnosis of this genus is presented below and the first record of its presence in Australia is noted.

Variichthys, nom. nov.

Varia Allen, 1991: 128 (preoccupied by *Varia* Walker 1867, a genus of Lepidoptera).

Type species

Therapon jamaerensis Mees, 1971: 214.

Diagnosis

A genus of terapontid fishes characterised by the following combination of features: posttemporal bone covered externally with skin and scales, not expanded posteriorly, and not having a serrate posterior margin; two fin spines arising from the first proximal dorsal pterygiophore; lateral-line scale count 55-66; each jaw with a band of fixed (non-depressible), conical teeth, the outer row somewhat enlarged; dorsal rays XII to XIV, 10 or 11; anal rays III, 8 or 9; pectoral rays 14 or 15; gill rakers on first arch 5 or 6+1+10 or 11; vertebrae 11+14.

Remarks

The genus contains two species: *V. jamaerensis* (Mees), **comb. nov.** known only from Lake

* Department of Aquatic Zoology, Western Australian Museum, Francis Street, Perth, Western Australia 6000

Jamur (sometimes spelled Jamoer or Yamur) in Irian Jaya (approximately 3°39'S, 134°58'E) and *V. lacustris* (Mees and Kailola), **comb. nov.** formerly thought to be confined to the Fly River Delta region of Papua New Guinea (see comments below).

FIRST RECORD OF *V. LACUSTRIS* FROM AUSTRALIA

A specimen of *V. lacustris* was recently presented to the author by Gordon Stables of Cairns, Queensland. A number of individuals were caught (all released except one) with a seine net by Mr. Stables in Twelve-Mile Lagoon, Mitchell River Drainage, Cape York Peninsula, Queensland (approximately 16°24'S, 143°09'E) in September 1992. This specimen is now deposited in the collection of the Western Australian Museum (reg. no. WAM P.30652-001). The species was previously known only from the Fly River Delta of Papua New Guinea, including the Balimo area, Morehead River, Lake Murray, and the lower Fly River (Allen, 1991). It is apparently common in Twelve Mile Lagoon in the Mitchell River system and future collecting will probably reveal a broader distribution on Cape York Peninsula.

Counts and measurements for the specimen (164.2 mm SL) from Twelve-Mile Lagoon are as follows: dorsal rays XIII,10; anal rays III,8; pectoral rays 14; tubed lateral-line scales 63; scales in horizontal row just above lateral line 73; gill rakers on first arch 6+1+10; maximum body depth 66 mm; maximum body width 28.5 mm; head length 46.0 mm; snout length 13.2 mm; upper jaw length 13.8 mm; eye diameter 11.1 mm; predorsal length 64.6 mm; prepelvic length 62.3 mm; preanal length 111.1 mm; least depth of caudal peduncle 20.8; length of caudal peduncle 32.3 mm; length of dorsal fin base 92.4 mm; longest (fourth) dorsal spine 30.2 mm; longest (first) soft dorsal ray 23.8 mm; length of anal fin base 29.5 mm; longest (second) anal spine 26.9 mm SL; longest (first) soft anal ray 26.5 mm; length of caudal fin 43.0 mm.

ACKNOWLEDGEMENTS

I thank Gordon Stables for donating the specimen of *V. lacustris* to the WAM fish collection. Special thanks are also due Carl Ferraris, Jr. for his discovery of the preoccupied status of *Varia*.

REFERENCES

- Allen, G.R. (1991). *Field guide to the freshwater fishes of New Guinea*. Christensen Research Institute (Papua New Guinea) Publ. No. 9, 268 pp.
- Mees, G.F. (1971). Revisional notes on some species of the genus *Therapon* (Pisces, Theraponidae). *Zool. Med.* **45**: 197-224.
- Mees, G.F. and Kailola, P.J. (1977). The freshwater Therapontidae of New Guinea. *Zool. Verhand.* **153**: 1-89.
- Walker, F. (1867). Characters of some undescribed Heterocerous Lepidoptera. *J. Linn. Soc.* **9**: 181-199.
- Vari, R.P. (1978). The terapon perches (Percoidei, Teraponidae), a cladistic analysis and taxonomic revision. *Bull. Amer. Mus. Nat. Hist.* **159**(5): 175-340.

Guide to Authors

Subject Matter:

Reviews, observations and results of research into all branches of natural science and human studies will be considered for publication. However, emphasis is placed on studies pertaining to Western Australia. Full length papers should not normally exceed 30 typed pages. Short communications should not normally exceed three typed pages and this category of paper is intended to accommodate observations, results or new records of *significance*, that otherwise might not get into the literature, or for which there is a particular urgency for publication. All material must be original and not have been published elsewhere.

Presentation:

Authors are advised to follow the layout and style in the most recent issue of the *Rec. West. Aust. Mus.* including headings, tables, illustrations and references.

The title should be concise, informative and contain key words necessary for retrieval by modern searching techniques. Names of new taxa must not be included. An abridged title (not exceeding 50 letter spaces) should be included for use as a running head.

An abstract must be given in full length papers but not short communications, summarizing the scope of the work and principal findings. It should normally not exceed 2% of the paper and should be suitable for reprinting in reference periodicals. Contrary to Recommendation 23 of the International Code of Zoological Nomenclature it may include names of new taxa.

Footnotes are to be avoided, except in papers dealing with historical subjects.

The International System of units should be used.

Numbers should be spelled out from one to nine in descriptive text; figures used for 10 or more. For associated groups, figures should be used consistently, e.g. 5 to 10, not five to 10.

Spelling should follow the *Concise Oxford Dictionary*.

Systematic papers must conform with the International Code of Botanical and Zoological Nomenclature and, as far as possible, with their recommendations.

Synonymies should be given in the short form (taxon, author, date, page) and the full reference cited at the end of the paper.

Manuscripts:

The original and two copies of manuscripts and figures should be submitted to the Editorial Committee, c/- Publications Department, Western Australian Museum, Francis Street, Perth, Western Australia 6000. They must be in double-spaced typescript on A4 sheets. All margins should be at least 30 mm wide. Tables plus heading and legends to illustrations should be typed on separate pages. The desired position for insertion of tables and illustrations in the text should be indicated in pencil. Tables should be numbered consecutively, have headings which make them understandable without reference to the text, and be referred to in the text.

High quality illustrations are required to size (13.5 cm x 18 cm) or no larger than 32 cm x 40 cm with sans serif lettering suitable for reduction to size. Photographs must be good quality black and white prints, not exceeding 13 cm x 18 cm (5 inches x 7 inches). If scale line and lettering are required on photographs *do not* place directly on to print. They should be positioned on a clear paper or film overlay. Scale must be indicated on illustrations. All maps, line drawings, photographs and graphs, should be numbered in sequence and referred to as Figure/s in the text and captions. Each must have a brief, fully explanatory caption. On acceptance an IBM compatible disk containing all corrections should be sent with amended manuscript. The disk should be marked with program (e.g. Wordperfect, Wordstar, etc) and exact catchline used.

In papers dealing with historical subjects references may be cited as footnotes. In all other papers references must be cited in the text by author and date and all must be listed alphabetically at the end of the paper. The names of journals are abbreviated according to *World List of Scientific Periodicals*. The use of 'unpublished data' or 'personal communication' is discouraged.

Processing:

Papers and short communications are reviewed by at least two referees and acceptance or rejection is then decided by the editors.

The senior author is sent one set of page proofs which must be returned promptly.

The senior author will receive fifty free offprints of the paper. Additional offprints can be ordered at page proof stage.

CONTENTS

G.B. Trotter The Martini-Enfield rifle in Western Australia. Part I. The Western Australian Pattern Martini-Enfield	283
G.B. Trotter The Martini-Enfield rifle in Western Australia. Part II. The procurement of .303 inch calibre Martini Arms by Western Australia	297
G.B. Trotter The Martini-Enfield rifle in Western Australia. Part III. The Pattern 1888 Sword Bayonet in colonial W.A.	309
J.D. Roberts, C.M. Majors Range extensions, range definitions and call structures for frogs from Western Australia	315
C.L. Deeleman-Reinhold Description of a new cave-dwelling pholcid spider from north-western Australia, with an identification key to the genera of Australian Pholcidae (Araneae)	323
K.J. McNamara First record of a spatangoid echinoid from the Paleocene of the Perth Basin, Western Australia	331
B. Saeed, W. Ivanstoft, L.E.L.M. Crowley A new species of the surf-inhabiting atheriniform <i>Iso</i> (Pisces: Isonidae)	337
B. Baehr, M. Baehr New species and new records of Hersiliidae from Australia, with an updated key to all Australian species (Arachnida, Araneae, Hersiliidae)	347
H. Hoch A new troglitic planthopper species (Hemiptera: Fulgoroidea: Meenoplidae) from Western Australia	393
D.J. Kitchener, W.C. Packer, I. Maryanto Taxonomic status of <i>Nyctimene</i> (Chiroptera: Pteropodidae) from the Banda, Kai and Aru Islands, Maluku, Indonesia – implications for biogeography	399
G.M. Shea, M. Peterson Notes on the biology of the genus <i>Pletholax</i> Cope (Squamata: Pygopodidae)	419
D.J. Kitchener, I. Maryanto A new species of <i>Melomys</i> (Rodentia: Muridae) from Kai Besar Island, Maluku Tengah, Indonesia	427
Yu Wen Early Ordovician gastropods from the Canning Basin, Western Australia	437
SHORT COMMUNICATION	
G.R. Allen <i>Variichthys</i> , a replacement name for the terapontid fish genus <i>Varia</i> and first record of <i>V. lacustris</i> from Australia	459